

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 February 2002 (14.02.2002)

PCT

(10) International Publication Number
WO 02/12263 A1

(51) International Patent Classification⁷: **C07H 19/16**,
21/00, C12Q 1/68

(21) International Application Number: PCT/EP01/08850

(22) International Filing Date: 31 July 2001 (31.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
00116816.0 3 August 2000 (03.08.2000) EP
01109438.0 24 April 2001 (24.04.2001) EP

(71) Applicant (for DE only): **ROCHE DIAGNOSTICS GMBH** [DE/DE]; Sandhofer Strasse 116, 68305 Mannheim (DE).

(71) Applicant (for all designated States except DE, US): **F.HOFFMANN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse 124, CH-4070 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SEELA, Frank** [DE/DE]; Gutenbergstrasse 44, 49076 Osnabrueck (DE). **BERGMANN, Frank** [DE/DE]; Faltergatter 5, 82393 Ifeldorf (DE). **VON DER ELTZ, Herbert** [DE/DE]; In der

Au 21, 82362 Weilheim (DE). **HEINDL, Dieter** [DE/DE]; Waldschmidtstrasse 1, 82327 Tutzing (DE). **SEIDEL, Christoph** [DE/DE]; Ammerstrasse 39, 82362 Weilheim (DE). **BECHER, Georg** [DE/DE]; Spindelstrasse 24, 49080 Osnabrueck (DE).

(81) Designated States (national): AU, CA, JP, US.

(84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).

Declarations under Rule 4.17:

- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AU, CA, JP, European patent (AT, BE, CH, CY, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR)
- of inventorship (Rule 4.17(iv)) for US only
- of inventorship (Rule 4.17(iv)) for US only
- of inventorship (Rule 4.17(iv)) for US only

Published:

- with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/12263 A1

(54) Title: NUCLEIC ACID BINDING COMPOUNDS CONTAINING PYRAZOLO[3,4-D]PYRIMIDINE ANALOGUES OF PURIN-2,6-DIAMINE AND THEIR USES

(57) Abstract: The present invention is in the field of nucleic acid binding compounds comprising 7-substituted 7-deaza-8-aza-2,6-diamino-purine bases, compounds useful for the preparation of such compounds, various uses thereof and methods for the determination of nucleic acids using said compounds in the field of diagnostics.

Nucleic acid binding compounds containing pyrazolo[3,4-d]pyrimidine analogues of purin-2,6-diamine and their uses

5

The present invention is in the field of nucleic acid binding compounds comprising 7-substituted 7-deaza-8-aza-2,6-diamino-purine bases, compounds useful for the preparation of such compounds, various uses thereof and methods for the determination of nucleic acids using said compounds in the field of diagnostics.

10

Background of the Invention

Nucleic acids have been found to be useful analytes for the determination of the presence or absence of genes or microorganisms in human body fluids, food or environment in the field of health care. Nucleic acid analysis has found widespread use after the introduction of nucleic acid amplification, like the Polymerase Chain Reaction (PCR, see US-A-4,683,202). Thus, sufficient amounts of nucleic acids are available from each sample. The nucleic acids can be determined from this pretreated sample using a variety of different techniques, dependent on the particular purpose. Most assays require the use of a probe which is either immobilized or immobilizable or is labelled by attachment of one or more reporter groups.

20

A reporter group has the characteristics to be itself capable to be determined or it can be reacted with reagents that make the probe determinable via said reporter group. Thus, for example, probes that are labelled by reporter groups can be determined, as can be hybrids that contain the probe and a nucleic acid to be determined. In case of immobilized probes, the hybrid between the probe and the nucleic acid to be determined is determined at the solid phase to which the probe is bound. In a particular form of assays, not only one nucleic acid having a specific sequence, but a large number of nucleic acids of different sequence is determined. For this purpose, the probes are immobilized in tiny spots in an array on a flat surface such as a glass chip (EP-A-0 476 014 and TIBTECH (1997), Vol. 15, 465-469, WO89/10977, WO89/11548, US 5,202,231, US5,002,867, WO 93/17126). Further development has provided methods for making very large arrays of oligonucleotide probes in very small areas. (US 5,143,854, WO 90/15070, WO 92/10092). Microfabricated arrays of large numbers of oligonucleotide probes, called "DNA chips" offer great promise for a wide variety of applications (see e.g. US 6,156,501 and US 6,022,963).

35

However, nucleic acid determinations often suffer from the problem that the base pairing possibilities between the natural bases A and T and C and G have different stability. This can be attributed to the different capability of these bases to form hydrogen bonding. Thus,

the dA-dT-base pair has two hydrogen bridges, while the dG-dC-base pair has three hydrogen bridges. This results in different melting temperatures (T_m) of hybrids, depending on the GC content [1 - 3]. The higher the GC content, the higher the T_m . The hybridisation strength or the degree of hybridization may be investigated by the measurement of the T_m of the resulting duplex. This can be done by exposing a duplex in solution to gradually increasing temperature and monitoring the denaturation of the duplex, for example, by absorbance of ultraviolet light, which increases with the unstacking of base pairs that accompanies denaturation. The T_m is generally defined as the temperature midpoint of the transition from a fully duplex structure to complete denaturation, i.e. the formation of two isolated single strands.


Therefore in routine nucleic acid analysis, there is often the wish to change the T_m of a nucleic acid molecule. For example, for certain purposes it may be advantageous to equalize or harmonize the T_m of nucleic acids of the same length or to make it even independent from the length of the nucleic acid or the binding region in order to be in the position to apply similar hybridization conditions for all assays. This is particularly necessary for assays using arrays, as on such arrays the hybridizing conditions for each probe must be identical. One solution was the use of low hybridization temperatures. Under such conditions, many nucleic acids having a low degree of base sequence complementarity will bind to the probe. This is called unspecific binding which does not allow discrimination between similar sequences. Another proposal was directed to the use of chemical reagents in the hybridization mixture, for example the addition of tetramethylammonium chloride (TMAC). This reagent reduces the difference between the stability of dG-dC and dA-dT base pairs but the effect is insufficient for short oligonucleotides. Further the addition of salts such as TMAC may not be appreciated as it complicates the optimization of the assay. Another proposal was directed to the use of different concentrations of each different (immobilized) probe in one assay. This was found to be technically complex if not impossible on a chip surface. As a further option the substitution of ribonucleotides in an oligonucleotide composed of deoxyribonucleotides, and vice versa was applied for the adaptation of DNA stability, Hoheisel (1996), Nucleic Acids Res. 24, 430-432.

However, it may be also advantageous to increase the T_m of a given nucleic acid. This is interesting in the field of nucleic acids used for antisense therapy, mismatch discrimination and for nucleic acids used in diagnostics. The nucleic acids may be used as primers or probes. The aim is to allow a more simple design of primers and probes used in multiplex

- reactions and to synthesize shorter capture probes used on chips, as the chemical synthesis of oligonucleotides on a chip surface used for arrays is not as effective as in routine oligonucleotide synthesis. The relative contribution of each base pair to the melting temperature of a hybrid is the higher the shorter an oligonucleotide is. In consequence, the difference in stability between a mismatch and a perfect match is higher for shorter oligonucleotides. However, short oligonucleotides hybridize weakly and, therefore, the hybridization reaction has to be performed at low stringency. In consequence, the potential higher ability of discrimination between different sequences by shorter oligonucleotides can only be used under conditions of low stringency. It would be of considerable advantage to provide bases which allow to achieve a high level of mismatch discrimination under stringent conditions, in particular for short oligonucleotides at temperatures used e.g. in amplification reactions. Further, there is the desire in the state of the art to use short oligonucleotides with high discriminatory power in arrays as the chemical synthesis of oligonucleotides on solid supports used for arrays is not as effective as in routine synthesis. Therefore, the ability to use shorter oligonucleotides under stringent conditions would be of considerable advantage. If bases are found that lead to an increase of the T_m of an oligonucleotide hybridized to its complementary strand, other bases may then be used in the same oligonucleotide to further adjust the T_m according to the preferences of the test system to be used.
- Theoretically, oligonucleotide duplexes forming other tridentate base pairs should exhibit a similar or higher stability, e.g. those with 2-aminoadenine opposite to thymine. Nevertheless, it has been shown that 2-aminoadenine-thymine/uracil base pairs exhibit only a low thermal stability [4 - 10]. From the data published so far one can conclude that the additional NH_2 -group of 2'-deoxyadenosin-2-amine (molecule 1 (see below); n^2A_d) contributes very little to the base pair stability of a DNA duplex. The T_m -increase is in the range of only 1-2° C. Furthermore, this stabilization does not correspond to the total number of n^2A_d -residues incorporated in the duplex instead of dA [11]. A stronger stabilization as reported for duplex DNA is found for duplex RNA or for DNA-RNA hybrids [9] [10] [12]. A rather high base pair stability is observed when 2-aminoadenine is introduced into PNA [13] or hexitol nucleic acids [14]. Modified backbones other than of DNA or of RNA appear to enhance the stability of the 2-aminoadenine-thymine/uracil pair.
- The unusual behavior of oligonucleotide duplexes containing n^2A_d -dT residues is interesting for the development of an adenine-thymine recognition motif showing the same


or even a higher stability than a guanine-cytosine base pair. In the following compounds the purine moiety of compound 1 is replaced by an 8-aza-7-deazapurine (pyrazolo[3,4-d]pyrimidine) or a 7-deazapurine (pyrrolo[2,3-d]pyrimidine) leading to nucleosides (2a [15], 2b, 2c or 3 [16], [17] see below).

5




1

purine numbering



2a: R = H
2b: R = Br
2c: R = I

systematic numbering



3

systematic numbering

Compounds of similar chemical architecture were investigated in the prior art. The synthesis of 7-substituted-7-deaza and 8-aza-7-deazapurine 2'-deoxyribonucleotides, their incorporation into oligonucleotides, and the stability of the corresponding duplexes has been investigated (Seela et al. (1997) *Nucleosides & Nucleotides* 16, 963-966). This document does not contain a disclosure of 7-substituted 7-deaza-8-aza-diamino-purines. Stabilization of duplexes by pyrazolopyrimidine base analogues have been reported (Seela et al. (1988) *Helv. Chim. Acta* 71, 1191-1198; Seela et al. (1988) *Helv. Chim Acta* 71, 1813-1823; and Seela et al. (1989) *Nucleic Acids Res.* 17, 901-910)

Pyrazolo[3,4-d]pyrimidine residues in oligonucleotides are also useful as sites of attachment of various groups (WO90/14353). Oligonucleotides having incorporated one or more pyrazolo[3,4-d]pyrimidine have an enhanced tendency to form triplexes (Belousov et al. (1998). *Nucleic Acids Res.* 26, 1324-1328).

The compounds 7-iodo, 7-cyano and 7-propynyl-7-deaza-2-amino-2'-deoxyadenosine were synthesized by Balow et al. (1997, *Nucleosides & Nucleotides* 16, 941-944) and incorporated into oligonucleotide sequences. These oligonucleotides exhibit enhanced binding affinities to RNA complements relative to unmodified sequences. However, no

corresponding 8-aza-compounds were made and investigated. Seela et al. (1999, Nucleosides & Nucleotides 18, 1399-1400) disclose 7-substituted 8-aza-7-deazapurine DNA, its synthesis and duplex stability. The authors do not address possible uses of the disclosed compounds.

5

WO 90/03370 discloses 3,4-disubstituted and 3,4,6-trisubstituted pyrazolo-[3,4-d]-pyrimidines, more particularly 4,6-diamino-pyrazolo-[3,4-d]-pyrimidines with a linker at the C3-position to which an intercalator, an electrophilic cross linker or a reporter group is attached. These compounds may be attached to sugars or incorporated into
10 oligonucleotides and thereby used for the identification, isolation, localization and/ or detection of complementary nucleic acid sequences of interest. US 5,594,121 discloses novel oligomers with enhanced abilities to form duplexes or triplexes. The oligomers may contain 7-substituted 8-aza-7-deaza-diamino-purines with propinyl and aryls as substituents at the 7-position. Compositions containing these oligomers may used for
15 diagnostic purposes.

There is still a need to provide probes with a high discriminatory power and with a short length, the T_m of which is high under stringent conditions and which can be used in various methods useful in the field of diagnostics as e.g. in the Lightcycler® system (Roche,
20 Mannheim, Germany), TaqMan® (WO92/02638 and corresponding US patents US 5,210,015, US 5,804,375, US 5,487,972) or other applications involving fluorescence energy transfer.

Terms and definitions

25 Conventional techniques of molecular biology and nucleic acid chemistry, which are within the skill of the art, are fully explained fully in the literature. See, for example, Sambrook et al., 1989, Molecular Cloning--A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York; Oligonucleotide Synthesis (M. J. Gait, ed., 1984); Nucleic Acid Hybridization (B. D. Hames and S. J. Higgins. eds., 1984); and a series, Methods in
30 Enzymology (Academic Press, Inc.), all of which are incorporated herein by reference. All patents, patent applications, and publications mentioned herein, both supra and infra, are incorporated herein by reference

The terms "nucleic acid" and "oligonucleotide" refer to polydeoxyribonucleotides
35 (containing 2-deoxy-D-ribose), to polyribonucleotides (containing D-ribose), and to any other type of polynucleotide which is an N glycoside of a purine or pyrimidine base, or

modified purine or pyrimidine base. There is no intended distinction in length between the terms "nucleic acid" and "oligonucleotide", and these terms will be used interchangeably. These terms refer only to the primary structure of the molecule. Thus, these terms include double- and single-stranded DNA, as well as double- and single-stranded RNA. The term
5 "polynucleotide" shall be used interchangeably for "nucleic acid".

The term „backbone“ or "nucleic acid backbone" for a nucleic acid binding compound according to the invention refers to the structure of the chemical moiety linking nucleobases in a nucleic acid binding compound. The bases are attached to the backbone
10 and take part in base pairing to a complementary nucleic acid binding compound via hydrogen bonds. This may include structures formed from any and all means of chemically linking nucleotides, e.g. the natural occurring phosphodiester ribose backbone or unnatural linkages as e.g. phosphorthioates, methyl phosphonates, phosphoramidates and phosphotriesters. Peptide nucleic acids have unnatural linkages. Therefore, a „modified
15 backbone“ as used herein includes modifications to the chemical linkage between nucleotides as described above, as well as other modifications that may be used to enhance stability and affinity, such as modifications to the sugar structure. For example an α -anomer of deoxyribose may be used, where the base is inverted with respect to the natural β -anomer. In an embodiment, the 2'-OH of the sugar group may be altered to 2'-O-alkyl or
20 2'-O-alkyl-n(O-alkyl), which provides resistance to degradation without comprising affinity. An unmodified nucleotide sequence having a phosphodiester backbone is "comparable" to a nucleobase-containing sequence having a modified backbone if the two sequences have identical base sequencing. Thus, the backbones of such sequences are also comparable.

25 The term „nucleic acid binding compound" refers to substances which associate with nucleic acids of any sequence and are able to function as binding partner to a substantially complementary nucleic acid. The binding preferably occurs via hydrogen bonding between complementary base pairs when the nucleic acid binding compound is in a single-stranded
30 form. Preferably, non-natural bases, the subject of the invention, attached to the backbone of the nucleic acid binding compound may be also involved in hydrogen-bonding, however, these may also be able to form hydrogen bonds to only some or all natural occurring bases as e.g. inosine. The expert in the field recognizes that the most well-known „nucleic acid binding compounds" are nucleic acids as DNA or RNA.

35

- The term „probe“ refers to synthetic or biologically produced nucleic acids (DNA or RNA) which, by design or selection, contain specific nucleotide sequences that allow them to hybridize under defined predetermined stringencies, specifically (i.e., preferentially) to target nucleic acids. A „probe“ can be identified as a „capture probe“ meaning that it
- 5 "captures" the target nucleic acid so that it can be separated from undesirable materials which might obscure its detection. Once separation is accomplished, detection of the captured target nucleic acid can be achieved using a suitable procedure. „Capture probes“ are often already attached to a solid phase.
- 10 The term "hybridization" refers the formation of a duplex structure by two single-stranded nucleic acids due to complementary base pairing. Hybridization can occur between fully complementary nucleic acid strands or between "substantially complementary" nucleic acid strands that contain minor regions of mismatch. Conditions under which only fully complementary nucleic acid strands will hybridize are referred to as "stringent
- 15 hybridization conditions" or "sequence-specific hybridization conditions". Stable duplexes of substantially complementary sequences can be achieved under less stringent hybridization conditions; the degree of mismatch tolerated can be controlled by suitable adjustment of the hybridization conditions. Those skilled in the art of nucleic acid technology can determine duplex stability empirically considering a number of variables
- 20 including, for example, the length and base pair concentration of the oligonucleotides, ionic strength, and incidence of mismatched base pairs, following the guidance provided by the art (see, e.g., Sambrook et al., 1989, Molecular Cloning--A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.; and Wetmur, 1991, Critical Review in Biochem. and Mol. Biol. 26(3/4):227-259; both incorporated herein by reference).
- 25 The term "primer" refers to an oligonucleotide capable of acting as a point of initiation of DNA synthesis under conditions in which synthesis of a primer extension product complementary to a nucleic acid strand is induced, i.e., either in the presence of four different nucleoside triphosphates and an agent for extension (e.g., a DNA polymerase or
- 30 reverse transcriptase) in an appropriate buffer and at a suitable temperature. As used herein, the term "primer" is intended to encompass the oligonucleotides used in ligation-mediated amplification processes, in which one oligonucleotide is "extended" by ligation to a second oligonucleotide which hybridizes at an adjacent position. Thus, the term "primer extension", as used herein, refers to both the polymerization of individual nucleoside
- 35 triphosphates using the primer as a point of initiation of DNA synthesis and to the ligation of two primers to form an extended product. A primer is preferably a single-stranded DNA.

The appropriate length of a primer depends on the intended use of the primer but typically ranges from 6 to 50 nucleotides. Short primer molecules generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. A primer need not reflect the exact sequence of the template nucleic acid, but must be sufficiently
5 complementary to hybridize with the template. The design of suitable primers for the amplification of a given target sequence is well known in the art and described in the literature cited herein. Primers can incorporate additional features which allow for the detection or immobilization of the primer but do not alter the basic property of the primer, that of acting as a point of initiation of DNA synthesis. For example, primers may contain
10 an additional nucleic acid sequence at the 5' end which does not hybridize to the target nucleic acid, but which facilitates cloning of the amplified product. The region of the primer which is sufficiently complementary to the template to hybridize is referred to herein as the hybridizing region.

15 The terms "target", "target sequence", "target segment", "target region", and "target nucleic acid" refer to a region or subsequence of a nucleic acid which is to be amplified or investigated.

As used herein, a primer is "specific" for a target sequence if the number of mismatches
20 present between the primer sequence and the target sequence is less than the number of mismatches present between the primer sequence and non-target sequences which may be present in the sample. Hybridization conditions can be chosen under which stable duplexes are formed only if the number of mismatches present is no more than the number of mismatches present between the primer sequence and the target sequence. Under such
25 conditions, the primer can form a stable duplex only with a target sequence. Thus, the use of target-specific primers under suitably stringent amplification conditions enables the specific amplification of those target sequences which contain the target primer binding sites. The use of sequence-specific amplification conditions enables the specific
amplification of those target sequences which contain the exactly complementary primer
30 binding sites.

Halogen means a fluoro, chloro, bromo or iodo group. The most preferred halogen groups are -I and -Br.

35 Alkyl groups are preferably chosen from alkyl groups containing from 1 to 10 carbon atoms, either arranged in linear, branched or cyclic form. The actual length of the alkyl

group will depend on the steric situation at the specific position where the alkyl group is located. If there are steric constraints, the alkyl group will generally be smaller, the methyl and ethyl group being most preferred. All alkyl, alkenyl and alkynyl groups can be either unsubstituted or substituted. Substitution by hetero atoms as outlined above, will help to
5 increase solubility in aqueous solutions.

Alkenyl groups are preferably selected from alkenyl groups containing from 2 to 10 carbon atoms. For the selections similar considerations apply as for alkyl groups. They also can be linear, branched and cyclic. The most preferred alkenyl group is the ethylene group.

10 Alkynyl groups have preferably from 2 to 10 carbon atoms. Again, those carbon atoms can be arranged in linear, branched and cyclic manner. Further, there can be more than one triple bond in the alkynyl group. The most preferred alkynyl group is the 3-propargyl-group.

15 Alkoxy groups preferably contain from 1 to 6 carbon atoms and are attached to the rest of the moiety via the oxygen atom. For the alkyl group contained in the alkoxy groups, the same considerations apply as for alkyl groups. The most preferred alkoxy group is the methoxy group.

20 By "aryl" and "heteroaryl" (or "heteroaromatic") is meant a carbocyclic or heterocyclic group comprising at least one ring having physical and chemical properties resembling compounds such as an aromatic group of from 5 to 6 ring atoms and comprising 4 to 20 carbon atoms, usually 4 to 9 or 4 to 12 carbon atoms, in which one to three ring atoms is N,
25 S or O, provided that no adjacent ring atoms are O--O, S--S, O--S or S--O. Aryl and heteroaryl groups include, phenyl, 2-, 4- and 5-pyrimidinyl, 2-, 4- and 5-thiazoyl, 2-s-triazinyl, 2-, 4-imidazolyl, 2-, 4- and 5-oxazolyl, 2-, 3- and 4-pyridyl, 2- and 3-thienyl, 2- and 3-furanyl, 2- and 3-pyrrolyl optionally substituted preferably on a ring C by oxygen, alkyl of 1-4 carbon atoms or halogen. Heteroaryl groups also include optional substitution
30 on a ring N by alkyl of 1-4 carbon atoms or haloalkyl of 1-4 carbon atoms and 1-4 halogen atoms. Exemplary substituents on the aryl or heteroaryl group include methyl, ethyl, trifluoromethyl and bromo. Such substituted aryl and heteroaryl groups include benzyl and the like. "Heteroaryl" also means systems having two or more rings, including bicyclic moieties such as benzimidazole, benzotriazole, benzoxazole, and indole. Aryl groups are
35 the phenyl or naphthyl moiety, either unsubstituted or substituted by one or more of amino, -cyano, -aminoalkyl, -O-(C₁-C₁₀)-alkyl, -S-(C₁-C₁₀)-alkyl, -(C₁-C₁₀)-alkyl, sulfonyl,

- 10 -

sulfenyl, sulfinyl, nitro and nitroso. Most preferred aryl group is the phenyl group. Preferred arylalkyl group is the benzyl group. The preferred alkylamino group is the ethylamino group. The preferred -COO(C₁-C₄) alkyl group contains one or two carbon atoms in the alkyl moiety (methyl or ethyl esters). Other aryl groups are heteroaryl groups as e.g. pyrimidine, purine, pyrrol, or pyrazole. Aryl and heteroaryl. According to the present invention the term aryl shall also include all heteroaryls.

Aryloxy groups preferably contain from 6 to 20 carbon atoms. Those carbon atoms may be contained in one or more aromatic rings and further in side chains (for example, alkyl chains) attached to the aromatic moiety. Preferred aryloxy groups are the phenoxy and the benzyloxy group.

A "protecting group" is a chemical group that is attached to a functional moiety (for example to the oxygen in a hydroxyl group or the nitrogen in an amino group, replacing the hydrogen) to protect the functional group from reacting in an undesired way. A protecting group is further defined by the fact that it can be removed without destroying the biological activity of the molecule formed, here the binding of the nucleic acid binding compound to a nucleic acid. Suitable protecting groups are known to a man skilled in the art. Especially preferred protecting groups for example for hydroxyl groups at the 5'-end of a nucleotide or oligonucleotide are selected from the trityl groups, for example dimethoxytrityl. Preferred protecting groups at exocyclic amino groups in formula I are acyl groups, most preferred the benzoyl group (Bz), phenoxyacetyl or acetyl or formyl, and the amidine protecting groups as e.g. the N,N-dialkylformamidine group, preferentially the dimethyl-, diisobutyl-, diisobutyl and the di-n-butylformamidine group. Preferred O-protecting groups are the aroyl groups, the diphenylcarbamoyl group, the acyl groups, and the silyl groups. Among these most preferred is the benzoyl group. Preferred silyl groups are the trialkylsilyl groups, like, trimethylsilyl, triethylsilyl and tertiary butyl-dimethyl-silyl. Another preferred silyl group is the trimethylsilyl-oxy-methyl group (TOM)(Swiss Patent Application 01931/97). Further, preferred protecting groups are groups as ortho nitro-benzyl protecting groups like 2-(4-nitrophenyl)ethoxycarbonyl (NPEOC) or photoactivable compounds as 2-nitrophenylpropyloxycarbonyl (NPPOC) (Giegrich et al., Nucleosides & Nucleotides 1998, 17, 1987). . According to the invention, also the phthaloyl group may be used as protecting group.

Any atom in the definitions within the formulae presented herein is not limited to a specific isotope. Thus, a phosphorous atom (P) can either mean the regular ³¹P or the radioactive

³²P or a mixture thereof. The same applies for hydrogen (H/D/T), carbon (C), iodine (Cl, Br, I) and nitrogen (N).

During chemical synthesis, any reactive groups as e.g. -OH, -SH, -NH₂, -NH-alkyl, -NH-alkenylene, -NH-alkynylene, or -NH-aryl (including those groups in reporter groups) should be protected by suitable protecting groups, i.e. that the present invention contemplates compounds for the synthesis of oligonucleotides wherein the formulas or substituents are chosen with the proviso that one or two hydrogen atoms of any -OH, -SH, -NH₂, -NH-alkyl, -NH-alkenylene, -NH-alkynylene, or -NH-aryl group are substituted by a protecting group. Further, during chemical synthesis, the compound will be attached for convenience to a solid phase. In these cases, the definitions of the substituents given above will be selected accordingly.

Reporter groups are generally groups that make the nucleic acid binding compound as well as any nucleic acids bound thereto distinguishable from the remainder of the liquid, i.e. the sample (nucleic acid binding compounds having attached a reporter group can also be termed labeled nucleic acid binding compounds, labeled probes or just probes). The term reporter group and the specific embodiments preferably include a linker which is used to connect the moiety intended to be used (the actual solid phase or the fluorophoric moiety) to the position of attachment as the reporter group. The linker will provide flexibility such that the nucleic acid binding compound can bind the nucleic acid sequence to be determined without major hindrance by the solid phase. Linkers, especially those that are not hydrophobic, for example based on consecutive ethylenoxy units, for example as disclosed in DE 3943522 are known to a man skilled in the art.

By "array" is meant an arrangement of addressable locations on a device. The locations can be arranged in two dimensional arrays, three dimensional arrays, or other matrix formats. The number of locations can range from several to at least hundreds of thousands. Most importantly, each location represents a totally independent reaction site. Each location carries a nucleic acid binding compound which can serve as a binding partner for a second nucleic acid binding compound, a nucleic acid, in particular a target nucleic acid.

The term "building block" or "subunit" refers to a compound which can be used in oligonucleotide synthesis wherein subsequently single building blocks are chemically linked to form a more complex structure, i.e. an oligonucleotide precursor. Examples for building blocks are phosphoramidites or phosphonates.

The term "substituted compound" shall mean that a compound carries further chemical groups, moieties or substituents other than the compound itself. These substituents shall in principle include but are not limited to halogens or alkyl, alkenyl, alkynyl, or aryl
 5 compounds optionally substituted with further heteroatoms

Figure legends

Figure 1a: Ultraviolet spectra of demination of compound 2a

Figure 1b: High pressure liquid chromatography (HPLC) profile of compounds 1, 2a and
 10 2b

Figure 2a: High pressure liquid chromatography (HPLC) profile of a snake venom diesterase/ alkaline phosphatase hydrolysate of an oligonucleotide containing 2a

Figure 2b: High pressure liquid chromatography (HPLC) profile of a snake venom diesterase/ alkaline phosphatase hydrolysate of an oligonucleotide containing 2b

15 Figure 3: Circular dichroism spectra of antiparallel DNA/DNA duplexes (Figure 3a) and antiparallel DNA/ RNA duplexes (Figure 3b)

Figure 4: Circular dichroism spectra of parallel DNA/DNA duplexes (Figure 4a) and parallel DNA/ RNA duplexes (Figure 4b)

Figure 5. HPLC profiles of the nucleosides 1, 2a-c (a) and 3a-c and dT (b). The mixtures of
 20 the nucleosides were analyzed on reversed-phase HPLC at 260 nm (RP-18, gradient: 0-30 min in 0.1M (Et₃NH)OAc (pH7.0) / MeCN (95:5), 0.7 ml/min). HPLC profile of oligonucleotides 100 (c) and 304 (d) after enzymatic hydrolysis with snake-venom phosphodiesterase followed by alkaline phosphatase in 0.1 M Tris-HCl buffer (pH 8.3) at 37°. HPLC was performed on a RP-18 column (200 x 10 mm), 0-30 min in 0.1 M
 25 (Et₃NH)OAc (pH 7.0) / MeCN (95:5), 0.7 ml/min. Peak detection: 260 nm.

Figure 6. The tendency of T_m -increase vs. the number of 2b residues.

Figure 7. (a) CD spectra of duplexes 102 • 103, 108 • 109, 110 • 111, 104 • 105 (in 100 mM NaCl, 100 mM MgCl₂, and 60 mM Na-cacodylate (pH 7.0). (b) CD Spectra of duplexes 102 • 403, 108 • 403, 110 • 403, 104 • 403 in 100 mM NaCl, 100 mM MgCl₂, and

60 mM Na-cacodylate (pH 7.0), the concentration of the oligonucleotides is 5 μ M + 5 μ M (single strand concentration).

Description of the Invention

- 5 The above-mentioned problem could be solved by the findings of the present invention which discloses nucleic acid binding compounds wherein 7-substituted 8-aza-7-deaza-2,6-diamino-purines are incorporated, derivatives of these oligonucleotides and compounds useful for the synthesis thereof. Further, the present invention discloses methods wherein the compounds according to the present invention may be used.
- 10 Surprisingly, an oligonucleotide having incorporated a 7-iodo or 7-bromo-8-aza-7-deaza-2,6-diamino-purine in place of an adenine residue shows an unexpected increase in the melting temperature. This finding is in contrast to the behaviour of 5-halogen substituted pyrimidines. This effect can be observed for parallel as well as for antiparallel duplexes.
- 15 These findings could not be foreseen on the basis of the state of the art, however, the surprising finding can be generalized to other 8-aza-7-deaza-2,6-diamino-purines substituted at the 7-position (purine numbering) with a hydrophobic residue as the halogen substituent is hydrophobic or a residue containing a hydrophobic moiety attached to the 7-position. Further, the teachings can be generalized to an electron-withdrawing
- 20 substituent. Therefore, preferred other substituents at this position as e.g. alkyl, alkenyl and alkynyl, preferably with a length of 6 carbon atoms, more preferably with a length of 3 carbon atoms, are suitable in the present invention. The 7-position allows the substituents to extend into the major groove of the DNA where they find enough space and do not disturb the DNA double helix. A possible explanation for this effect could be removal of
- 25 water from the major groove and the concomitant hydrophobisation of the major groove thereby increasing the stacking of the DNA bases. Substituents fulfilling these requirements may also carry other functional groups as e.g. reporter groups.


- 14 -

In summary, the present invention discloses nucleic acid binding compounds comprising 7-substituted 7-deaza-8-aza-2,6-diamino-purine bases, compounds useful for the preparation of such compounds, various uses thereof and methods for the determination of nucleic acids using said compounds in the field of diagnostics.

5

The subject of the present invention is a nucleic acid binding compound comprising a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic group is a group of the general formula I

Formula I



10 wherein

R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of

- (1) -F, -Cl, -Br or -I,
- (2) Nitro
- 15 (3) Cyano
- (4) $-\text{COO}^-$
- (5) $-(\text{C}_1-\text{C}_{10})$ -alkyl substituted according to (10)
- (6) $-(\text{C}_2-\text{C}_{10})$ -alkenyl substituted according to (10)
- (7) $-(\text{C}_2-\text{C}_{10})$ -alkynyl substituted according to (10)
- 20 (8) $-(\text{C}_6-\text{C}_{22})$ -aryl substituted according to (10)
- (9) $-\text{W}-(\text{C}_1-\text{C}_{10})$ -alkyl, $-\text{W}-(\text{C}_2-\text{C}_{10})$ -alkenyl, $-\text{W}-(\text{C}_2-\text{C}_{10})$ -alkynyl, $-\text{W}-(\text{C}_6-\text{C}_{22})$ -aryl or W-H, wherein W = -S-, -O-, -NH-, -S-S-, -CO-, -COO-, -CO-NH-, -NH-CO-, -NH-CO-NH-, -NH-CS-NH-, $-(\text{CH}_2)_n-[\text{O}-(\text{CH}_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 0 or 1 independently from r and s,
- 25 (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -SH, -NO₂, -CN, -S-(C₁-C₆)-alkyl, $-\text{NR}^5\text{R}^6$, $-\text{N}^+\text{R}^5\text{R}^6\text{R}^{12}$, -OR¹², -COR¹¹, -NH-CO-NR⁵R⁶, -NH-CS-NR⁵R⁶ and $-(\text{CH}_2)_n-[\text{O}-(\text{CH}_2)_r]_s-$ NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,
- 30

- 15 -

wherein R^{11} is selected from the group consisting of $-NHR^{12}$, OR^{12} , and $-SR^{12}$

wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of $-H$, $-(C_1-C_{10})$ -alkyl, $-(C_2-C_{10})$ -alkenyl, $-(C_2-C_{10})$ -alkynyl, $-(C_6-C_{22})$ -aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,

said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being

unsubstituted or substituted by one or more moieties selected from the group consisting of $-halogen$, $-SH$, $-S-(C_1-C_6)$ -alkyl, $-(C_1-C_6)$ -alkoxy, $-OH$, $-NR^5R^6$, $-COR^{11}$, $-NH-CONR^5R^6$, $-NH-CSNR^5R^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ,

with the proviso that R^5 , R^6 or R^{12} is not a reporter group if the radicals (5) to (7) are substituted by $-NR^5R^6$, NHR^{12} , OR^{12} , or SR^{12} ;

R^2 , R^3 is independent from X , R^1 , R^2 and R^3 and is selected from the group of,

- (1) $-H$
- (2) (C_1-C_{10}) -alkyl,
- (3) (C_2-C_{10}) -alkenyl,
- (4) (C_2-C_{10}) -alkynyl,
- (5) (C_6-C_{22}) -aryl,
- (6) $-Z-(C_1-C_{10})$ -alkyl, $-Z-(C_2-C_{10})$ -alkenyl, $-Z-(C_2-C_{10})$ -alkynyl, $-Z-(C_6-C_{22})$ -aryl or $Z-H$, wherein $Z = -CO-$, $-CO-NH-$, $-CS-NH-$, $-(CH_2)_n-[O-(CH_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s ,
- (7) substituents (2) to (6)

wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of $-halogen$, $-NO_2$, $-OR^{12}$, $-CN$, $-SH$, $-S-(C_1-C_6)$ -alkyl, $-NR^5R^6$, $-N^+R^5R^6R^{12}$, $-COR^{11}$, $-NH-CONR^5R^6$, $-NH-CSNR^5R^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ,

wherein R^{11} is selected from the group consisting of $-NHR^{12}$ and OR^{12} ,

- 16 -

wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of -H, $-(C_1-C_{10})$ -alkyl, $-(C_2-C_{10})$ -alkenyl, $-(C_2-C_{10})$ -alkynyl, $-(C_6-C_{22})$ -aryl and a reporter group,

said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, $-S-(C_1-C_6)$ -alkyl, $-(C_1-C_6)$ -alkoxy, -OH, $-NR^5R^6$, $-COR^{11}$, $-NH-CONR^5R^6$, $-NH-CSNR^5R^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s;

X is independent from R^1 , R^2 or R^3 and is selected from the group consisting of N and CH

D is the position of attachment of the group to the rest of the nucleic acid binding compound.

and any salts thereof.

The nucleobases, to which the nucleic acid binding compound according to the invention may bind, can be nucleobases occurring in nature, as e.g. adenosine, guanosine, uridin or cytidin or the 2'-desoxyderivatives thereof, or nucleobases not occurring in nature as e.g. nucleobases with heterocyclic groups according to the invention, heterocyclic groups as pyrrolo-[2,3-d]-pyrimidine or pyrazolo[3,4-d]-pyrimidine or analogues thereof, preferably the said analogues of guanine or adenine or 7-deaza-guanine. Other non-natural heterocyclic groups are known to the person skilled in the art and need not to be mentioned explicitly herein.

In another preferred embodiment, the invention relates to nucleic acid binding compounds with the general formula I and residues R^2 and R^3 as defined above, that have electron-withdrawing or hydrophobic substituents at the 7-position of 7-deaza-8-aza-2,6-diaminopurine. Therefore, in addition to the preferred halogen substituents other preferred substituents at the 7-position are hydrophobic in nature as e.g. alkyl, alkenyl and alkynyl residues. It is, however, sufficient if the first residues extending into the major groove are hydrophobic. In detail, other preferred substituents at the 7-position are the residues

- (1) $-(C_1-C_{10})$ -alkyl substituted according to (5)
- (2) $-(C_2-C_{10})$ -alkenyl substituted according to (5)
- (3) $-(C_2-C_{10})$ -alkynyl substituted according to (5)
- (4) $-(C_6-C_{22})$ -aryl substituted according to (5)

- 17 -

- (5) substituents (1) to (4) wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -SH, -NO₂, -CN, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -OR¹², -COR¹¹, -NH-CO-NR⁵R⁶, -NH-CS-NR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s, wherein R¹¹ is selected from the group consisting of -NHR¹², OR¹², and -SR¹² wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving, said alkyl, alkenyl, alkynyl or aryl in substituents (1) to (4) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s, with the proviso that R⁵, R⁶ or R¹² is not a reporter group if the radicals (1) to (3) are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;

More preferably, the residue (1) has a length of between 1 and 6 carbon atoms, more preferably a length of between 1 to 3 carbon atoms, and the residues (2) and (3) have a length of between 2 and 6 carbon atoms, more preferably a length of between 2 to 3 carbon atoms.

In the most preferred embodiment the heterocyclic group is 7-bromo-7-deaza-8-aza-2,6-diamino-purine or 7-iodo-7-deaza-8-aza-2,6-diamino-purine. The nucleic acid binding compounds according to the invention further possess the advantage of having a very stable glycosidic bond in contrast to 2-amino-adenosine which has an extremely acid labile glycosidic bond and may only be used in oligonucleotide synthesis if specific conditions are used. In a further embodiment of the invention, the halogenides in the 7-position of the 7-deaza-8-aza-2,6-diamino-purine may be substituted by pseudohalogenides as e.g. -SCN or -CN.

The preferred substituents mentioned above are also preferred in the methods and uses according to the present invention.

- 5 The heterocyclic groups of formula I are mainly characterized by the following properties:
- The base is linked to the backbone, preferred to a sugar moiety, via the 9-position (purine numbering).
 - The base contains an aromatic π -electron system which is capable to form stacking interactions with other nucleic acid constituents.
 - 10 - The base contains donor and/ or acceptor site(s) for hydrogen bonding to the natural nucleoside T.

In order to increase the T_m , in a nucleic acid binding compound one or more A in a strand complementary to a T in the nucleic acid to be determined could be replaced by the

15 heterocyclic groups according to the invention. The oligonucleotide would then bind specifically to the target sequence containing T opposite to the bases according to the invention with a binding energy in the order of a G-C base pair but with higher stability than a A-T base pair. This works for antiparallel or parallel duplexes equally well whereby natural A-T base pairs have equal abilities to bind in parallel or antiparallel duplexes but

20 with a lower binding energy in the parallel duplex. However, the heterocyclic group according to the invention, when incorporated into a nucleic acid binding compound, will bind to a T in the opposite strand equally well in a parallel or in an antiparallel duplex. In order to effect antiparallel binding in a duplex, natural G or C bases normally forming G-C base pairs in antiparallel duplexes have to be substituted by non-natural base pairs as e.g.

25 G-iC (isocytosine) or better G-^miC (methylated isocytosine) or C-iG (isoguanosine) as disclosed above. A summary of parallel and antiparallel duplexes can be found in EP 0 624 161 or Seela et al. (Tetrahedron 55 (1999), 9481-9500) and is shown schematically in scheme 11 and 12 in example 4.

30 This general principle of course is not limited, as bases showing the same characteristics in the 6-membered ring would be expected to have the same properties based on the above explanation due to their containing the structure according to the invention. Particularly, the farther the part of the heterocyclic group from the part participating in the base pairing, the more tolerant will the oligomer be over modifications in the chemical structure, for

35 example the attachment of groups to this part of the heterocyclic rings. In the following,

when reference is made to the heterocyclic group of the invention, there is made reference to a heterocyclic group according to general formula I.

5 The present invention also contemplates tautomeric forms and salts of heterocyclic groups of formula I.

The nucleic acid binding compound according to the invention preferably has a length of less than 100 subunits, more preferably of from 10 to 30 subunits. In order to be active as nucleic acid binding compound, the substituents should be chosen such that hydrogen
10 bonds to heterocyclic groups at the nucleic acid to be bound are enabled, preferably by base pairing, for example by Watson-Crick or Hoogsteen base pairing. Compounds in which the substituents do not enable such preferred hydrogen bonding, can be useful as intermediates for the preparation of nucleic acid binding compounds. Preferred nucleic acid binding compounds of the invention are those which are chemically synthesized.

15 The nucleic acid binding compound will be constructed such that it contains a nucleobase sequence which is substantially complementary to the nucleic acid to be determined or the nucleic acid to which it is intended to be bound by base pairing dependent on the formation of a parallel or an antiparallel duplex. As those nucleic acids will usually contain
20 at least once any of the naturally occurring nucleobases Ade, Cyt, Gua and Thy or Ura, the nucleic acid binding compound according to the invention will also contain any of those four bases. However, according to the invention, at least one of the heterocyclic groups is replaced by the heterocyclic base of formula I.

25 If the nucleic acid binding compound is to be used as a probe for the determination of a nucleic acid, or any other identification of the compound or the nucleic acid is intended, any of the substituents are selected such as to contain a reporter group. While as many reporter groups can be attached as useful to label the nucleic acid binding compound sufficiently, it is preferred to attach only a limited number of reporter groups to a single
30 subunit, such that recognition of nucleic acids, affinities to nucleic acids and solubility is not affected such that the probe would not be useful in hybridization assays. In a very preferred case, there will be only from 1 to 4, most preferably 1 or 2 or most preferred only one reporter group in each nucleic acid binding compound. There are formats for the nucleic acid determination which require more than one reporter group attached to the
35 probe. An example for such formats is disclosed in WO92/02638. In this case, one of the reporter groups will be a fluorescence emitter, while the other is a fluorescence quencher.

- 20 -

The reporter group may also be attached to a heterocyclic base which is not according to formula I.

In a preferred embodiment of the invention, R^1 is alkynyl-amino $-C\equiv C-E-NR^5R^6$, alkenyl-amino $-CH=CH-E-NR^5R^6$ and $-E-NR^5R^6$ wherein E is $-[(CH_2)_rF]_s-(CH_2)_r-$ wherein F = O or S and r and s are independently from one another an integer from 1 to 18, wherein R^5 and R^6 are selected independently from the group consisting of -H, $-(C_1-C_{10})$ -alkyl, $-(C_2-C_{10})$ -alkenyl, $-(C_2-C_{10})$ -alkynyl, $-(C_6-C_{22})$ -aryl and a reporter group, wherein R^{11} is selected from the group consisting of $-NHR^{12}$ and OR^{12} , wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of -H, $-(C_1-C_{10})$ -alkyl, $-(C_2-C_{10})$ -alkenyl, $-(C_2-C_{10})$ -alkynyl, $-(C_6-C_{22})$ -aryl and a reporter group, said alkyl, alkenyl, alkynyl or aryl being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, $-S-(C_1-C_6)$ -alkyl, $-(C_1-C_6)$ -alkoxy, -OH, $-NR^5R^6$, $-COR^{11}$, $-NH-CONR^5R^6$, $-NH-CSNR^5R^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s.

Reporter groups are generally groups that make the nucleic acid binding compound as well as any nucleic acids bound thereto distinguishable from the remainder of the liquid, i.e. the sample (nucleic acid binding compounds having attached a reporter group can also be termed labeled nucleic acid binding compounds, labeled probes or just probes). This distinction can be either effected by selecting the reporter group from the group of directly or indirectly detectable groups or from the groups of immobilized or immobilizable groups. Directly detectable groups are for example fluorescent compounds, like fluorescein and its derivatives, like hexachlorofluorescein and hexafluorofluorescein, rhodamines, psoralenes squaraines, porphyrines, fluorescent particles, bioluminescent compounds, like acridinium esters and luminol, or the cyanine dyes, like Cy-5. Examples of such compounds are disclosed in EP 0 680 969. Further, spin labels like TEMPO, electrochemically detectable groups, ferrocene, viologene, heavy metal chelates and electrochemiluminescent labels, like ruthenium bispyridyl complexes, and naphthoquinones, quencherdyes, like dabcyI, and nuclease active complexes, for example of Fe and Cu, are useful detectable groups. Other examples of such compounds are europium complexes. Indirectly detectable groups are groups that can be recognized by another moiety which is directly or indirectly labelled. Examples of such indirect detectable groups are haptens, like digoxigenin or biotin. Digoxigenin for example can be recognized by antibodies against digoxigenin. Those antibodies may either be labelled directly or can be recognized by labelled antibodies directed against the (antidigoxigenin) antibodies.

- 21 -

Formats based on the recognition of digoxigenin are disclosed in EP-B-0 324 474. Biotin can be recognized by avidin and similar compounds, like streptavidin and other biotin binding compounds. Again, those compounds can be labelled directly or indirectly. Further interesting labels are those directly detectable by atomic force microscopy (AFM) or scanning tunneling microscopy (STM). The reporter group can further be a nucleotide sequence which does not interfere with other nucleotide sequences in the sample. The sequence can therefore be specifically recognized by nucleotide containing a complementary sequence. This nucleotide sequence can be labelled directly or indirectly or can be immobilizable or immobilized. A reporter group can further be a solid phase.

Attachment of the nucleic acid binding compound with solid phase can be either directly or indirectly as pointed out above for the detectable group. Examples of such solid phases are latex beads or gold particles. In another embodiment of the invention, a further reporter group attached to the nucleic acid binding compound may be any positively or negatively charged group, preferably a carboxylate group or an ammonium $N^+R^5R^6R^{12}$ with substituents as specified under formula I as described above. These may be attached e.g. via a propargylen linker to the base and enhance the sensitivity of MALDI-TOF mass spectroscopy (MALDI-TOF: matrix-assisted laser desorption/ionization time-of-flight) in the positive or negative mode. The substituents of the ammonium group are preferably introduced into the oligonucleotide via post-labelling, i.e. binding compounds can be postlabeled with reporter groups when a suitable reactive group is introduced during oligonucleotide synthesis, for example, an amino group protected in the oligonucleotide synthesis precursor with a phthaloyl group.


Direct labelling can be effected by covalent coupling of a nucleic acid binding compound to a reactive group on the solid phase, i.e. preferably via a linker. Indirect labelling can be made similar as disclosed above for the detectable groups. Preferably, indirect attachment is non-covalently by biospecific interactions, for example those selected from the group of hapten-antibody, vitamin-receptor and nucleic acid-complementary nucleic acid. Again, those interactions and their use in nucleic acid assays is known to a man skilled in the art.

Solid phases that are useful for immobilization of the probe according to the invention are preferably selected from the group of polystyrene, polyethylene, polypropylene, glass, SiO_2 and TiO_2 . The formats of such solid phases can be selected according to the needs of the instrumentation and format of the assay. For example, a solid phase may assume the form of a bead or a vessel.

- 22 -

The most popular backbone is the naturally occurring sugar phosphate backbone of nucleic acids containing either ribonucleoside subunits (RNA) or deoxyribonucleoside subunits (DNA). Therefore, in a preferred embodiment, the backbone of the nucleic acid binding compound comprises phosphodiester linkages and ribose. In the last years, there were descriptions of nucleic binding compounds that have similar properties like oligonucleotides, but differ in their backbone, which have structures formed from any and all means of chemically linking nucleotides in contrast to the natural occurring phosphodiester ribose backbone. Therefore, it is evident that the invention would still work, even if the backbone of the nucleic acid binding compound is not an oligonucleotide in the strict sense, i.e. it has a modified backbone. The backbone may include e.g. phosphorothioates, methyl phosphonates, phosphoramidates and phosphorotriesters linkages. Peptide nucleic acids also have unnatural linkages. The modifications in the backbone may vary the properties of the nucleic acid binding compound, i.e. it may enhance stability and affinity. Therefore, in a preferred embodiment, the nucleic acid binding compounds are those, wherein the backbone comprises one or more moieties of the general formula II

Formula II




- wherein
- A is selected from the group consisting of O, S, CH₂, N-CO-(C₁-C₁₀)-alkyl,
- L is selected from the group consisting of oxy, sulfanediyl, -CH₂- and -NR²²-,
- T is selected from the group consisting of oxo, thioxo and selenoxo,
- U is selected from the group consisting of -OH, O⁻, -O-reporter group, -SH, -S reporter group -SeH, -(C₁-C₁₀)-alkoxy, (C₁-C₁₀)-alkyl, -(C₆-C₂₂)-aryl, -(C₆-C₁₄)-aryl-(C₁-C₁₀)-alkyl, -NR²³R²⁴, and -(-O-(C₁-C₁₀)-alkyl)_n-R²⁵, wherein n can be any integer between 1 and 6, or wherein -NR²³R²⁴ can together with N be a 5-6-membered heterocyclic ring,

- 23 -

- V is selected from the group consisting of oxy, sulfanediyl, $-\text{CH}_2-$, or $-\text{NR}^{22}-$,
 R^{14} is selected from the group consisting of $-\text{H}$, $-\text{OH}$, $-(\text{C}_1-\text{C}_6)\text{-alkyl}$, $-(\text{C}_1-\text{C}_{10})\text{-alkoxy}$, $-(\text{C}_2-\text{C}_{10})\text{-alkenyloxy}$, $-\text{halogen}$, $-\text{azido}$, $-\text{O-allyl}$, $-\text{O-alkynyl}$, and $-\text{NH}_2$,
 R^{22} is independently selected from the group of $-\text{H}$ and $-(\text{C}_1-\text{C}_{10})\text{-alkyl}$,
 5 R^{23} and R^{24} are independently selected from the group consisting of $-(\text{C}_1-\text{C}_{10})\text{-alkyl}$, $-(\text{C}_1-\text{C}_{20})\text{-aryl}$, $-(\text{C}_6-\text{C}_{14})\text{-aryl}$, $-(\text{C}_1-\text{C}_{10})\text{-alkyl}$, $-(\text{C}_1-\text{C}_6)\text{-alkyl}$, $[\text{NH}(\text{CH}_2)_c]_d\text{-NR}^{26}\text{R}^{27}$ and a reporter group,
 R^{25} is selected from the group consisting of $-\text{H}$, $-\text{OH}$, $-\text{halogen}$, $-\text{amino}$, $-(\text{C}_1-\text{C}_{18})\text{-alkylamino}$, $-\text{COOH}$, $-\text{CONH}_2$ and $-\text{COO}(\text{C}_1-\text{C}_4)\text{-alkyl}$ and a reporter group,
 10 R^{26} and R^{27} are independently selected from the group consisting from $-\text{H}$, $-(\text{C}_1-\text{C}_6)\text{-alkyl}$, and $-(\text{C}_1-\text{C}_4)\text{-alkoxy}$, $-(\text{C}_1-\text{C}_6)\text{-alkyl}$ and a reporter group,
 c is an integer from 2 to 6,
 d is an integer from 0 to 6, and
 15 B is a moiety of formula I

Formula I



wherein

- R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of
 (1) $-\text{F}$, $-\text{Cl}$, $-\text{Br}$ or $-\text{I}$,
 20 (2) Nitro
 (3) Cyano
 (4) $-\text{COO}^-$
 (5) $-(\text{C}_1-\text{C}_{10})\text{-alkyl}$ substituted according to (10)
 (6) $-(\text{C}_2-\text{C}_{10})\text{-alkenyl}$ substituted according to (10)
 25 (7) $-(\text{C}_2-\text{C}_{10})\text{-alkynyl}$ substituted according to (10)
 (8) $-(\text{C}_6-\text{C}_{22})\text{-aryl}$ substituted according to (10)
 (9) $-\text{W}-(\text{C}_1-\text{C}_{10})\text{-alkyl}$, $-\text{W}-(\text{C}_2-\text{C}_{10})\text{-alkenyl}$, $-\text{W}-(\text{C}_2-\text{C}_{10})\text{-alkynyl}$, $-\text{W}-(\text{C}_6-\text{C}_{22})\text{-aryl}$ or W-H , wherein $\text{W} = -\text{S}-$, $-\text{O}-$, $-\text{NH}-$, $-\text{S-S}-$, $-\text{CO}-$, $-\text{COO}-$, $-\text{CO-NH}-$, $-\text{NH-CO}-$, $-\text{NH-CO-NH}-$, $-\text{NH-CS-NH}-$, $-(\text{CH}_2)_n\text{-[O-(CH}_2)_r]_s-$, where r and s are, independently of each other, an integer
 30 between 1 to 18 and n is 0 or 1 independently from r and s,

- 24 -

- (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -SH, -NO₂, -CN, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -OR¹², -COR¹¹, -NH-CO-NR⁵R⁶, -NH-CS-NR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s, wherein R¹¹ is selected from the group consisting of -NHR¹², OR¹², and -SR¹² wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,
- said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s, with the proviso that R⁵, R⁶ or R¹² is not a reporter group if the radicals (5) to (7) are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;
- R², R³ is independent from X, R¹, R² and R³ and is selected from the group of,
- (1) -H
 - (2) (C₁-C₁₀)-alkyl,
 - (3) (C₂-C₁₀)-alkenyl,
 - (4) (C₂-C₁₀)-alkynyl,
 - (5) (C₆-C₂₂)-aryl,
 - (6) -Z-(C₁-C₁₀)-alkyl, -Z-(C₂-C₁₀)-alkenyl, -Z-(C₂-C₁₀)-alkynyl, -Z-(C₆-C₂₂)-aryl or Z-H, wherein Z = -CO-, -CO-NH-, -CS-NH-, -(CH₂)_n-[O-(CH₂)_r]_s-, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s,
 - (7) substituents (2) to (6)
- wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -NO₂, -OR¹², -CN, -SH, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -

- 25 -

- COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s, wherein R¹¹ is selected from the group consisting of -NHR¹² and OR¹²,
- 5 wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group, said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of -
- 10 halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s;
- 15 X is independent from R¹, R² or R³ and is selected from the group consisting of N and CH; and D is the position of attachment of the group to the rest of the nucleic acid binding compound
- 20 or any salts thereof.

The preferred definitions of the groups as defined under formula I apply to formula II and the following formulae, if not indicated otherwise.

- 25 Preferably, in compounds of formula II, R¹⁴ is hydrogen. Preferred definition of L is oxy. Preferred definition of U is -OH and -O-reporter group. Preferred definition of V is oxy. Preferred definition of c is an integer from 2 to 4, and of d an integer from 0 to 2. Compounds of formula II are especially suited to contain the heterocyclic moiety of the invention as an integrated part (preferably not at one of the termini) of the nucleic acid
- 30 binding compound. The group NR²³R²⁴ is preferably selected from the group consisting of dialkylamino groups. In case of this group together with the forming of 5- or 6-membered heterocyclic ring, it assumes preferably the definition of morpholinyl, pyrrolidinyl or piperidinyl.
- 35 In a further preferred embodiment, the sugar configuration is selected from the group consisting of the α-D-, β-D-, α-L- and β-L-configurations, most preferred the compound

- 26 -

contains at least one 2'-deoxy- β -D-erythro-pentofuranosyl moiety or one β -D-ribofuranosyl moiety. In a preferred embodiment of the invention, D is the glycosid C-1 of a sugar moiety of the compound according to the invention.


- 5 In another embodiment of the invention the sugar is in a locked conformation. LNA (Locked Nucleic Acid) is a novel class of nucleic acid analogue. LNA oligomers obey the Watson-Crick base pairing rules and hybridize to complementary oligonucleotides. However, when compared to DNA and other nucleic acid derivatives, LNA provides vastly improved hybridization performance. LNA/DNA or LNA/RNA duplexes are much more
- 10 thermally stable than the similar duplexes formed by DNA or RNA. In fact, LNA has the highest affinity towards complementary DNA and RNA ever to be reported. In general, the thermal stability of a LNA/DNA duplex is increased 3°C to 8°C per modified base in the oligo. Within the fields of general molecular biology and molecular diagnostics, five major fields for the application of LNA have been identified which are capture probes, sample
- 15 preparation, detection of SNP's (Single Nucleotide Polymorphisms), allele specific PCR, and hybridization probes, Molecular Beacons, Padlock probes, Taqman probes (WO92/02638 and corresponding US patents US 5,210,015, US 5,804,375, US 5,487,972) and probes for *in-situ* hybridizations. In most respects, LNA may be handled like DNA. LNA is at least as stable as DNA and is soluble in aqueous buffers. LNA can be ethanol
- 20 precipitated, dried and resuspended, and can be analyzed on gels, HPLC and MALDI-TOF. LNAs are novel nucleic acid analogs that can dramatically increase the performance of not only diagnostic assays that probe and evaluate genetic information but also of antisense and other genetic medicine approaches. These analogs, which can be utilized in most applications just like their natural counterparts, lock the nucleic acid into the most
- 25 productive conformation for hybridization. Hybridization, or complementary docking of genetic probes, is the predominant form of evaluation of genetic information in diagnostics. A broad variety of applications for LNA have been developed including a number of extremely sensitive and specific assays able to detect specific disease-causing single base mutations in an individual's genes. In the detection of SNPs (Single Nucleotide
- 30 Polymorphisms), which are the small variations in our genes, that may cause a predisposition to disease, there are data to show that LNA capture probes of only eight nucleotides in length are able to more effectively discriminate between mutated and wildtype genes in a sample than much longer conventional nucleic acid capture probes. Therefore the invention also contemplates compounds according to the invention wherein
- 35 e.g. at least one atom of the sugar moiety e.g. a carbon or an oxygen atom is connected to at least one other atom of the sugar moiety via at least one bridging moiety containing at least

- 27 -

one atom whereby a conformationally constrained sugar is formed as outlined above. Thereby the sugar is fixed in a locked conformation.

For the synthesis of the compounds according to the invention, the reader is referred to
 5 Chemistry of Nucleosides and Nucleotides Part 1, edited by L. B. Townsend, Plenum Press New York (1988), Chapter 2: Synthesis and Properties of Purine Nucleosides and Nucleotides, page 113-281 or to US patent 5,594,121. However, more information is provided below.

10 Different chemical structures can be used in the backbone of the nucleic acid binding compound. The expert skilled in the field appreciates the fact that the nucleic acid binding compound may also possess a modified 3'-end. Therefore, a preferred subject of the invention is a nucleic acid binding compound as outlined above, wherein the backbone comprises one or more moieties of the general formula III, wherein t is 0 or 1,



15


wherein in the case that t=1,

- A is selected from the group consisting of O, S, CH₂ and N-(C₁-C₆)-alkyl,
 M is selected from the group consisting of oxy, sulfanediyl, -NR²²-, -(C₁-C₁₀)-alkyl-, or -O-(C₁-C₁₀)-alkyl-O-, and -S-(C₁-C₁₀)-alkyl-O- and -NR²²-(C₁-C₆)-alkyl-O-,
 20 R²² is selected from the group of -H, -(C₁-C₁₀)-alkyl, a protecting group and a reporter group,
 R¹⁴ is selected from the group consisting of -H, -OH, -(C₁-C₃)-alkyl, -(C₁-C₆)-allyl, -(C₁-C₁₀)-alkoxy, -(C₂-C₁₀)-alkenyloxy, -(C₂-C₁₀)-alkynyloxy, -(O-CH₂)_n - wherein n may be an integer of from 1 to 18, -halogen, -azido, SH, -(C₁-C₁₀)-alkylmercapto, O-reporter group, O-solid phase and -NH₂,
 25 R¹⁵ is selected from the group consisting of -H, -(C₁-C₆)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₂-C₁₀)-alkyl-carbonyl, -(C₃-C₁₉)-alkenyl-

- 28 -

carbonyl, $-(C_3-C_{19})$ -alkynyl-carbonyl, $-(C_6-C_{14})$ -aryl- (C_1-C_{10}) -alkyl, a solid phase and a group of formula IV

Formula IV




wherein

- 5 T is selected from the group consisting of oxo, thioxo and selenoxo, and
- U is selected from the group consisting of -OH, O⁻, -O-reporter group, -SH, -SeH, $-(C_1-C_{10})$ -alkoxy, $-(C_1-C_{10})$ -alkyl, $-(C_6-C_{22})$ -aryl, $-(C_6-C_{14})$ -aryl- (C_1-C_{10}) -alkyl, $-NR^{23}R^{24}$, and $-(-O-(C_1-C_{10})$ -alkyl)- R^{25} , wherein N can be any integer between 1 and 6, or
- 10 wherein $NR^{23}R^{24}$ can together with N be a 5-6-membered heterocyclic ring,
- R^{23} and R^{24} are independently selected from the group consisting of - (C_1-C_{10}) -alkyl, $-(C_1-C_{20})$ -aryl, $-(C_6-C_{14})$ -aryl- (C_1-C_{10}) -alkyl, $-(C_1-C_6)$ -alkyl- $[NH(CH_2)_c]_d-NR^{26}R^{27}$,
- 15 R^{25} is selected from the group consisting of -H, -OH, -halogen, -amino, $-(C_1-C_{18})$ -alkylamino, -COOH, -CONH₂ and -COO (C_1-C_4) -alkyl,
- R^{26} and R^{27} are independently selected from the group consisting from -H, $-(C_1-C_6)$ -alkyl, and $-(C_1-C_4)$ -alkoxy- (C_1-C_6) -alkyl
- R^{29} is selected from the group consisting of -OR³⁰ and -SR³⁰,
- 20 R^{30} is selected from the group consisting of -H, $-(C_1-C_{10})$ -alkyl, $-(C_2-C_{10})$ -alkenyl, $-(C_6-C_{22})$ -aryl, a protecting group, a solid phase and a reporter group
- c is an integer from 2 to 6,
- d is an integer from 0 to 6, and
- 25 B is a moiety of formula I

- 29 -

Formula I



wherein

R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of

- (1) -F, -Cl, -Br or -I,
 - (2) Nitro
 - (3) Cyano
 - (4) $-\text{COO}^-$
 - (5) $-(\text{C}_1\text{-C}_{10})$ -alkyl substituted according to (10)
 - (6) $-(\text{C}_2\text{-C}_{10})$ -alkenyl substituted according to (10)
 - (7) $-(\text{C}_2\text{-C}_{10})$ -alkynyl substituted according to (10)
 - (8) $-(\text{C}_6\text{-C}_{22})$ -aryl substituted according to (10)
 - (9) $-\text{W}-(\text{C}_1\text{-C}_{10})$ -alkyl, $-\text{W}-(\text{C}_2\text{-C}_{10})$ -alkenyl, $-\text{W}-(\text{C}_2\text{-C}_{10})$ -alkynyl, $-\text{W}-(\text{C}_6\text{-C}_{22})$ -aryl or W-H, wherein W = -S-, -O-, -NH-, -S-S-, -CO-, -COO-, -CO-NH-, -NH-CO-, -NH-CO-NH-, -NH-CS-NH-, $-(\text{CH}_2)_n\text{-[O-(CH}_2)_r\text{]}_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 0 or 1 independently from r and s,
 - (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -SH, $-\text{NO}_2$, -CN, $-\text{S}-(\text{C}_1\text{-C}_6)$ -alkyl, $-\text{NR}^5\text{R}^6$, $-\text{N}^+\text{R}^5\text{R}^6\text{R}^{12}$, $-\text{OR}^{12}$, $-\text{COR}^{11}$, $-\text{NH-CO-NR}^5\text{R}^6$, $-\text{NH-CS-NR}^5\text{R}^6$ and $-(\text{CH}_2)_n\text{-[O-(CH}_2)_r\text{]}_s-$ NR^5R^6 , r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s, wherein R^{11} is selected from the group consisting of $-\text{NHR}^{12}$, OR^{12} , and $-\text{SR}^{12}$
- wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of -H, $-(\text{C}_1\text{-C}_{10})$ -alkyl, $-(\text{C}_2\text{-C}_{10})$ -alkenyl, $-(\text{C}_2\text{-C}_{10})$ -alkynyl, $-(\text{C}_6\text{-C}_{22})$ -aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,

- 30 -

- said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,
- with the proviso that R⁵, R⁶ or R¹² is not a reporter group if the radicals (5) to (7) are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;
- R², R³ is independent from X, R¹, R² and R³ and is selected from the group of,
- (1) -H
 - (2) (C₁-C₁₀)-alkyl,
 - (3) (C₂-C₁₀)-alkenyl,
 - (4) (C₂-C₁₀)-alkynyl,
 - (5) (C₆-C₂₂)-aryl,
 - (6) -Z-(C₁-C₁₀)-alkyl, -Z-(C₂-C₁₀)-alkenyl, -Z-(C₂-C₁₀)-alkynyl, -Z-(C₆-C₂₂)-aryl or Z-H, wherein Z = -CO-, -CO-NH-, -CS-NH-, -(CH₂)_n-[O-(CH₂)_r]_s-, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s,
 - (7) substituents (2) to (6)
- wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -NO₂, -OR¹², -CN, -SH, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,
- wherein R¹¹ is selected from the group consisting of -NHR¹² and OR¹²,
- wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group,
- said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s;

- 31 -

X is independent from R^1 , R^2 or R^3 and is selected from the group consisting of N and CH; and

D is the position of attachment of the group to the rest of the nucleic acid binding compound

5

and wherein in the case that $t=0$, R^{15} is -H,

or any salts thereof.

10

For the definitions and preferences the particulars apply as outlined for the substituents under formulae I and II, if not specified otherwise specifically for formula III. In a preferred embodiment the 3'-end possesses a 2',3'-dideoxyribose, i.e. wherein $t=0$, R^{15} is -H and R^{14} is -H, or an analogue thereof. This is of interest if an enzymatical termination is necessary when the nucleic acid binding compound according to the invention is extended with triphosphate compounds also occurring in nature as the triphosphates of adenosine, guanosine, uridin, cytidine or thymidine or the desoxyderivates of the triphosphates of adenosine, guanosine, cytidine or thymidine. However, the invention also relates to the extension of primers containing only nucleotides occurring in nature or nucleic acid binding compounds according to the invention with triphosphate compounds according to the invention with the general formula VIII.

15

20

Nucleic acid binding compounds, wherein the group of formula I is attached to subunits, for example the nucleotide, at the 3'-terminus of the compound, are useful either as starting compound for the synthesis of longer compounds or/and as end-labeled probes. This group of compounds is especially preferred because the terminal position of probes generally is the most tolerant in view of attachment of chemical moieties.


25

In view of the modifications to the 3'-end of the nucleic acid binding compound, it is evident that also the 5'-end of the nucleic acid binding compound may be modified. Therefore, another preferred subject of the invention is a nucleic acid binding compound as outlined above comprising a backbone moiety of the formula V

30

- 32 -

Formula V




wherein

A is selected from the group consisting of O, S, CH₂ and N-(C₁-C₆)-alkyl,M' is selected from the group consisting of oxy, sulfanediyl, -NR²²-, -(C₁-C₁₀)-alkyl, or -O-(C₁-C₁₀)-alkyl-O-, and -S-(C₁-C₁₀)-alkyl-O- and -NR²²-(C₁-C₆)-alkyl-O-,R²² is selected from the group of -H, a protecting group, a reporter group and -(C₁-C₁₀)-alkyl,R¹⁴ is selected from the group consisting of -H, -OH, -(C₁-C₁₀)-alkoxy, -(C₂-C₁₀)-alkenyloxy, -(C₂-C₁₀)-alkynyloxy, -halogen, azido, -SH, -S-(C₁-C₆)-alkylmercapto, O-reporter group, O-solid phase and NH₂,R¹⁶ is selected from the group consisting of -H, -(C₁-C₈)-alkyl, -(C₂-C₁₈)-alkenyl, -(C₂-C₁₈)-alkynyl, -(C₂-C₁₈)-alkyl-carbonyl, -(C₃-C₁₉)-alkenyl-carbonyl, -(C₃-C₁₉)-alkynyl-carbonyl, -(C₆-C₁₄)-aryl-(C₁-C₈)-alkyl, a protective group or a compound of formula IV

wherein

Formula IV




T is selected from the group consisting of oxo, thioxo and selenoxo,

U is selected from the group consisting of -OH, -SH, -SeH, -(C₁-C₁₀)-alkoxy, -(C₁-C₁₀)-alkyl, -(C₆-C₂₂)-aryl, -(C₆-C₁₄)-aryl-(C₁-C₁₀)-alkyl, -NR²³R²⁴, and -(O-(C₁-C₁₀)-alkyl)-R²⁵, wherein n can be any integer between 1 and 6, wherein NR²³R²⁴ can together with N be a 5-6-membered heterocyclic ring,R²³ and R²⁴ are independently selected from the group consisting of -(C₁-C₁₀)-alkyl, -(C₁-C₂₀)-aryl, -(C₆-C₁₄)-aryl-(C₁-C₁₀)-alkyl, -(C₁-C₆)-alkyl-[NH(CH₂)_c]-NR²⁶R²⁷,R²⁵ is selected from the group consisting of -H, -OH, -halogen, -amino, -(C₁-C₁₈)-alkylamino, -COOH, -CONH₂ and -COO(C₁-C₄)-alkyl,

- 33 -

- R^{26} and R^{27} are independently selected from the group consisting from -H,
 -(C₁-C₆)-alkyl, and -(C₁-C₄)-alkoxy-(C₁-C₆)-alkyl
 R^{29} is selected from the group consisting of -OR³⁰ and -SR³⁰,
 R^{30} is selected from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -
 (C₆-C₂₂)-aryl, a protecting group, a solid phase and a reporter group, and
 c is an integer from 2 to 6,
 d is an integer from 0 to 6, and
 B is a moiety of formula I

Formula I



- wherein
 R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of
 (1) -F, -Cl, -Br or -I,
 (2) Nitro
 (3) Cyano
 (4) -COO⁻
 (5) -(C₁-C₁₀)-alkyl substituted according to (10)
 (6) -(C₂-C₁₀)-alkenyl substituted according to (10)
 (7) -(C₂-C₁₀)-alkynyl substituted according to (10)
 (8) -(C₆-C₂₂)-aryl substituted according to (10)
 (9) -W-(C₁-C₁₀)-alkyl, -W-(C₂-C₁₀)-alkenyl, -W-(C₂-C₁₀)-alkynyl, -W-(C₆-
 C₂₂)-aryl or W-H, wherein W= -S-, -O-, -NH-, -S-S-, -CO-, -COO-, -
 CO-NH-, -NH-CO-, -NH-CO-NH-, -NH-CS-NH-, -(CH₂)_n-[O-
 (CH₂)_r]_s-, where r and s are, independently of each other, an integer
 between 1 to 18 and n is 0 or 1 independently from r and s,
 (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be
 substituted by one or more moieties selected from the group consisting of
 -halogen, -SH, -NO₂, -CN, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -OR¹²,
 -COR¹¹, -NH-CO-NR⁵R⁶, -NH-CS-NR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-
 NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18
 and n is 0 or 1 independently from r and s,

wherein R^{11} is selected from the group consisting of $-NHR^{12}$, OR^{12} , and $-SR^{12}$

wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of $-H$, $-(C_1-C_{10})$ -alkyl, $-(C_2-C_{10})$ -alkenyl, $-(C_2-C_{10})$ -alkynyl, $-(C_6-C_{22})$ -aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,

said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being

unsubstituted or substituted by one or more moieties selected from the group consisting of $-halogen$, $-SH$, $-S-(C_1-C_6)$ -alkyl, $-(C_1-C_6)$ -alkoxy, $-OH$, $-NR^5R^6$, $-COR^{11}$, $-NH-CONR^5R^6$, $-NH-CSNR^5R^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ,

with the proviso that R^5 , R^6 or R^{12} is not a reporter group if the radicals (5) to (7) are substituted by $-NR^5R^6$, NHR^{12} , OR^{12} , or SR^{12} ;

R^2 , R^3 is independent from X , R^1 , R^2 and R^3 and is selected from the group of,

- (1) $-H$
- (2) (C_1-C_{10}) -alkyl,
- (3) (C_2-C_{10}) -alkenyl,
- (4) (C_2-C_{10}) -alkynyl,
- (5) (C_6-C_{22}) -aryl,
- (6) $-Z-(C_1-C_{10})$ -alkyl, $-Z-(C_2-C_{10})$ -alkenyl, $-Z-(C_2-C_{10})$ -alkynyl, $-Z-(C_6-C_{22})$ -aryl or $Z-H$, wherein $Z = -CO-$, $-CO-NH-$, $-CS-NH-$, $-(CH_2)_n-[O-(CH_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s ,
- (7) substituents (2) to (6)

wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of $-halogen$, $-NO_2$, $-OR^{12}$, $-CN$, $-SH$, $-S-(C_1-C_6)$ -alkyl, $-NR^5R^6$, $-N^+R^5R^6R^{12}$, $-COR^{11}$, $-NH-CONR^5R^6$, $-NH-CSNR^5R^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ,

wherein R^{11} is selected from the group consisting of $-NHR^{12}$ and OR^{12} ,

- 35 -

wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of -H, $-(C_1-C_{10})$ -alkyl, $-(C_2-C_{10})$ -alkenyl, $-(C_2-C_{10})$ -alkynyl, $-(C_6-C_{22})$ -aryl and a reporter group,

said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, $-S-(C_1-C_6)$ -alkyl, $-(C_1-C_6)$ -alkoxy, -OH, $-NR^5R^6$, $-COR^{11}$, $-NH-CONR^5R^6$, $-NH-CSNR^5R^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s;

X is independent from R^1 , R^2 or R^3 and is selected from the group consisting of N and CH; and

D is the position of attachment of the group to the rest of the nucleic acid binding compound

and any salts thereof.

A very preferred compound is a compound of formula V, wherein M' is O, R^{16} is H and R^{14} is selected from the group consisting of hydrogen and hydroxyl.

Those compounds can for example be used as 5'-terminally labeled probes. Regarding the definitions of the substituents, the definitions as given above apply if not indicated otherwise.

The backbone of the nucleic acid binding compound has the function to bear the base pairing heterocycles such that the compound can bind to a nucleic acid having a complementary sequence. Preferably, the degree of complementarity in the naturally occurring bases will be in the range from 70 % up to 100 % in a stretch of bases in a region effecting binding, compared to the stretch of same length in the region of the nucleic acid to be bound. Deletions and insertions of subunits in each sequence will therefor, in this calculation, be counted as gaps until the next fitting base and thus reduce complementarity by as many bases as the gap contains.

Preferred backbone contains sugar-phosphate moieties. From these, deoxy sugar containing backbones are further preferred.

Each moiety in the backbone bearing a moiety capable of base pairing to a nucleic acid of complementary sequence, including the moieties of the invention, are termed a subunit.

Compounds are known that have backbones mixed of different kinds of subunits. Recently, a new kind of non-natural nucleic acid binding compounds was described. They are termed Peptide Nucleic Acids (PNA), as they contain at least one peptide bond between the subunits (WO 92/20702). The nucleic acid binding compound of the present invention can
5 have any length. However, due to the convenience of chemical synthesis, compounds of a length of less than 100, more preferably from 10 to 30 subunits, for example nucleosides, are preferred.

Altering the thermal stability (T_m) of a duplex formed between a nucleic acid binding
10 compound according to the invention, e.g. used as a probe, and a second nucleic acid binding compound using the heterocyclic groups according to the invention and other analogues allows for optimization of duplex stability and mismatch discrimination (see e.g. Kwok, Shirley; Chang, Sheng Yung; Sninsky, John J.; Wang, Alice. A guide to the design and use of mismatched and degenerate primers. PCR Methods Appl. (1994), 3(4), 39-47).

15 One useful aspect of altering the T_m arises from the fact that Adenine-Thymine (A-T) duplexes have a lower T_m than Guanine-Cytosine (G-C) duplexes, due in part to the fact that the A-T duplexes have 2 hydrogen bonds per base-pair, while the G-C duplexes have 3 hydrogen bonds per base pair. For example in heterogeneous oligonucleotide arrays, in which there is a non-uniform distribution of bases, it can be difficult to optimize
20 hybridization conditions for all probes simultaneously. Thus, in some embodiments, it is desirable to destabilize G-C-rich duplexes and/or to increase the stability of A-T-rich duplexes while maintaining the sequence specificity of hybridization. This results in a harmonization or equalization of the contribution of each base pair to the melting temperature of a duplex. This is accomplished, e.g. by replacing one or more of the
25 heterocyclic groups in the nucleic acid binding compound used as a probe (or as the target nucleic acid) with certain modified, non-standard bases. Therefore, in another embodiment, the invention relates to nucleic acid binding compounds according to the invention wherein the nucleic acid binding compounds in addition to a heterocyclic group of formula I further contain a heterocyclic group different from the group of the general
30 formula I, i.e. at least one other heterocyclic group. Substitution of guanine residues with 7-deazaguanine, for example, will generally destabilize duplexes, whereas substituting adenine residues with 2,6-diaminopurine will enhance duplex stability. A variety of other modified bases are also incorporated into nucleic acids to enhance or decrease overall duplex stability while maintaining specificity of hybridization. The incorporation of 6-aza-
35 pyrimidine analogs into oligonucleotide probes generally decreases their binding affinity for complementary nucleic acids. Many 5-substituted pyrimidines substantially increase the

- stability of hybrids in which they have been substituted in place of the native pyrimidines in the sequence. Examples include 5-bromo-, 5-methyl-, 5-propynyl-, 5-(imidazol-2-yl)- and 5-(thiazol-2-yl)-derivatives of cytosine and uracil. Preferably the additional heterocyclic group is a pyrrolo-[2,3-d]-pyrimidine or a pyrazolo[3,4-d]-pyrimidine or an analogue thereof, in particular the said analogues of adenine or guanine. It should be emphasized that the invention also relates to the case where all other heterocyclic groups are those occurring in nature as adenine, guanine, uracil, cytosine or thymine. Many modified nucleosides, nucleotides and various bases suitable for incorporation into nucleosides are commercially available from a variety of manufacturers, including the SIGMA chemical company (Saint Louis, Mo.), R&D systems (Minneapolis, Minn.), Pharmacia LKB Biotechnology (Piscataway, N.J.), CLONTECH Laboratories, Inc. (Palo Alto, Calif.), Chem Genes Corp., Aldrich Chemical Company (Milwaukee, Wis.), Glen Research, Inc., GIBCO BRL Life Technologies, Inc. (Gaithersburg, Md.), Fluka Chemica-Biochemika Analytika (Fluka Chemie AG, Buchs, Switzerland), Invitrogen, San Diego, Calif., and Applied Biosystems (Foster City, Calif.), as well as many other commercial sources known to one of skill. Methods of attaching bases to sugar moieties to form nucleosides are known. See, e.g., Lukevics and Zablocka (1991), Nucleoside Synthesis: Organosilicon Methods Ellis Horwood Limited Chichester, West Sussex, England and the references therein. Methods of phosphorylating nucleosides to form nucleotides, and of incorporating nucleotides into oligonucleotides are also known. See, e.g., Agrawal (ed) (1993) Protocols for Oligonucleotides and Analogues, Synthesis and Properties, Methods in Molecular Biology volume 20, Humana Press, Towota, N.J., and the references therein. See also, Crooke and Lebleu, and Sanghvi and Cook, and the references cited therein, both supra.
- In yet another embodiment, the invention relates to nucleic acid binding compounds according to the invention wherein the nucleic acid binding compound according to the invention further contains at the 3'-end in addition to a heterocyclic group of formula I a heterocyclic group different from the group of the general formula I, i.e. at least one other heterocyclic group. Preferably, the nucleic acid binding compound according to the invention additionally comprises a heterocyclic group which is a pyrrolo-[2,3-d]-pyrimidine or a pyrazolo[3,4-d]-pyrimidine or an analogue thereof, preferably the said analogues of guanine or adenine.

- The invention further contemplates the binding product of a nucleic acid binding compound according to the invention and a second nucleic acid binding compound, the nucleic acid binding compound according to the invention and the second nucleic acid

binding compound being bound to each other by base pairing in parallel or antiparallel orientation. In addition to heterocyclic groups with formula I with substituents as defined above, the nucleic acid binding compound according to the invention may contain other natural nucleobase or nucleobases not occurring in nature as e.g. nucleobases with
5 heterocyclic groups according to the invention, heterocyclic groups as pyrrolo-[2,3-d]-pyrimidine or pyrazolo[3,4-d]-pyrimidine or analogues thereof, preferably the said analogues of guanine or adenine or 7-deaza-guanine. Further non-natural heterocyclic groups are known to the person skilled in the art.

10 Another embodiment of the invention is a nucleic acid binding compound wherein the heterocyclic group of formula I as defined above is incorporated to compensate for a decrease of the melting point created by the attachment of the reporter groups, preferably 1 to 5 nucleotides separated from the nucleotide to which a reporter group is attached. This is because a reporter group leads to disturbances of the hybridization efficiency of a nucleic
15 acid binding compound close to the point or nucleotide whereto the reporter group is attached.

Another embodiment of the invention is a nucleic acid binding compound wherein the heterocyclic group of formula I as defined above is incorporated to compensate for a
20 decrease of the melting point created by mismatches discrimination. This problem has been discussed by Kwok, Shirley; Chang, Sheng Yung; Sninsky, John J.; Wang, Alice. A guide to the design and use of mismatched and degenerate primers. PCR Methods Appl. (1994), 3(4), 39-47. This is particularly useful for the amplification of viral subtypes where the hybridization stretch does not contain complementary bases over the total length of the
25 stretch.

In an embodiment, the invention relates to nucleic acid binding compounds according to the invention wherein a protecting group substitutes one or two hydrogen atoms of a -OH, -SH, -NH₂, -NH-alkyl, -NH-alkenylene, -NH-alkynylene, or a NH-aryl group, although it
30 is preferred that the nucleic acid binding compounds according to the invention only contain a few protecting groups or even none.

One particular preferred embodiment, is the use of the nucleic acid binding compounds in field of arrays of nucleic acid binding compounds bound to a solid surface (see e.g.
35 US5,143,854, US 6,022,963, US 6,156,501, WO90/15070, WO 92/10092), which has the properties as described in these references and can be manufactured as described therein or


by Niemeyer and Blohm (Angew. Chem. Int. Ed. 1999, 38, 2865-2869). Therefore, in a preferred embodiment, the invention relates to a composition for analyzing interactions between nucleic acid binding compounds whereby one nucleic acid binding compound is a target nucleic acid. The composition comprises an array of a plurality of nucleic acid binding compounds having different sequences, wherein said plurality of nucleic acid binding compounds are coupled to a solid substrate at known locations and are selected to bind to complementary nucleic acid binding compounds or target nucleic acids whereby only the nucleic acid binding compounds or the nucleic acid binding compounds and the complementary nucleic acid binding compounds (or target nucleic acids) together are nucleic acid binding compounds comprising a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases, wherein a heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof. Different kinds of supports are possible as e.g. nonporous supports or other solid supports less porous than porous beads, resins, or other supports work well and are often preferable. One such support is a resin in the form of beads. In general, the bead size is in the range of 1 nm to 100 μm , but a more massive solid support of up to 1 mm in size may sometimes be used. Particularly preferred resins include Sasrin resin (a polystyrene resin available from Bachem Bioscience, Switzerland); and TentaGel S AC, TentaGel PHB, or TentaGel S NH_2 resin (polystyrene-polyethylene glycol copolymer resins available from Rappe Polymere, Tubingen, Germany). Other preferred supports are commercially available and described by Novabiochem, La Jolla, Calif. In other embodiments, the solid substrate is flat, or alternatively, may take on alternative surface configurations. For example, the solid substrate may contain raised or depressed regions on which synthesis takes place or is coated with porous SiO_2 / glass. In some embodiments, the solid substrate will be chosen to provide appropriate light-absorbing characteristics. For example, the substrate may be a polymerized Langmuir Blodgett film, functionalized glass, Si, Ge, GaAs, GaP, SiO_2 , SiN_4 , modified silicon, or any one of a variety of gels or polymers such as (poly)tetrafluorethylene, (poly)vinylidenedifluoride, polystyrene, polycarbonate, or combinations thereof. Other suitable solid substrate material will be readily apparent to those of skill in the art. Preferably, the surface of the solid substrate will contain reactive groups, which could be carboxyl, amino, hydroxyl, thiol, or the like. More preferably, the surface will be optically transparent and will have surface Si--OH functionalities, such as are found on silica surfaces. Particularly preferred is therefore a composition, wherein the solid substrate is selected from the group consisting of silica, polymeric materials, glass, porous glass, beads, chips, and slides.

Preferred is a composition according to the invention wherein only the nucleic acid binding compounds or the nucleic acid binding compounds and the complementary nucleic acid binding compounds (or target nucleic acid) are nucleic acid binding compounds according to the invention, i.e. they contain a heterocyclic group with formula I as described above. The complementary nucleic acid binding compounds (or target nucleic acid) may contain a heterocyclic group according to formula I as described above when they are amplified with e.g. the polymerase chain reaction in the presence of a triphosphate according to the invention containing a heterocyclic group of formula I as described above.

Preferred is a composition according to the invention, wherein the nucleic acid binding compounds comprise a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic groups is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof. In a preferred embodiment the substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof is a substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof or a 7-substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof. In another embodiment of the invention the substituted pyrazolo[3,4-d]pyrimidine analogue is a substituted pyrazolo[3,4-d]pyrimidine analogue of adenine or guanine or a 7-substituted pyrazolo[3,4-d]pyrimidine analogue of adenine or guanine, wherein the adenine or guanine analogues may preferably carry the same substituents R^1 in the 7-position or N-substituents R^2 and R^3 as set out directly below for the substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof or a 7-substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof.

In a very preferred embodiment of the invention the substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof or the 7-substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof has the general formula I

Formula I



30

R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of

- (1) -F, -Cl, -Br or -I,
- (2) Nitro
- (3) Cyano
- (4) -COO^-
- 5 (5) $\text{-(C}_1\text{-C}_{10}\text{)-alkyl}$ substituted according to (10)
- (6) $\text{-(C}_2\text{-C}_{10}\text{)-alkenyl}$ substituted according to (10)
- (7) $\text{-(C}_2\text{-C}_{10}\text{)-alkynyl}$ substituted according to (10)
- (8) $\text{-(C}_6\text{-C}_{22}\text{)-aryl}$ substituted according to (10)
- 10 (9) $\text{-W-(C}_1\text{-C}_{10}\text{)-alkyl}$, $\text{-W-(C}_2\text{-C}_{10}\text{)-alkenyl}$, $\text{-W-(C}_2\text{-C}_{10}\text{)-alkynyl}$, $\text{-W-(C}_6\text{-C}_{22}\text{)-aryl}$ or W-H , wherein $\text{W= -S-, -O-, -NH-, -S-S-, -CO-, -COO-, -CO-NH-, -NH-CO-, -NH-CO-NH-, -NH-CS-NH-, -(CH}_2\text{)}_n\text{-[O-(CH}_2\text{)}_r\text{]}_s\text{-}$, where r and s are, independently of each other, an integer between 1 to 18 and n is 0 or 1 independently from r and s ,
- 15 (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -SH, -NO_2 , -CN, $\text{-S-(C}_1\text{-C}_6\text{)-alkyl}$, $\text{-NR}^5\text{R}^6$, $\text{-N}^+\text{R}^5\text{R}^6\text{R}^{12}$, -OR^{12} , -COR^{11} , $\text{-NH-CO-NR}^5\text{R}^6$, $\text{-NH-CS-NR}^5\text{R}^6$ and $\text{-(CH}_2\text{)}_n\text{-[O-(CH}_2\text{)}_r\text{]}_s\text{-NR}^5\text{R}^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ,
- 20 wherein R^{11} is selected from the group consisting of -NHR^{12} , OR^{12} , and -SR^{12}
 wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of -H, $\text{-(C}_1\text{-C}_{10}\text{)-alkyl}$, $\text{-(C}_2\text{-C}_{10}\text{)-alkenyl}$, $\text{-(C}_2\text{-C}_{10}\text{)-alkynyl}$, $\text{-(C}_6\text{-C}_{22}\text{)-aryl}$ and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,
- 25 said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, $\text{-S-(C}_1\text{-C}_6\text{)-alkyl}$, $\text{-(C}_1\text{-C}_6\text{)-alkoxy}$, -OH, $\text{-NR}^5\text{R}^6$, -COR^{11} , $\text{-NH-CONR}^5\text{R}^6$, $\text{-NH-CSNR}^5\text{R}^6$ and $\text{-(CH}_2\text{)}_n\text{-[O-(CH}_2\text{)}_r\text{]}_s\text{-NR}^5\text{R}^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ,
- 30 with the proviso that R^5 , R^6 or R^{12} is not a reporter group if the radicals (5) to (7) are substituted by $\text{-NR}^5\text{R}^6$, NHR^{12} , OR^{12} , or SR^{12} ;
- 35 R^2 , R^3 is independent from X , R^1 , R^2 and R^3 and is selected from the group of,

- (1) -H
- (2) (C₁-C₁₀)-alkyl,
- (3) (C₂-C₁₀)-alkenyl,
- (4) (C₂-C₁₀)-alkynyl,
- 5 (5) (C₆-C₂₂)-aryl,
- (6) -Z-(C₁-C₁₀)-alkyl, -Z-(C₂-C₁₀)-alkenyl, -Z-(C₂-C₁₀)-alkynyl, -Z-(C₆-C₂₂)-aryl or Z-H, wherein Z = -CO-, -CO-NH-, -CS-NH-, -(CH₂)_n-[O-(CH₂)_r]_s-, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s,
- 10 (7) substituents (2) to (6)
- wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -NO₂, -OR¹², -CN, -SH, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,
- 15 wherein R¹¹ is selected from the group consisting of -NHR¹² and OR¹²,
- wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group,
- 20 said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s;
- 25 X is independent from R¹, R² or R³ and is selected from the group consisting of N and CH
- 30 D is the position of attachment of the group to the rest of the nucleic acid binding compound.

In a very preferred embodiment, the nucleic acid binding compound is a nucleic acid binding compound according to the invention, preferably R¹=Br or R¹=I. Further preferred

is a composition which comprises an array of nucleic acid binding compounds 5 to 20 nucleotides in length.

The invention is further related to a binding product of a first nucleic acid binding
5 compound according to the invention or a composition according to the invention with a second nucleic acid binding compound or a second a nucleic acid binding compound according to the invention, wherein the first nucleic acid binding compound or the composition and the second nucleic acid binding compound being bound to each other by base pairing in parallel or antiparallel orientation.

10

The invention is further related to methods for the synthesis of the nucleic acid binding compounds according to the invention and to compounds useful in these methods. The nucleic acid binding compound of the present invention can be prepared in solution or, preferably, on a solid phase, where appropriate using an automatic synthesis device. The
15 oligomers can be assembled stepwise by successively condensing a mononucleotide, which in each case possesses a nucleotide base, onto an appropriately derivatized support or onto a growing oligomer chain. Alternatively, the nucleic acid binding compounds can be assembled by joining dinucleotides or trinucleotides together [S. Beaucage et al., Tetrahedron, 48 (12), 2223-2311, (1992); and Tetrahedron, 48 (28), 6123-6194, (1993)].
20 This is particularly advantageous when synthesizing oligonucleotides which posses modified phosphate bridges.


The oligonucleotides are assembled using methods which are known to the person skilled in the art, such as the triester method, the H-phosphonate method or the phosphoramidite
25 method [E. Sonveaux, (1986), Bioorganic Chemistry, 14, 274-325; S. L. Beaucage et al., (1992), Tetrahedron, 48, 2223-2311].

The compounds according to the present invention can be advantageously used in oligonucleotide synthesis as the ammonia hydrolysis of the protecting groups of the
30 pyrazolo[3,4-d]pyrimidine nucleosides is quicker than the slow ammonia hydrolysis of 2-amino-adenosine which takes several days.

A further subject of the invention is therefore a method for the chemical synthesis of nucleic acid binding compounds of the present invention using activated subunits, wherein
35 said subunit contains a group of formula I. The most preferred method of chernical synthesis uses the phosphoramidite approach. A particularly preferred method uses a

activated subunit one or more compounds of general formula VII. This method has the advantage that it is very convenient and the reagents necessary, for example a phosphoramidite containing a group of formula I, are possible to be included easily.

- 5 A further subject of the invention are therefore compounds of the general formula VII




wherein

- 10 A is selected from the group consisting of O, S, CH₂ and N-(C₁-C₆)-alkyl,
M and M' are independently selected from the group consisting of oxy, sulfanediyl, -
NR²², -(C₁-C₁₀)-alkyl, or -O-(C₁-C₁₀)-alkyl-O-, and -S-
(C₁-C₁₀)-alkyl-O- and -NR²²-(C₁-C₆)-alkyl-O-,
R²² is selected from the group of -H and -(C₁-C₁₀)-alkyl,
15 R¹⁴ is selected from the group consisting of -H, -OR³¹, -(C₁-C₁₀)-alkoxy,
-(C₂-C₁₀)-alkenyloxy, -(C₂-C₁₀)-alkynyloxy, -halogen, -azido, NHR³¹, SR³¹,
R³¹ is a protecting group or a reporter group,
R³² and R¹⁷ are independently selected from the group consisting of -H, -(C₁-C₁₀)-
alkyl, -(C₂-C₁₀)-alkenyl, -(C₆-C₂₂)-aryl, or wherein NR³²R¹⁷ can form together
20 with N a 5-6-membered heterocyclic ring,
R¹⁸ is selected from the group consisting of -(C₂-C₆)-alkenyloxy, substituted or
unsubstituted -(C₁-C₆)-alkyl, unsubstituted -(C₁-C₆)-alkoxy or -(C₁-C₆)-alkoxy
substituted one or more times by a group selected from the group consisting of
-halogen, p-nitroaryloxy and -cyano, and
25 B is a group of formula I

- 45 -

Formula I




R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of

- (1) -F, -Cl, -Br or -I,
- (2) Nitro
- (3) Cyano
- (4) $-\text{COO}^-$
- (5) $-(\text{C}_1\text{-C}_{10})\text{-alkyl}$ substituted according to (10)
- (6) $-(\text{C}_2\text{-C}_{10})\text{-alkenyl}$ substituted according to (10)
- (7) $-(\text{C}_2\text{-C}_{10})\text{-alkynyl}$ substituted according to (10)
- (8) $-(\text{C}_6\text{-C}_{22})\text{-aryl}$ substituted according to (10)
- (9) $-\text{W}-(\text{C}_1\text{-C}_{10})\text{-alkyl}$, $-\text{W}-(\text{C}_2\text{-C}_{10})\text{-alkenyl}$, $-\text{W}-(\text{C}_2\text{-C}_{10})\text{-alkynyl}$, $-\text{W}-(\text{C}_6\text{-C}_{22})\text{-aryl}$ or W-H , wherein $\text{W} = -\text{S}-$, $-\text{O}-$, $-\text{NH}-$, $-\text{S-S}-$, $-\text{CO}-$, $-\text{COO}-$, $-\text{CO-NH}-$, $-\text{NH-CO}-$, $-\text{NH-CO-NH}-$, $-\text{NH-CS-NH}-$, $-(\text{CH}_2)_n\text{-[O-(CH}_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 0 or 1 independently from r and s,
- (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -SH, $-\text{NO}_2$, -CN, $-\text{S}-(\text{C}_1\text{-C}_6)\text{-alkyl}$, $-\text{NR}^5\text{R}^6$, $-\text{N}^+\text{R}^5\text{R}^6\text{R}^{12}$, $-\text{OR}^{12}$, $-\text{COR}^{11}$, $-\text{NH-CO-NR}^5\text{R}^6$, $-\text{NH-CS-NR}^5\text{R}^6$ and $-(\text{CH}_2)_n\text{-[O-(CH}_2)_r]_s-\text{NR}^5\text{R}^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s, wherein R^{11} is selected from the group consisting of $-\text{NHR}^{12}$, OR^{12} , and $-\text{SR}^{12}$ wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of -H, $-(\text{C}_1\text{-C}_{10})\text{-alkyl}$, $-(\text{C}_2\text{-C}_{10})\text{-alkenyl}$, $-(\text{C}_2\text{-C}_{10})\text{-alkynyl}$, $-(\text{C}_6\text{-C}_{22})\text{-aryl}$ and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,

(11)

- 46 -



wherein Nuc is the link to formula I and n is any integer from 1 to 18

said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being

unsubstituted or substituted by one or more moieties selected from the group
 consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -
 COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r
 and s are independently of each other an integer of from 1 to 18 and n is 0 or 1
 independently from r and s,

with the proviso that R⁵, R⁶ or R¹² is not a reporter group if the radicals (5) to (7)
 are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;

said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being

unsubstituted or substituted by one or more moieties selected from the group
 consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -
 COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r
 and s are independently of each other an integer of from 1 to 18 and n is 0 or 1
 independently from r and s,

with the proviso that R⁵, R⁶ or R¹² is not a reporter group if the radicals (5) to (7)
 are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;

R², R³ is independent from X, R¹, R² and R³ and is selected from the group of,

- (1) -H
- (2) (C₁-C₁₀)-alkyl,
- (3) (C₂-C₁₀)-alkenyl,
- (4) (C₂-C₁₀)-alkynyl,
- (5) (C₆-C₂₂)-aryl,

- (6) $-Z-(C_1-C_{10})\text{-alkyl}$, $-Z-(C_2-C_{10})\text{-alkenyl}$, $-Z-(C_2-C_{10})\text{-alkynyl}$, $-Z-(C_6-C_{22})\text{-aryl}$ or $Z\text{-H}$, wherein $Z = -CO-$, $-CO-NH-$, $-CS-NH-$, $-(CH_2)_n-[O-(CH_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s ,
- 5 (7) substituents (2) to (6)
- wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of $-\text{halogen}$, $-\text{NO}_2$, $-\text{OR}^{12}$, $-\text{CN}$, $-\text{SH}$, $-\text{S}-(C_1-C_6)\text{-alkyl}$, $-\text{NR}^5\text{R}^6$, $-\text{N}^+\text{R}^5\text{R}^6\text{R}^{12}$, $-\text{COR}^{11}$, $-\text{NH-CONR}^5\text{R}^6$, $-\text{NH-CSNR}^5\text{R}^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-\text{NR}^5\text{R}^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ,
- 10 wherein R^{11} is selected from the group consisting of $-\text{NHR}^{12}$ and OR^{12} ,
- wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of $-\text{H}$, $-(C_1-C_{10})\text{-alkyl}$, $-(C_2-C_{10})\text{-alkenyl}$, $-(C_2-C_{10})\text{-alkynyl}$, $-(C_6-C_{22})\text{-aryl}$ and a reporter group,
- 15 said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of $-\text{halogen}$, $-\text{SH}$, $-\text{S}-(C_1-C_6)\text{-alkyl}$, $-(C_1-C_6)\text{-alkoxy}$, $-\text{OH}$, $-\text{NR}^5\text{R}^6$, $-\text{COR}^{11}$, $-\text{NH-CONR}^5\text{R}^6$, $-\text{NH-CSNR}^5\text{R}^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-\text{NR}^5\text{R}^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ;
- 20
- X is independent from R^1 , R^2 or R^3 and is selected from the group consisting of N and CH
- 25 D is the position of attachment of the group to the rest of the nucleic acid binding compound.

with the proviso that one or two hydrogen atoms of any $-\text{OH}$, $-\text{SH}$, $-\text{NH}_2$, $-\text{NH-alkyl}$, $-\text{NH-alkenylene}$, $-\text{NH-alkynylene}$, or $-\text{NH-aryl}$ group are substituted by a protecting group,

30

and any salts thereof.


After suitable protective groups for the amino groups at position 2 and 6 and for the free 5'-hydroxyl group of the sugar moiety have been introduced, the monomers are converted

35 into the corresponding phosphonate or phosphoramidite derivatives. Suitable amino protective groups, for example in the form of acyl protective groups (e.g. isobutyryl, acetyl

or phenoxyacetyl), are inserted using well-known methods [J. C. Schulhof, D. Molko, R. Teoule, (1987), *Nucleic Acids Res.*, 15, 397-416]. An example of a suitable protective group for the free 5'-OH group of the sugar is the 4,4'-dimethoxytrityl residue, whose insertion is likewise effected using known methods [C. B. Reese (1978), *Tetrahedron*, 34, 3143; D. Flockerzi et al., (1981), *Liebigs Ann. Chem.*, 1568]. The monomers which have been protected in this way can be converted into the corresponding phosphonates in accordance with a protocol due to Froehler et al. [B. C. Froehler et al., (1986), *Nucleic Acids Res.*, 14, 5399]. Cyanoethyl-phosphoramidite derivatives can, for example, be prepared by reacting the monomers with chloro- β -cyanoethoxy-(N,N-diisopropylamino)phosphane in anhydrous dichloromethane [N. D. Sinha et al., (1984), *Nucleic Acids Res.*, 12, 4539].

Further subject of the invention are compounds of the general formula IX

Formula IX



15


wherein

- A is selected from the group consisting of O, S, CH₂ and N-(C₁-C₆)-alkyl,
M and M' are independently selected from the group consisting of oxy, sulfanediyl, -
NR²², -(C₁-C₁₀)-alkyl, or -O-(C₁-C₁₀)-alkyl-O-, and -S-(C₁-C₁₀)-alkyl-O- and -
NR²²-(C₁-C₆)-alkyl-O-,
R²² is selected from the group of -H and -(C₁-C₁₀)-alkyl,
R¹⁴ is selected from the group consisting of -H, -OR³¹, -(C₁-C₁₀)-alkoxy,
-(C₂-C₁₀)-alkenyloxy, -(C₂-C₁₀)-alkynyloxy, -halogen, -azido, NHR³¹, SR³¹, or
O-reporter group,
R³¹ is a protecting group or a reporter group,
R³² and R¹⁷ are independently selected from the group consisting of -H, -(C₁-C₁₀)-
alkyl, -(C₂-C₁₀)-alkenyl, -(C₆-C₂₂)-aryl, or wherein NR³²R¹⁷ can form together
with N a 5-6-membered heterocyclic ring,

R^{18} is selected from the group consisting of $-(C_2-C_6)$ -alkenyloxy, substituted or unsubstituted $-(C_1-C_6)$ -alkyl, unsubstituted $-(C_1-C_6)$ -alkoxy or $-(C_1-C_6)$ -alkoxy substituted one or more times by a group selected from the group consisting of -halogen, p-nitroaryloxy and -cyano, and

5 B is a group of formula I

Formula I



R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of


- (1) -F, -Cl, -Br or -I,
- (2) Nitro
- 10 (3) Cyano
- (4) $-\text{COO}^-$
- (5) $-(C_1-C_{10})$ -alkyl substituted according to (10)
- (6) $-(C_2-C_{10})$ -alkenyl substituted according to (10)
- (7) $-(C_2-C_{10})$ -alkynyl substituted according to (10)
- 15 (8) $-(C_6-C_{22})$ -aryl substituted according to (10)
- (9) $-\text{W}-(C_1-C_{10})$ -alkyl, $-\text{W}-(C_2-C_{10})$ -alkenyl, $-\text{W}-(C_2-C_{10})$ -alkynyl, $-\text{W}-(C_6-C_{22})$ -aryl or $\text{W}-\text{H}$, wherein $\text{W} = -\text{S}-, -\text{O}-, -\text{NH}-, -\text{S}-\text{S}-, -\text{CO}-, -\text{COO}-, -\text{CO}-\text{NH}-, -\text{NH}-\text{CO}-, -\text{NH}-\text{CO}-\text{NH}-, -\text{NH}-\text{CS}-\text{NH}-, -(\text{CH}_2)_n-[\text{O}-(\text{CH}_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 0 or 1 independently from r and s,
- 20 (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -SH, $-\text{NO}_2$, -CN, $-\text{S}-(C_1-C_6)$ -alkyl, $-\text{NR}^5\text{R}^6$, $-\text{N}^+\text{R}^5\text{R}^6\text{R}^{12}$, $-\text{OR}^{12}$, $-\text{COR}^{11}$, $-\text{NH}-\text{CO}-\text{NR}^5\text{R}^6$, $-\text{NH}-\text{CS}-\text{NR}^5\text{R}^6$ and $-(\text{CH}_2)_n-[\text{O}-(\text{CH}_2)_r]_s-\text{NR}^5\text{R}^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,
- 25 wherein R^{11} is selected from the group consisting of $-\text{NHR}^{12}$, OR^{12} , and $-\text{SR}^{12}$

30 wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of -H, $-(C_1-C_{10})$ -alkyl, $-(C_2-C_{10})$ -alkenyl, $-(C_2-C_{10})$ -alkynyl, $-(C_6-C_{22})$ -aryl and a reporter group, a group which facilitates intracellular

- 50 -

uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,

(11)



5

wherein Nuc is the link to formula I and n is any integer from 1 to 18 said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,

10

with the proviso that R⁵, R⁶ or R¹² is not a reporter group if the radicals (5) to (7) are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;

15 R², R³

is independent from X, R¹, R² and R³ and is selected from the group of,

(1) -H

(2) (C₁-C₁₀)-alkyl,(3) (C₂-C₁₀)-alkenyl,(4) (C₂-C₁₀)-alkynyl,

20

(5) (C₆-C₂₂)-aryl,

(6) -Z-(C₁-C₁₀)-alkyl, -Z-(C₂-C₁₀)-alkenyl, -Z-(C₂-C₁₀)-alkynyl, -Z-(C₆-C₂₂)-aryl or Z-H, wherein Z = -CO-, -CO-NH-, -CS-NH-, -(CH₂)_n-[O-(CH₂)_r]_s-, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s,

25

(7) substituents (2) to (6)


- wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -NO₂, -OR¹², -CN, -SH, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,
- wherein R¹¹ is selected from the group consisting of -NHR¹² and OR¹²,
- wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group,
- said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s;
- X is independent from R¹, R² or R³ and is selected from the group consisting of N and CH
- D is the position of attachment of the group to the rest of the nucleic acid binding compound.

with the proviso that one or two hydrogen atoms of any -OH, -SH, -NH₂, -NH-alkyl, -NH-alkenylene, -NH-alkynylene, or -NH-aryl group are substituted by a protecting group, and any salts thereof.

Those compounds can be used like those of formula VII in chemical synthesis.

A further subject of the invention are compounds of the general formula X wherein

Formula X



M and M' are independently selected from the group consisting of oxy, sulfanediyl, -NR²², -
(C₁-C₁₀)-alkyl, or -O-(C₁-C₁₀)-alkyl-O-, and -S-


5 (C₁-C₁₀)-alkyl-O- and -NR²²-(C₁-C₆)-alkyl-O-,

R²² is selected from the group of -H and -(C₁-C₁₀)-alkyl,

R¹⁴ is selected from the group consisting of -H, -OR³¹, -(C₁-C₁₀)-alkoxy,
-(C₂-C₁₀)-alkenyloxy, -(C₂-C₁₀)-alkynyloxy, -halogen, -azido, NHR³¹, SR³¹, or
O-reporter,

10 R³¹ is a protecting group or a reporter group,

Formula I



R¹ is independent from X, R² or R³ and is selected from the group consisting of

- (1) -F, -Cl, -Br or -I,
- (2) Nitro
- 15 (3) Cyano
- (4) -COO⁻
- (5) -(C₁-C₁₀)-alkyl substituted according to (10)
- (6) -(C₂-C₁₀)-alkenyl substituted according to (10)
- (7) -(C₂-C₁₀)-alkynyl substituted according to (10)
- 20 (8) -(C₆-C₂₂)-aryl substituted according to (10)
- (9) -W-(C₁-C₁₀)-alkyl, -W-(C₂-C₁₀)-alkenyl, -W-(C₂-C₁₀)-alkynyl, -W-(C₆-
C₂₂)-aryl or W-H, wherein W= -S-, -O-, -NH-, -S-S-, -CO-, -COO-, -


CO-NH-, -NH-CO-, -NH-CO-NH-, -NH-CS-NH-, $-(CH_2)_n-[O-(CH_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 0 or 1 independently from r and s,

- (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -SH, -NO₂, -CN, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -OR¹², -COR¹¹, -NH-CO-NR⁵R⁶, -NH-CS-NR⁵R⁶ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,

wherein R¹¹ is selected from the group consisting of -NHR¹², OR¹², and -SR¹²

wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,

(11)



wherein Nuc is the link to formula I and n is any integer from 1 to 18 said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r

and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,

with the proviso that R^5 , R^6 or R^{12} is not a reporter group if the radicals (5) to (7) are substituted by $-NR^5R^6$, NHR^{12} , OR^{12} , or SR^{12} ;

- 5 R^2, R^3 is independent from X, R^1 , R^2 and R^3 and is selected from the group of,
- (1) $-H$
 - (2) (C_1-C_{10}) -alkyl,
 - (3) (C_2-C_{10}) -alkenyl,
 - (4) (C_2-C_{10}) -alkynyl,
 - 10 (5) (C_6-C_{22}) -aryl,
 - (6) $-Z-(C_1-C_{10})$ -alkyl, $-Z-(C_2-C_{10})$ -alkenyl, $-Z-(C_2-C_{10})$ -alkynyl, $-Z-(C_6-C_{22})$ -aryl or $Z-H$, wherein $Z = -CO-$, $-CO-NH-$, $-CS-NH-$, $-(CH_2)_n-$, $[O-(CH_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s,
 - 15 (7) substituents (2) to (6)
- wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, - NO_2 , $-OR^{12}$, $-CN$, $-SH$, $-S-(C_1-C_6)$ -alkyl, $-NR^5R^6$, $-N^+R^5R^6R^{12}$, $-COR^{11}$, $-NH-CONR^5R^6$, $-NH-CSNR^5R^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,
- 20 wherein R^{11} is selected from the group consisting of $-NHR^{12}$ and OR^{12} ,
- wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of $-H$, $-(C_1-C_{10})$ -alkyl, $-(C_2-C_{10})$ -alkenyl, $-(C_2-C_{10})$ -alkynyl, $-(C_6-C_{22})$ -aryl and a reporter group,
- 25 said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, $-SH$, $-S-(C_1-C_6)$ -alkyl, $-(C_1-C_6)$ -alkoxy, $-OH$, $-NR^5R^6$, $-COR^{11}$, $-NH-CONR^5R^6$, $-NH-CSNR^5R^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s;
- 30 X is independent from R^1 , R^2 or R^3 and is selected from the group consisting of N and CH
- 35 D is the position of attachment of the group to the rest of the nucleic acid binding compound.


with the proviso that one or two hydrogen atoms of any -OH, -SH, -NH₂, -NH-alkyl, -NH-alkenylene, -NH-alkynylene, or -NH-aryl group are substituted by a protecting group, and any salts thereof. Those compounds are useful in chemical synthesis of nucleic acid binding
 5 compounds as mentioned above and the precursors thereof.

In another option which is more suited for long oligomers and those based on natural backbones, the oligomers are produced enzymatically. In this case, a starting oligomer is reacted with a polymerase and a triphosphate or modified triphosphate such that a
 10 monophosphate or a modified monophosphate is attached to a terminus of the oligomer, thus elongating the oligomer. Also for this method, the man skilled in the art will know several possible formates, like the nick-translation approach, or the simple primer extension (J. Sambrook, E.F. Fritsch, T. Maniatis, Molecular Cloning - A laboratory Manual, Cold Spring Harbor Laboratory Press 1989).

15 A further subject of the invention is therefore a method for the enzymatic synthesis of a nucleic acid binding compound according to the invention comprising reacting a triphosphate subunit with a primer using a nucleic acid as a template for the elongation of the primer, wherein the triphosphate subunit contains a heterocyclic group of formula I.
 20 Preferably, the triphosphate subunit has the formula VI. For example, 7- or 8-substituted 7-deaza-2'-deoxyadenosine and guanosine-triphosphates can be easily incorporated enzymatically into DNA by various DNA polymerases (WO 00/68422).

A further subject of the present invention are therefore compounds of the general formula
 25 VI


Formula VI



wherein

- A is selected from the group consisting of O, S, CH₂ and N-(C₁-C₆)-alkyl,
 M and M' are independently selected from the group consisting of oxy, sulfanediyl, -
 5 NR²²-, -(C₁-C₁₀)-alkyl-, or -O-(C₁-C₁₀)-alkyl-O-, and -S-(C₁-C₁₀)-alkyl-O- and
 -NR²²-(C₁-C₆)-alkyl-O-,
 R²² is selected from the group of -H, -(C₁-C₁₀)-alkyl, a protecting group and a
 reporter group,
 R¹⁴ is selected from the group consisting of -H, -OR³¹, -(C₁-C₁₀)-alkoxy, O-
 10 protecting group, S-protecting group, NH₂-protecting group, -(C₂-C₁₀)-
 alkenyloxy, -(C₂-C₁₀)-alkynyloxy, -halogen, -azido, SH, -(C₁-C₁₀)-
 alkylmercapto, and -O-solid phase,
 R¹⁵ and R¹⁶ are independently selected from the group consisting of -H, -(C₁-C₆)-
 alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₂-C₁₀)-alkyl-carbonyl, -(C₃-C₁₉)-
 15 alkenyl-carbonyl, -(C₃-C₁₉)-alkynyl-carbonyl, -(C₆-C₁₄)-aryl-(C₁-C₁₀)-alkyl,
 protecting group and a solid phase
 B is the link to a moiety of formula I,

Formula I



- 20 R¹ is independent from X, R² or R³ and is selected from the group consisting of
 (1) -F, -Cl, -Br or -I,
 (2) Nitro
 (3) Cyano
 (4) -COO⁻
 25 (5) -(C₁-C₁₀)-alkyl substituted according to (10)
 (6) -(C₂-C₁₀)-alkenyl substituted according to (10)
 (7) -(C₂-C₁₀)-alkynyl substituted according to (10)
 (8) -(C₆-C₂₂)-aryl substituted according to (10)
 (9) -W-(C₁-C₁₀)-alkyl, -W-(C₂-C₁₀)-alkenyl, -W-(C₂-C₁₀)-alkynyl, -W-(C₆-
 30 C₂₂)-aryl or W-H, wherein W= -S-, -O-, -NH-, -S-S-, -CO-, -COO-, -
 CO-NH-, -NH-CO-, -NH-CO-NH-, -NH-CS-NH-, -(CH₂)_n-[O-

- 57 -

$(\text{CH}_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 0 or 1 independently from r and s ,


- (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of
- 5 -halogen, -SH, -NO₂, -CN, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -OR¹², -COR¹¹, -NH-CO-NR⁵R⁶, -NH-CS-NR⁵R⁶ and $-(\text{CH}_2)_n-[\text{O}-(\text{CH}_2)_r]_s-$ NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ,

wherein R¹¹ is selected from the group consisting of -NHR¹², OR¹², and -SR¹²

wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound

15 hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,

(11)



wherein Nuc is the link to formula I and n is any integer from 1 to 18

20 said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and $-(\text{CH}_2)_n-[\text{O}-(\text{CH}_2)_r]_s-$ NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1

25 independently from r and s ,

- with the proviso that R^5 , R^6 or R^{12} is not a reporter group if the radicals (5) to (7) are substituted by $-NR^5R^6$, NHR^{12} , OR^{12} , or SR^{12} ;
- R^2 , R^3 is independent from X , R^1 , R^2 and R^3 and is selected from the group of,
- (1) $-H$
 - (2) (C_1-C_{10}) -alkyl,
 - (3) (C_2-C_{10}) -alkenyl,
 - (4) (C_2-C_{10}) -alkynyl,
 - (5) (C_6-C_{22}) -aryl,
 - (6) $-Z-(C_1-C_{10})$ -alkyl, $-Z-(C_2-C_{10})$ -alkenyl, $-Z-(C_2-C_{10})$ -alkynyl, $-Z-(C_6-C_{22})$ -aryl or $Z-H$, wherein $Z = -CO-$, $-CO-NH-$, $-CS-NH-$, $-(CH_2)_n-$, $[O-(CH_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s ,
 - (7) substituents (2) to (6)
- wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of $-halogen$, $-NO_2$, $-OR^{12}$, $-CN$, $-SH$, $-S-(C_1-C_6)$ -alkyl, $-NR^5R^6$, $-N^+R^5R^6R^{12}$, $-COR^{11}$, $-NH-CONR^5R^6$, $-NH-CSNR^5R^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ,
- wherein R^{11} is selected from the group consisting of $-NHR^{12}$ and OR^{12} ,
- wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of $-H$, $-(C_1-C_{10})$ -alkyl, $-(C_2-C_{10})$ -alkenyl, $-(C_2-C_{10})$ -alkynyl, $-(C_6-C_{22})$ -aryl and a reporter group,
- said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of $-halogen$, $-SH$, $-S-(C_1-C_6)$ -alkyl, $-(C_1-C_6)$ -alkoxy, $-OH$, $-NR^5R^6$, $-COR^{11}$, $-NH-CONR^5R^6$, $-NH-CSNR^5R^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ;
- X is independent from R^1 , R^2 or R^3 and is selected from the group consisting of N and CH
- D is the position of attachment of the group to the rest of the nucleic acid binding compound.


whereby optionally at least one protecting group substitutes one or two hydrogen atoms of a -OH, -SH, NH₂, NH-alkyl, -NH-alkenylene, -NH-alkynylene, or a -NH-aryl group, and any salts thereof.

- 5 Most preferred in these compounds -M'R¹⁶ is a triphosphate group and -MR¹⁵ is OH. The most preferred compound is the one in which R¹⁴ is -H.

Most preferred compounds for enzymatic synthesis of a nucleic acid binding compound according to the invention are of formula VIII

10

Formula VIII




wherein

- 15 PPP is a triphosphate group, a thiotriphosphate group or analogues thereof,
 R¹⁴ is selected from the group consisting of -H, -OH, -(C₁-C₁₀)-alkoxy,
 -(C₂-C₁₀)-alkenyloxy, -(C₂-C₁₀)-alkynyloxy halogen, -azido and NH₂,
 R³⁶ is selected from the group of -H and -OH, and
 B is a group of formula I.

20

Formula I



wherein

- R¹ is independent from X, R² or R³ and is selected from the group consisting of
- (1) -F, -Cl, -Br or -I,
 - (2) Nitro
 - (3) Cyano
 - (4) -COO⁻

25

- (5) $-(C_1-C_{10})$ -alkyl substituted according to (10)
 (6) $-(C_2-C_{10})$ -alkenyl substituted according to (10)
 (7) $-(C_2-C_{10})$ -alkynyl substituted according to (10)
 (8) $-(C_6-C_{22})$ -aryl substituted according to (10)
 5 (9) $-W-(C_1-C_{10})$ -alkyl, $-W-(C_2-C_{10})$ -alkenyl, $-W-(C_2-C_{10})$ -alkynyl, $-W-(C_6-C_{22})$ -aryl or $W-H$, wherein $W = -S-, -O-, -NH-, -S-S-, -CO-, -COO-, -CO-NH-, -NH-CO-, -NH-CO-NH-, -NH-CS-NH-, -(CH_2)_n-[O-(CH_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 0 or 1 independently from r and s ,
 10 (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of $-halogen, -SH, -NO_2, -CN, -S-(C_1-C_6)$ -alkyl, $-NR^5R^6, -N^+R^5R^6R^{12}, -OR^{12}, -COR^{11}, -NH-CO-NR^5R^6, -NH-CS-NR^5R^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ,
 15 wherein R^{11} is selected from the group consisting of $-NHR^{12}, OR^{12}$, and $-SR^{12}$
 wherein R^5, R^6 and R^{12} are selected independently from the group consisting of $-H, -(C_1-C_{10})$ -alkyl, $-(C_2-C_{10})$ -alkenyl, $-(C_2-C_{10})$ -alkynyl, $-(C_6-C_{22})$ -aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,
 20 said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being unsubstituted or substituted by one or more moieties selected from the group consisting of $-halogen, -SH, -S-(C_1-C_6)$ -alkyl, $-(C_1-C_6)$ -alkoxy, $-OH, -NR^5R^6, -COR^{11}, -NH-CONR^5R^6, -NH-CSNR^5R^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ,
 25 with the proviso that R^5, R^6 or R^{12} is not a reporter group if the radicals (5) to (7) are substituted by $-NR^5R^6, NHR^{12}, OR^{12}$, or SR^{12} ;
 30 R^2, R^3 is independent from X, R^1, R^2 and R^3 and is selected from the group of,
 (1) $-H$
 (2) (C_1-C_{10}) -alkyl,
 35 (3) (C_2-C_{10}) -alkenyl,
 (4) (C_2-C_{10}) -alkynyl,

- 61 -

- (5) (C₆-C₂₂)-aryl,
 (6) -Z-(C₁-C₁₀)-alkyl, -Z-(C₂-C₁₀)-alkenyl, -Z-(C₂-C₁₀)-alkynyl, -Z-(C₆-C₂₂)-aryl or Z-H, wherein Z = -CO-, -CO-NH-, -CS-NH-, -(CH₂)_n-[O-(CH₂)_r]_s-, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s,
 (7) substituents (2) to (6)

wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -NO₂, -OR¹², -CN, -SH, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s, wherein R¹¹ is selected from the group consisting of -NHR¹² and OR¹², wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group, said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s;

25

X is independent from R¹, R² or R³ and is selected from the group consisting of N and CH

D is the position of attachment of the group to the rest of the nucleic acid binding compound.

30

whereby optionally at least one protecting group substitutes one or two hydrogen atoms of a -OH, -SH, NH₂, NH-alkyl, -NH-alkenylene, -NH-alkynylene, or a -NH-aryl group, and any salts thereof.

3' deoxy- and 2'-3'-dideoxytriphosphate subunits according to formula VIII for example can be used as terminating nucleotides in sequencing methods.

35

More preferable, above mentioned method for enzymatic synthesis uses as a triphosphate subunit a compound of formula VIII as defined above.

By the above methods, it is principally possible to introduce only one monomer containing the moiety of the invention into one nucleic acid binding component, but also more than one, as the case may be. This is especially possible using chemical methods for the synthesis of nucleic acid binding compounds.

These nucleic acid compounds according to the invention can be usefully applied in hybridization methods. Therefore, a further subject of the invention is a method for the determination of a nucleic acid comprising the steps of providing a sample suspected to contain said nucleic acid, providing a nucleic acid binding compound, which is essentially complementary to a part or all of said nucleic acid, contacting said sample with said nucleic acid binding compound under conditions for binding said nucleic acid binding compound to said nucleic acid, and determining the degree of hybridization or the binding product formed from said nucleic acid and said nucleic acid binding compound as a measure of the presence of said nucleic acid.

Methods for determination of nucleic acids by hybridization are generally known, for example from Sambrook et al. (cited above). They can easily adopted for the use of probes of the present invention.

Probes of the present invention also allow the determination of pathogens like bacteria or viruses, for example hepatitis A, B or C virus (HBV, HCV), the human immunodeficiency virus (HIV), the human papilloma virus or parvovirus B19. However, any other viruses are possible.

In a preferred embodiment of the invention, a nucleic acid binding compound, hereinafter termed a first nucleic acid binding compound, is used in a hybridization reaction to form a parallel or antiparallel duplex with a second nucleic acid binding compound wherein the first and/or the second nucleic acid binding compound comprise a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic groups, i.e. at least one of said heterocyclic groups, is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof. Preferably, the hybridization reaction is a multiplex hybridization reaction, i.e. multiple target nucleic acids as second nucleic acid binding compounds and multiple first nucleic acid binding compounds are

present. This is done preferably in the form of an array, i.e. the first nucleic acid binding compound comprises a multitude of different nucleic acid binding compounds with different sequences and is attached in the form of an array to a solid phase on different addressable locations. In a further embodiment, a nucleic acid binding compound is used
5 as a capture probe, whereby the nucleic acid binding compound has a backbone whereto heterocyclic groups capable of base pairing to nucleobases are attached characterized in that a heterocyclic group, i.e. at least one of said heterocyclic groups, is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof. Most preferred the nucleic acid binding compound is a nucleic acid binding compound with a heterocyclic group of the formula I
10 and the substituents described therefor. Most preferred are the halogen substituents in the 7-position as e.g. brom and iod.

In another preferred embodiment, the substituted pyrazolo[3,4-d]pyrimidine or the analogue thereof is used in place of a heterocyclic group in a first nucleic acid binding
15 compound to increase the melting temperature of a parallel or antiparallel duplex with a second nucleic acid binding compound whereby the increase in melting temperature is increased in comparison to the melting temperature of a duplex of the first nucleic acid binding compound with the second nucleic acid binding compound wherein the heterocyclic group in the first nucleic acid binding compound is complementary to a
20 heterocyclic group in the second nucleic acid binding compound. Preferably, a natural heterocyclic group as an adenine base is substituted in the first nucleic acid binding compound by the heterocyclic group according to the invention.

In another preferred embodiment, the substituted pyrazolo[3,4-d]pyrimidine or the
25 analogue thereof is used in place of a heterocyclic group in a first nucleic acid binding compound used as a probe in an amplification reaction, to increase the melting temperature of a duplex with a second nucleic acid binding compound in comparison to the melting temperature of a primer used in the amplification reaction, whereby the increase in melting temperature is compared to the melting temperature of a duplex of the
30 first nucleic acid binding compound with the second nucleic acid binding compound wherein the heterocyclic group in the first nucleic acid binding compound is complementary to a heterocyclic group in the second nucleic acid binding compound. Preferably, the amplification reaction is in the TaqMan® format which is described in more detail below. Preferably, a naturally occurring heterocyclic group as an adenine base is
35 substituted by the heterocyclic group according to the invention.

In another embodiment the substituted pyrazolo[3,4-d]pyrimidine or the analogue thereof is used in place of a heterocyclic group in a first nucleic acid binding compound to harmonize the contribution of each base pair to the melting temperature of a parallel or antiparallel duplex with a second nucleic acid binding compound. This is particularly
5 interesting when other non-natural compounds are present which contribute to the melting temperature in the order of the contribution of the heterocyclic group according to the invention. Then it is of interest to use the heterocyclic group according to the invention to equalize (or harmonize) the contribution of each heterocyclic group or base. This has already been described supra. This use is particularly interesting for multiplex
10 reactions and in arrays.

In a further preferred embodiment, the substituted pyrazolo[3,4-d]pyrimidine or the analogue thereof is used in place of a heterocyclic group in a first nucleic acid binding compound for enhanced detection of sequences in a second nucleic acid binding
15 compound having mismatches in a duplex with the first nucleic acid binding compound. Preferably the second nucleic acid binding compound is a target nucleic acid e.g. different subtypes of a virus. The substituted pyrazolo[3,4-d]pyrimidine or the analogue thereof can in principle be positioned anywhere in the nucleic acid binding compound. This use is of particular interest when the second nucleic acid binding compound is a target nucleic acid,
20 in particular a viral target, and different subtypes of a virus have to be amplified and detected. Therefore, in another embodiment of the invention the substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof is used in place of a heterocyclic group in a nucleic acid binding compound for enhanced detection of subtypes in a target nucleic acid.

25 In another embodiment, the substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof is used in place of a heterocyclic group in a nucleic acid binding compound to increase the melting temperature of an intramolecular duplex or hairpin of the nucleic acid binding compound whereby the increase in melting temperature is compared to the melting temperature of the intramolecular duplex of the nucleic acid binding compound wherein
30 the heterocyclic group in the nucleic acid binding compound is complementary to a heterocyclic group in the hybridizing part of the nucleic acid binding compound. This is particularly interesting in the Molecular beacons, Scorpion and TaqMan technology (WO92/02638 and corresponding US patents US 5,210,015, US 5,804,375, US 5,487,972) when two fluorescent labels have to be brought into close proximity for efficient
35 quenching.

Selecting the length of nucleic acid binding compounds or probes is also an important consideration when optimizing hybridization specificity. In general, shorter probe sequences are more specific than longer ones, in that the occurrence of a single-base mismatch has a greater destabilizing effect on the hybrid duplex. However, as the overall thermodynamic stability of hybrids decreases with length, in some embodiments it is desirable to enhance duplex stability for short probes globally. Therefore, in a further embodiment the substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof is used in place of a heterocyclic group in a nucleic acid binding compound to reduce the length of the nucleic acid binding compound in detection reactions as the heterocyclic group according to the invention has a high contribution to the melting temperature and therefore duplex stability of short probes.

Preferably, in the above described uses, at least one, preferably one or two, reporter groups are attached to the nucleic acid binding compound or a probe. Preferably, the substituted pyrazolo[3,4-d]pyrimidine or the analogue thereof used in place of a heterocyclic group is 1 to 5 nucleotides separated from the point of attachment of one or of all of the reporter groups. Preferably, the substituted pyrazolo[3,4-d]pyrimidine or the analogue thereof is used in place of an adenine in the nucleic acid binding compound.


All uses described above are preferably performed in the form of multiplex hybridization reactions, i.e. multiple target nucleic acids as second nucleic acid binding compounds and multiple first nucleic acid binding compounds are present. This is done preferably in the form of an array, i.e. the first nucleic acid binding compound comprises a multitude of different nucleic acid binding compounds with different sequences and is attached in the form of an array to a solid phase on different addressable locations.

In all the uses of the invention, the substituted pyrazolo[3,4-d]pyrimidine analogue is preferably a substituted pyrazolo[3,4-d]pyrimidine analogue of adenine or guanine or a 7-substituted pyrazolo[3,4-d]pyrimidine analogue of adenine or guanine, wherein the adenine or guanine analogues may preferably carry the same substituents R^1 in the 7-position or N-substituents R^2 and R^3 as set out directly below for the substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof or a 7-substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof. More preferably, the substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof is a substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof or a 7-substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof. Even more preferred, the substituted 7-deaza-8-aza-2,6-diamino-purine or a

derivative thereof or the 7-substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof has the formula I with the substituents as defined in the following.

- 5 In the uses described above, in the most preferred embodiment the substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof or the 7-substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof has the general formula I

Formula I



wherein

- 10 R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of
- (1) -F, -Cl, -Br or -I,
 - (2) Nitro
 - (3) Cyano
 - (4) $-\text{COO}^-$
 - 15 (5) $-(\text{C}_1-\text{C}_{10})$ -alkyl substituted according to (10)
 - (6) $-(\text{C}_2-\text{C}_{10})$ -alkenyl substituted according to (10)
 - (7) $-(\text{C}_2-\text{C}_{10})$ -alkynyl substituted according to (10)
 - (8) $-(\text{C}_6-\text{C}_{22})$ -aryl substituted according to (10)
 - (9) $-\text{W}-(\text{C}_1-\text{C}_{10})$ -alkyl, $-\text{W}-(\text{C}_2-\text{C}_{10})$ -alkenyl, $-\text{W}-(\text{C}_2-\text{C}_{10})$ -alkynyl, $-\text{W}-(\text{C}_6-$
 - 20 $\text{C}_{22})$ -aryl or $\text{W}-\text{H}$, wherein $\text{W} = -\text{S}-, -\text{O}-, -\text{NH}-, -\text{S}-\text{S}-, -\text{CO}-, -\text{COO}-, -\text{CO}-\text{NH}-, -\text{NH}-\text{CO}-, -\text{NH}-\text{CO}-\text{NH}-, -\text{NH}-\text{CS}-\text{NH}-, -(\text{CH}_2)_n-[\text{O}-(\text{CH}_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 0 or 1 independently from r and s,
 - (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be
 - 25 substituted by one or more moieties selected from the group consisting of -halogen, -SH, $-\text{NO}_2$, -CN, $-\text{S}-(\text{C}_1-\text{C}_6)$ -alkyl, $-\text{NR}^5\text{R}^6$, $-\text{N}^+\text{R}^5\text{R}^6\text{R}^{12}$, $-\text{OR}^{12}$, $-\text{COR}^{11}$, $-\text{NH}-\text{CO}-\text{NR}^5\text{R}^6$, $-\text{NH}-\text{CS}-\text{NR}^5\text{R}^6$ and $-(\text{CH}_2)_n-[\text{O}-(\text{CH}_2)_r]_s-$ NR^5R^6 , r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,
 - 30 wherein R^{11} is selected from the group consisting of $-\text{NHR}^{12}$, OR^{12} , and $-\text{SR}^{12}$

- 67 -

- wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,
- 5 said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,
- 10 with the proviso that R^5 , R^6 or R^{12} is not a reporter group if the radicals (5) to (7) are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;
- 15 R^2 , R^3 is independent from X, R^1 , R^2 and R^3 and is selected from the group of,
- (1) -H
 - (2) (C₁-C₁₀)-alkyl,
 - (3) (C₂-C₁₀)-alkenyl,
 - (4) (C₂-C₁₀)-alkynyl,
 - 20 (5) (C₆-C₂₂)-aryl,
 - (6) -Z-(C₁-C₁₀)-alkyl, -Z-(C₂-C₁₀)-alkenyl, -Z-(C₂-C₁₀)-alkynyl, -Z-(C₆-C₂₂)-aryl or Z-H, wherein Z = -CO-, -CO-NH-, -CS-NH-, -(CH₂)_n-[O-(CH₂)_r]_s-, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s,
 - 25 (7) substituents (2) to (6)
- wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -NO₂, -OR¹², -CN, -SH, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,
- 30 wherein R^{11} is selected from the group consisting of -NHR¹² and OR¹²,
- wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group,
- 35

said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s;

X is independent from R¹, R² or R³ and is selected from the group consisting of N and CH; and
 D is the position of attachment of the group to the rest of the nucleic acid binding compound
 or any salts thereof.

In the most preferred embodiment, the nucleic acid binding compound is a nucleic acid binding compound according to the invention preferably wherein R¹=Br or R¹=I.

Nucleic acid binding compounds according to the present invention also can be applied in nucleic acid determination methods in the case the nucleic acid to be determined is amplified. Since the original publication of nucleic acid amplification, various primer-based nucleic acid amplification methods have been described including, but are not limited to, Ligase Chain Reaction (LCR, Wu and Wallace, 1989, Genomics 4:560-569 and Barany, 1991, Proc. Natl. Acad. Sci. USA 88:189-193); Polymerase Ligase Chain Reaction (Barany, 1991, PCR Methods and Applic. 1:5-16); Gap-LCR (PCT Patent Publication No. WO 90/01069); Repair Chain Reaction (European Patent Publication No. 439,182 A2), 3SR (Kwoh et al., 1989, Proc. Natl. Acad. Sci. USA 86:1173-1177; Guatelli et al., 1990, Proc. Natl. Acad. Sci. USA 87:1874-1878; PCT Patent Publication No. WO 92/0880A), and NASBA (U.S. Pat. No. 5,130,238). Further, there are strand displacement amplification (SDA), transcription mediated amplification (TMA), and Q β -amplification (for a review see e.g. Whelen and Persing (1996). Annu. Rev. Microbiol. 50, 349-373; Abramson and Myers, 1993, Current Opinion in Biotechnology 4:41-47). A preferred method is the polymerase chain reaction (PCR). The invention is also related to the amplification of the target nucleic acid in the presence of the triphosphates of heterocyclic groups according to the invention as pyrazolo-[3,4-d]-pyrimidines, substituted variants thereof or analogues thereof, particularly preferred are 7-substituted variants thereof. Most preferred are the triphosphates of the heterocyclic groups according to formula I with the substituents as defined above.

The nucleic acid binding compounds according to the present invention can be used as primers and probes as e.g. as a capture probe. In the case, that the nucleic acid binding compound should be used as probe, it will preferably contain a detectable reporter group. Any hybrids formed from the nucleic acid binding compound and a nucleic acid can then
5 be determined via the detectable reporter group. This group of assays can further be divided into two groups, one being the group of homogeneous assays and the other being the heterogeneous assays. In heterogeneous assays, preferably the hybrid (binding product) will be determined when bound to a solid phase. This embodiment has the advantage that any excess of probe and other components can be removed easily from the hybrid, thus
10 make the determination easier. The hybrid formed can be captured to a solid phase either covalently, noncovalently, specifically or unspecifically. There are several embodiments which are known to a man skilled in the art.


In the so-called homogeneous assays, the hybrid formed will not be bound to a solid phase,
15 but will be determined either directly or indirectly in solution. A preferred example of such assays is disclosed in PCT/US 91/05571 which is incorporated by reference here.

In particular, when using several nucleic acid binding compounds, for example when conducting PCR-, multiplex-PCR- or multiplex-hybridization-methods it is often difficult
20 to find appropriate hybridization conditions ensuring a good specificity without losing some specific hybridization complexes resulted from a lower T_m , which also means a lower stability. In the case of diagnostic methods this can lead to false negative results, which should be avoided. A further difficulty lies in the complexity of biological samples, for example blood or sputum. Such samples often have background nucleic acids, which may
25 disturb the determination method, for example leading to false positive results.

Therefore the heterocyclic groups of formula I can also be used in multiplex hybridization methods in order to increase the T_m of one or more hybridization complexes formed in an assay. By introducing a heterocyclic group of formula I instead of a natural base contained
30 in a nucleic acid binding compound used in that assay the T_m of the hybridization complex formed with its target nucleic acid can be increased. Such changes of the T_m still allows the specific hybridization of the nucleic acid compound with its target nucleic acid at a different temperature. A preferred application field are multiplex hybridization methods on chips which often use hundreds to thousands hybridization probes.

Also included in the present invention are intermediates and precursor compounds for the chemical synthesis of the described nucleic acid binding compounds. Preferred intermediates and precursor compounds are described below.

- 5 Preferred is a solid phase bound precursor for the synthesis of a nucleic acid binding compound comprising a backbone, wherein the backbone comprises a moiety of the general formula VI



10 wherein

A is selected from the group consisting of O, S, CH₂ and N-(C₁-C₆)-alkyl,
M and M' are independently selected from the group consisting of oxy, sulfanediyl, -NR²²-, -(C₁-C₁₀)-alkyl-, or -O-(C₁-C₁₀)-alkyl-O-, and -S-(C₁-C₁₀)-alkyl-O- and -NR²²-(C₁-C₆)-alkyl-O-,

15 R²² is selected from the group of -H, -(C₁-C₁₀)-alkyl, a protecting group and a reporter group,


R¹⁴ is selected from the group consisting of -H, -OR³¹, -(C₁-C₁₀)-alkoxy, O-protecting group, S-protecting group, NH₂-protecting group, -(C₂-C₁₀)-alkenyloxy, -(C₂-C₁₀)-alkynyloxy, -halogen, -azido, SH, -(C₁-C₁₀)-alkylmercapto, and -O-solid phase,

20 R¹⁵ and R¹⁶ are independently selected from the group consisting of -H, -(C₁-C₆)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₂-C₁₀)-alkyl-carbonyl, -(C₃-C₁₉)-alkenyl-carbonyl, -(C₃-C₁₉)-alkynyl-carbonyl, -(C₆-C₁₄)-aryl-(C₁-C₁₀)-alkyl, protecting group and a solid phase

25 B is the link to a moiety of formula I,

- 71 -

Formula I




wherein

R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of

- (1) -F, -Cl, -Br or -I,
 - (2) Nitro
 - (3) Cyano
 - (4) $-\text{COO}^-$
 - (5) $-(\text{C}_1-\text{C}_{10})$ -alkyl substituted according to (10)
 - (6) $-(\text{C}_2-\text{C}_{10})$ -alkenyl substituted according to (10)
 - (7) $-(\text{C}_2-\text{C}_{10})$ -alkynyl substituted according to (10)
 - (8) $-(\text{C}_6-\text{C}_{22})$ -aryl substituted according to (10)
 - (9) $-\text{W}-(\text{C}_1-\text{C}_{10})$ -alkyl, $-\text{W}-(\text{C}_2-\text{C}_{10})$ -alkenyl, $-\text{W}-(\text{C}_2-\text{C}_{10})$ -alkynyl, $-\text{W}-(\text{C}_6-\text{C}_{22})$ -aryl or W-H, wherein W = -S-, -O-, -NH-, -S-S-, -CO-, -COO-, -CO-NH-, -NH-CO-, -NH-CO-NH-, -NH-CS-NH-, $-(\text{CH}_2)_n-[\text{O}-(\text{CH}_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 0 or 1 independently from r and s,
 - (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -SH, $-\text{NO}_2$, -CN, $-\text{S}-(\text{C}_1-\text{C}_6)$ -alkyl, $-\text{NR}^5\text{R}^6$, $-\text{N}^+\text{R}^5\text{R}^6\text{R}^{12}$, $-\text{OR}^{12}$, $-\text{COR}^{11}$, $-\text{NH}-\text{CO}-\text{NR}^5\text{R}^6$, $-\text{NH}-\text{CS}-\text{NR}^5\text{R}^6$ and $-(\text{CH}_2)_n-[\text{O}-(\text{CH}_2)_r]_s-\text{NR}^5\text{R}^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s, wherein R^{11} is selected from the group consisting of $-\text{NHR}^{12}$, OR^{12} , and $-\text{SR}^{12}$
- wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of -H, $-(\text{C}_1-\text{C}_{10})$ -alkyl, $-(\text{C}_2-\text{C}_{10})$ -alkenyl, $-(\text{C}_2-\text{C}_{10})$ -alkynyl, $-(\text{C}_6-\text{C}_{22})$ -aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,

(11)

- 72 -



wherein Nuc is the link to formula I and n is any integer from 1 to 18
 R^2, R^3 is independent from X, R^1 , R^2 and R^3 and is selected from the group of,

- (1) -H
- (2) $(\text{C}_1-\text{C}_{10})$ -alkyl,
- (3) $(\text{C}_2-\text{C}_{10})$ -alkenyl,
- (4) $(\text{C}_2-\text{C}_{10})$ -alkynyl,
- (5) $(\text{C}_6-\text{C}_{22})$ -aryl,
- (6) -Z- $(\text{C}_1-\text{C}_{10})$ -alkyl, -Z- $(\text{C}_2-\text{C}_{10})$ -alkenyl, -Z- $(\text{C}_2-\text{C}_{10})$ -alkynyl, -Z- $(\text{C}_6-\text{C}_{22})$ -aryl or Z-H, wherein Z = -CO-, -CO-NH-, -CS-NH-, $-(\text{CH}_2)_n-$, $[\text{O}-(\text{CH}_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s,
- (7) substituents (2) to (6)

wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, - NO_2 , - OR^{12} , -CN, -SH, -S- (C_1-C_6) -alkyl, - NR^5R^6 , - $\text{N}^+\text{R}^5\text{R}^6\text{R}^{12}$, - COR^{11} , -NH- CONR^5R^6 , -NH-CSNR⁵R⁶ and $-(\text{CH}_2)_n-[\text{O}-(\text{CH}_2)_r]_s-\text{NR}^5\text{R}^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,

wherein R^{11} is selected from the group consisting of -NHR¹² and OR^{12} ,

wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of -H, $(\text{C}_1-\text{C}_{10})$ -alkyl, $(\text{C}_2-\text{C}_{10})$ -alkenyl, $(\text{C}_2-\text{C}_{10})$ -alkynyl, $(\text{C}_6-\text{C}_{22})$ -aryl and a reporter group,


- 73 -

said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s;

X is independent from R¹, R² or R³ and is selected from the group consisting of N and CH
D is the position of attachment of the group to the rest of the nucleic acid binding compound.

with the proviso that one or two hydrogen atoms of any -OH, -SH, -NH₂, -NH-alkyl, -NH-alkenylene, -NH-alkynylene, or -NH-aryl group are substituted by a protecting group, and any salts thereof. Such compounds of Formula VI can be used for chemical synthesis of nucleic acid binding compounds according to the invention as precursors. In this case the compounds are linked to a solid phase, preferred R¹⁴, R¹⁵, or R¹⁶ is O-solid phase, most preferred R¹⁵ is solid phase. It is also preferred that reactive groups are protected by protective groups.

Also included in the present invention are precursors and intermediates of a nucleic acid binding compound, wherein the backbone comprises a moiety of the general formula III




25

wherein

A is selected from the group consisting of O, S, CH₂ and N-(C₁-C₆)-alkyl,

- M is selected from the group consisting of oxy, sulfanediyl, $-NR^{22}$ -, $-(C_1-C_{10})$ -alkyl-, or $-O-(C_1-C_{10})$ -alkyl-O-, and $-S-(C_1-C_{10})$ -alkyl-O-, $O-CO$ -, $-NR^{22}-(C_1-C_6)$ -alkyl-O-,
- 5 R^{22} is selected from the group of $-H$ -, $-(C_1-C_{10})$ -alkyl, a protecting group and a reporter group,
- R^{14} is selected from the group consisting of $-H$ -, $-OH$ -, $-(C_1-C_{10})$ -alkoxy, $-(C_2-C_{10})$ -alkenyloxy, $-(C_2-C_{10})$ -alkynyloxy, -halogen, -azido, SH -, $-(C_1-C_{10})$ -alkylmercapto, O-reporter group, O-solid phase and $-NH_2$ linked to a protecting group,
- 10 R^{15} is selected from the group consisting of $-H$ -, $-(C_1-C_6)$ -alkyl, $-(C_2-C_{10})$ -alkenyl, $-(C_2-C_{10})$ -alkynyl, $-(C_2-C_{10})$ -alkyl-carbonyl, $-(C_3-C_{19})$ -alkenyl-carbonyl, $-(C_3-C_{19})$ -alkynyl-carbonyl, $-(C_6-C_{14})$ -aryl- $-(C_1-C_{10})$ -alkyl and a solid phase,
- B is the link to a moiety of formula I,

Formula I



15


wherein

- R^1 is independent from X , R^2 or R^3 and is selected from the group consisting of
- (1) $-F$ -, $-Cl$ -, $-Br$ or $-I$,
 - 20 (2) Nitro
 - (3) Cyano
 - (4) $-COO^-$
 - (5) $-(C_1-C_{10})$ -alkyl substituted according to (10)
 - (6) $-(C_2-C_{10})$ -alkenyl substituted according to (10)
 - 25 (7) $-(C_2-C_{10})$ -alkynyl substituted according to (10)
 - (8) $-(C_6-C_{22})$ -aryl substituted according to (10)
 - (9) $-W-(C_1-C_{10})$ -alkyl, $-W-(C_2-C_{10})$ -alkenyl, $-W-(C_2-C_{10})$ -alkynyl, $-W-(C_6-C_{22})$ -aryl or $W-H$, wherein $W = -S-$, $-O-$, $-NH-$, $-S-S-$, $-CO-$, $-COO-$, $-CO-NH-$, $-NH-CO-$, $-NH-CO-NH-$, $-NH-CS-NH-$, $-(CH_2)_n-[O-(CH_2)_r]_s-$, where r and s are, independently of each other, an integer
 - 30 between 1 to 18 and n is 0 or 1 independently from r and s ,

- 75 -

- (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -SH, -NO₂, -CN, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -OR¹², -COR¹¹, -NH-CO-NR⁵R⁶, -NH-CS-NR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s, wherein R¹¹ is selected from the group consisting of -NHR¹², OR¹², and -SR¹² wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,

(11)



wherein Nuc is the link to formula I and n is any integer from 1 to 18 said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s, with the proviso that R⁵, R⁶ or R¹² is not a reporter group if the radicals (5) to (7) are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²,

- 76 -

- R^2, R^3 is independent from X, R^1, R^2 and R^3 and is selected from the group of,
- (1) $-H$
 - (2) (C_1-C_{10}) -alkyl,
 - (3) (C_2-C_{10}) -alkenyl,
 - 5 (4) (C_2-C_{10}) -alkynyl,
 - (5) (C_6-C_{22}) -aryl,
 - (6) $-Z-(C_1-C_{10})$ -alkyl, $-Z-(C_2-C_{10})$ -alkenyl, $-Z-(C_2-C_{10})$ -alkynyl, $-Z-(C_6-C_{22})$ -aryl or $Z-H$, wherein $Z = -CO-, -CO-NH-, -CS-NH-, -(CH_2)_n-[O-(CH_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s ,
 - 10 (7) substituents (2) to (6)
- wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of $-halogen, -NO_2, -OR^{12}, -CN, -SH, -S-(C_1-C_6)$ -alkyl, $-NR^5R^6, -N^+R^5R^6R^{12}, -COR^{11}, -NH-CONR^5R^6, -NH-CSNR^5R^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ,
- wherein R^{11} is selected from the group consisting of $-NHR^{12}$ and OR^{12} ,
- 20 wherein R^5, R^6 and R^{12} are selected independently from the group consisting of $-H, -(C_1-C_{10})$ -alkyl, $-(C_2-C_{10})$ -alkenyl, $-(C_2-C_{10})$ -alkynyl, $-(C_6-C_{22})$ -aryl and a reporter group,
- said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of $-halogen, -SH, -S-(C_1-C_6)$ -alkyl, $-(C_1-C_6)$ -alkoxy, $-OH, -NR^5R^6, -COR^{11}, -NH-CONR^5R^6, -NH-CSNR^5R^6$ and $-(CH_2)_n-[O-(CH_2)_r]_s-NR^5R^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ;
- 25
- 30 X is independent from R^1, R^2 or R^3 and is selected from the group consisting of N and CH
- D is the position of attachment of the group to the rest of the nucleic acid binding compound
- with the proviso that one or two hydrogen atoms of any $-OH, -SH, -NH_2, -NH$ -alkyl, $-NH$ -alkenylene, $-NH$ -alkynylene, or $-NH$ -aryl group are substituted by a protecting group, and any salts thereof,
- 35


wherein R^{14} is O-solid phase or R^{15} is solid phase.

Solid phases for chemical synthesis of a nucleic acid binding compound according to the invention preferably also include linkers to fix the growing nucleic acid binding compound.

- 5 Such linkers are known in the art. Preferably such linkers can be cleaved after synthesis to free said nucleic acid binding compound and can for example also be used to generate a free 3'-hydroxy group in said nucleic acid binding compound. Such linkers are known in the art, for example succinic acid linked via an amide bond to the solid phase and via an ester to the precursor or intermediate. Preferred R^{15} is solid phase, but in the precursor for
- 10 chemical synthesis of a nucleic acid binding compound according to formula III alternatively R^{14} may also be solid phase. Reactive groups of said compound are preferably protected by a protective group.

- A more general formula of preferred precursors and intermediates according to the present
- 15 invention are compounds comprising a backbone, said backbone having attached heterocyclic groups characterized in that a heterocyclic group is a group of the general formula I

Formula I



- R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of
- 20 (1) -F, -Cl, -Br or -I,
 (2) Nitro
 (3) Cyano
 (4) $-\text{COO}^-$
 (5) $-(\text{C}_1-\text{C}_{10})$ -alkyl substituted according to (10)
 25 (6) $-(\text{C}_2-\text{C}_{10})$ -alkenyl substituted according to (10)
 (7) $-(\text{C}_2-\text{C}_{10})$ -alkynyl substituted according to (10)
 (8) $-(\text{C}_6-\text{C}_{22})$ -aryl substituted according to (10)
 (9) $-\text{W}-(\text{C}_1-\text{C}_{10})$ -alkyl, $-\text{W}-(\text{C}_2-\text{C}_{10})$ -alkenyl, $-\text{W}-(\text{C}_2-\text{C}_{10})$ -alkynyl, $-\text{W}-(\text{C}_6-\text{C}_{22})$ -aryl or W-H, wherein W= -S-, -O-, -NH-, -S-S-, -CO-, -COO-, -CO-NH-, -NH-CO-, -NH-CO-NH-, -NH-CS-NH-, $-(\text{CH}_2)_n$ -[O-
- 30

$(\text{CH}_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 0 or 1 independently from r and s ,

- (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of
- 5 -halogen, -SH, -NO₂, -CN, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -OR¹², -COR¹¹, -NH-CO-NR⁵R⁶, -NH-CS-NR⁵R⁶ and $-(\text{CH}_2)_n-[\text{O}-(\text{CH}_2)_r]_s-$ NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ,

wherein R¹¹ is selected from the group consisting of -NHR¹², OR¹², and -SR¹²

wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound

15 hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,

said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and $-(\text{CH}_2)_n-[\text{O}-(\text{CH}_2)_r]_s-$ NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ,

with the proviso that R⁵, R⁶ or R¹² is not a reporter group if the radicals (5) to (7) are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;

25 R², R³ is independent from X, R¹, R² and R³ and is selected from the group of,

- (1) -H
- (2) (C₁-C₁₀)-alkyl,
- (3) (C₂-C₁₀)-alkenyl,
- (4) (C₂-C₁₀)-alkynyl,
- 30 (5) (C₆-C₂₂)-aryl,
- (6) -Z-(C₁-C₁₀)-alkyl, -Z-(C₂-C₁₀)-alkenyl, -Z-(C₂-C₁₀)-alkynyl, -Z-(C₆-C₂₂)-aryl or Z-H, wherein Z = -CO-, -CO-NH-, -CS-NH-, $-(\text{CH}_2)_n-[\text{O}-(\text{CH}_2)_r]_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s ,

35 (7) substituents (2) to (6)

- 79 -

- wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -NO₂, -OR¹², -CN, -SH, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,
- wherein R¹¹ is selected from the group consisting of -NHR¹² and OR¹²,
- wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group,
- said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s;
- X is independent from R¹, R² or R³ and is selected from the group consisting of N and CH
- D is the position of attachment of the group to the rest of the nucleic acid binding compound.
- wherein said backbone is solid phase bound;
- with the proviso that one or two hydrogen atoms of any -OH, -SH, -NH₂, -NH-alkyl, -NH-alkenylene, -NH-alkynylene, or -NH-aryl group are substituted by a protecting group, and any salts thereof.
- Beside the possibility that the precursor compound or intermediate is coupled to the solid phase at the backbone, it can also be linked at the heterocyclic group of formula I included in said compound, for example using a -OH, -SH or -NH₂ groups as attachment site. Preferably the other reactive groups of said compound are protected by protective groups.
- A preferred embodiment of the invention is a method for the determination of the presence, absence or amount of a nucleic acid comprising the steps of providing a sample

suspected to contain the nucleic acid, providing a nucleic acid binding compound comprising a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof, which is essentially
5 complementary to a part or all of the nucleic acid, contacting said sample with the nucleic acid binding compound under conditions for binding the nucleic acid binding compound to the nucleic acid, determining the binding product or the degree of hybridization between the nucleic acid and the nucleic acid binding compound as a measure of the presence, absence or amount of the nucleic acid.

10

A further embodiment of the invention is a method for the determination of the presence, absence or amount of a nucleic acid wherein a nucleic acid binding compound is used as a capture probe, wherein the nucleic acid binding compound comprises a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases
15 characterized in that a heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof.

Yet another embodiment of the invention is a method for distinguishing related nucleotide sequences in a nucleic acid, the method comprising the steps of providing a nucleic acid
20 binding compound comprising a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof and having a defined sequence, providing a nucleic acid with two related nucleotide sequences, each of which comprises a target sequence, wherein one of the nucleotide sequence is a target sequence
25 that is perfectly complementary to the nucleic acid binding compound and at least one other of the segments is a related target sequence, incubating the nucleic acid with the nucleic acid binding compound under hybridization conditions, and determining the degree of hybridization between the nucleic acid binding compound and each of the segments.

30

Preferably any substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof is located in said compound to pair with dT and to increase the melting point of the nucleic binding compound hybridized to its complementary nucleic acid. The expert in the field is aware of the fact that the increase of the melting point of a nucleic acid binding compound
35 according to the invention is influenced by the environment of the heterocyclic group

according to the invention. Preferably however, the melting point is 4.5 to 7 °C higher, preferably 5 or 5.5 to 7 °C, more preferably 6 to 7 °C than the T_m of a dA-dT pair.

In a very preferred embodiment, in the methods according to the invention the nucleic acids are isolated from biological material, preferably from a human or an animal. Preferably the methods according to the invention are used in the diagnostical field.

In still another embodiment of the invention a method as described above is contemplated wherein the nucleic acid binding compound according to the invention comprises a reporter group which is a fluorescent label, preferably fluorescein. Preferably, the nucleic acid binding compound according to the invention comprises multiple fluorescent labels wherein the emission wavelengths of one of the fluorescent labels overlaps the absorption wavelengths of another of the fluorescent labels. The nucleic acid binding compound may further comprise a quenching agent which quenches the fluorescence emission of the fluorescent label, which can be fluorescein. Preferably the quenching agent is a fluorescent rhodamine or cyanine. Preferably the method further comprises the step of altering the spatial relationship between the fluorescent label and the quenching agent subsequent to hybridization, preferably by exonuclease hydrolysis of the nucleic acid binding compound whereby release of label occurs as a result of exonuclease hydrolysis. In a preferred embodiment, the degree of hybridization between the nucleic acid binding compound and the nucleic acid is determined by the quantity of label that is released from the nucleic acid binding compound subsequent to hybridization.

In a preferred embodiment of the invention, a method for distinguishing related nucleotide sequences is disclosed, wherein the related sequences preferably differ by a single nucleotide. Preferably, the degree of hybridization between the nucleic acid binding compound and the nucleic acid is determined by the priming ability of the nucleic acid binding compound, wherein most preferably priming occurs as part of an amplification reaction which may be an amplification reaction described above. The amplification reaction is preferably a polymerase chain reaction.

In methods for the determination of the presence, absence or amount of a nucleic acid or the method for distinguishing related nucleotide sequences more than one nucleic acid binding compound may be used, wherein the nucleic acid binding compound comprises a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic group is a substituted pyrazolo [3,4-

d]pyrimidine or an analogue thereof. Preferably, two nucleic acid binding compounds are used. In a preferred embodiment the first of the two nucleic acid binding compounds comprises a fluorescence donor and the second of the two nucleic acid binding compounds comprises a fluorescence acceptor, wherein the emission wavelengths of the fluorescence donor overlap the absorption wavelengths of the fluorescence acceptor. Then, the degree of hybridization can be measured by the quantity of light transferred between the fluorescence donor and the fluorescence acceptor and emitted by the fluorescence acceptor. In another embodiment the degree of hybridization is determined by the measurement of the melting temperature between the nucleic acid binding compound and the nucleic acid.

10 In yet another embodiment of the invention, a method for detecting the presence of a target sequence in a nucleic acid is disclosed, the method comprising the steps of providing a nucleic acid which is to be tested for the presence of the target sequence, providing a nucleic acid binding compound having a sequence that is substantially complementary to the target sequence and comprising a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic groups is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof, incubating the nucleic acid and the nucleic acid binding compound under hybridization conditions; and identifying hybridized nucleic acids. Preferably, multiple nucleic acids are tested for the presence of the target sequence, whereby the nucleic acids have related target sequences. Most preferably, the nucleic acids differ from one another by a single nucleotide within the target sequence. Preferably, the nucleic acid binding compound is a primer comprising an extendible 3'-hydroxyl group. In a preferred embodiment, the hybridized nucleic acids are identified by extending the primer with a polymerizing enzyme, which can be a thermostable enzyme and wherein the nucleic acid binding compound is a primer in an amplification reaction, preferably a polymerase chain reaction. Preferably the thermostable enzyme is the DNA polymerase from *Thermus aquaticus*, the so-called Taq-Polymerase.

30 In still another embodiment of the invention, a method for primer extension is disclosed which comprises the steps of providing a nucleic acid containing a target sequence, providing one or more nucleic acid binding compounds complementary to the target sequence, wherein nucleic acid binding compound comprises a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof, providing a polymerizing enzyme and nucleotide substrates, and incubating the nucleic acid, the nucleic acid binding compounds, the enzyme and the substrates under conditions

favorable for polymerization. Preferably, the method is part of an amplification reaction, most preferably a polymerase chain reaction. The method can be used in the synthesis of a cDNA molecule.

- 5 Another embodiment of the invention is a method for determining the nucleotide sequence of a nucleic acid, the method comprising the steps of providing an array of nucleic acid binding compounds having different known sequences and comprising a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic group is a substituted pyrazolo[3,4-
- 10 d]pyrimidine or an analogue thereof, with the proviso that the nucleic acid binding compounds do not contain a reporter group, incubating the nucleic acid with the array under hybridization conditions, and determining to which of the nucleic acid binding compounds in the array the nucleic acid hybridizes.
- 15 Still another embodiment of the invention is a method for determining the nucleotide sequence of a target sequence in a nucleic acid, the method comprising the steps of providing a nucleic acid comprising the target sequence, providing at least two nucleic acid binding compounds with a known sequence comprising a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a
- 20 heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof, and wherein one of the at least two nucleic acid binding compounds has a sequence that is perfectly complementary to the target sequence and at least one other of the nucleic acid binding compounds has a related target sequence, incubating the nucleic acid binding compounds with the nucleic acid under hybridization conditions, and determining the
- 25 degree of hybridization between each of the nucleic acid binding compounds and the nucleic acid. Preferably, the at least one other nucleic acid binding compounds has a single-nucleotide mismatch with the target sequence.
- Further, the invention contemplates a method for examining gene expression in a cell, the
- 30 method comprising the steps of providing a population of nucleic acids representative of the genes expressed in the cell, providing an array of nucleic acid binding compounds comprising a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof, with the proviso that the nucleic acid
- 35 binding compounds do not contain a reporter group, incubating the population of nucleic acids with the array under hybridization conditions, and determining which of the nucleic


acid binding compounds in the array become hybridized to nucleic acids which is optionally labelled by incorporation of e.g. of adenosinetriphosphate coupled to a label.

5 Still another embodiment of the invention is a method for identifying a mutation in a target sequence of a gene of interest, the method comprising the steps of providing a nucleic acid that comprises the target sequence, providing an array of nucleic acid binding compounds of different sequences, wherein the different sequences include the wild-type target sequence and different mutant target sequences, wherein the nucleic acid binding compounds comprise a backbone, said backbone having attached heterocyclic groups
10 capable of base pairing to nucleobases characterized in that a heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof with the proviso that the nucleic acid binding compounds do not contain a reporter group, incubating the nucleic acid with the array under hybridization conditions, and determining which of the nucleic acid binding compounds in the array become hybridized to the nucleic acid.

15 In all methods presented above the substituted pyrazolo[3,4-d]pyrimidine analogue may be preferably the substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof is a substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof or a 7-substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof. In another embodiment of the
20 invention the substituted pyrazolo[3,4-d]pyrimidine analogue is a substituted pyrazolo[3,4-d]pyrimidine analogue of adenine or guanine or a 7-substituted pyrazolo[3,4-d]pyrimidine analogue of adenine or guanine, wherein the adenine or guanine analogues may preferably carry the same substituents R^1 in the 7-position or N-substituents R^2 and R^3 as set out directly below for the substituted 7-deaza-8-aza-2,6-diamino-purine or a
25 derivative thereof or a 7-substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof.

In a preferred embodiment the substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof or the 7-substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative
30 thereof has the general formula I

Formula I



wherein

R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of

- (1) -F, -Cl, -Br or -I,
 - (2) Nitro
 - (3) Cyano
 - (4) $-\text{COO}^-$
 - (5) $-(\text{C}_1\text{-C}_{10})$ -alkyl substituted according to (10)
 - (6) $-(\text{C}_2\text{-C}_{10})$ -alkenyl substituted according to (10)
 - (7) $-(\text{C}_2\text{-C}_{10})$ -alkynyl substituted according to (10)
 - (8) $-(\text{C}_6\text{-C}_{22})$ -aryl substituted according to (10)
 - (9) $-\text{W}-(\text{C}_1\text{-C}_{10})$ -alkyl, $-\text{W}-(\text{C}_2\text{-C}_{10})$ -alkenyl, $-\text{W}-(\text{C}_2\text{-C}_{10})$ -alkynyl, $-\text{W}-(\text{C}_6\text{-C}_{22})$ -aryl or W-H, wherein W = -S-, -O-, -NH-, -S-S-, -CO-, $-\text{COO}^-$, -CO-NH-, -NH-CO-, -NH-CO-NH-, -NH-CS-NH-, $-(\text{CH}_2)_n\text{-[O-(CH}_2)_r\text{]}_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 0 or 1 independently from r and s,
 - (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -SH, $-\text{NO}_2$, -CN, $-\text{S}-(\text{C}_1\text{-C}_6)$ -alkyl, $-\text{NR}^5\text{R}^6$, $-\text{N}^+\text{R}^5\text{R}^6\text{R}^{12}$, $-\text{OR}^{12}$, $-\text{COR}^{11}$, $-\text{NH-CO-NR}^5\text{R}^6$, $-\text{NH-CS-NR}^5\text{R}^6$ and $-(\text{CH}_2)_n\text{-[O-(CH}_2)_r\text{]}_s-$ NR^5R^6 , r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s, wherein R^{11} is selected from the group consisting of $-\text{NHR}^{12}$, OR^{12} , and $-\text{SR}^{12}$
- wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of -H, $-(\text{C}_1\text{-C}_{10})$ -alkyl, $-(\text{C}_2\text{-C}_{10})$ -alkenyl, $-(\text{C}_2\text{-C}_{10})$ -alkynyl, $-(\text{C}_6\text{-C}_{22})$ -aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,

- said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,
- with the proviso that R⁵, R⁶ or R¹² is not a reporter group if the radicals (5) to (7) are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;
- R², R³ is independent from X, R¹, R² and R³ and is selected from the group of,
- (1) -H
 - (2) (C₁-C₁₀)-alkyl,
 - (3) (C₂-C₁₀)-alkenyl,
 - (4) (C₂-C₁₀)-alkynyl,
 - (5) (C₆-C₂₂)-aryl,
 - (6) -Z-(C₁-C₁₀)-alkyl, -Z-(C₂-C₁₀)-alkenyl, -Z-(C₂-C₁₀)-alkynyl, -Z-(C₆-C₂₂)-aryl or Z-H, wherein Z = -CO-, -CO-NH-, -CS-NH-, -(CH₂)_n-[O-(CH₂)_r]_s-, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s,
 - (7) substituents (2) to (6)
- wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -NO₂, -OR¹², -CN, -SH, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,
- wherein R¹¹ is selected from the group consisting of -NHR¹² and OR¹²,
- wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group,
- said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s;

- 87 -

X is independent from R¹, R² or R³ and is selected from the group consisting of N and CH; and

D is the position of attachment of the group to the rest of the nucleic acid binding compound

5

or any salts thereof.

In still another preferred embodiment, the nucleic acid binding compound is a nucleic acid binding compound according to the invention, preferably R¹ is a hydrophobic or electron-withdrawing substituent as defined previously, preferably a halogen substituent whereby Br or I is most preferred.

10

In all the methods or uses of the present invention the nucleic acid binding compound having incorporated a heterocyclic group according to the present invention can bind to an opposite strand to form a parallel or antiparallel duplex.

15

The invention furthermore relates to pharmaceutical compositions comprising one or more nucleic acid binding compounds containing a heterocyclic group of formula I, together with a physiologically acceptable excipients and, where appropriate, suitable additives and/or conventional auxiliary substances. Therefore, one embodiment of the invention is a pharmaceutical composition comprising a nucleic acid binding compound according to the invention. Another embodiment is a nucleic acid binding compound according to the invention for use in medicine. In a quite general manner, the present invention extends to the use of such nucleic acid binding compounds in therapeutically effective compositions. Such compositions are understood to mean to include the nucleic acid binding compounds according to the invention as antisense oligonucleotides, triple helix forming oligonucleotides, aptamers or ribozymes, in particular antisense oligonucleotides.

20

25

The invention further contemplates protecting groups derived from phthalic acid (see e.g. Ramzaeva et al. (1999). Nucleosides & Nucleotides 18, 1439-1440; Rich B. Meyer in Methods in Molecular Biology 26 (1994), pp. 73-91, Humana Press Inc. Totowa, NJ, USA; Griffey et al. (1996). J. Med. Chem. 39, 5100-5109; Gibson and Benkovic (1987). Nucl. Acids Res. 15, 6455-6467). Aminoalkinyl or aminoalkenyl side chains of the pyrazolo[3,4-d]pyrimidin analogues of desoxyadenosine, desoxy-guanosine or desoxy-isoguanosine may be protected with phthalic whereby an imide is formed. In contrast to the trifluoroacetyl protecting group which may be removed relatively easy from an oligonucleotide, the


30

35

- 88 -

removal of a phthaloyl group from aliphatic side chains is not easy as e.g. for peptides. Therefore, hydrazine would have to be used which cannot however be used in the case of oligonucleotides, as pyrimidine nucleosides are degraded. The amino groups of alkynyl compounds, in particular of propargylamine as well as those which are in allyl position on other positions in the chain are less basic. Thereby, a phthaloyl group may be more easily removed and no methyl amine has to be used. The deprotection is performed under standard conditions, i.e. concentrated ammonia solution at 60 °C, 12 hours. The advantage of the phthaloyl group is that both H-atom positions of the amino group are acylated. For normal acyl groups as e.g. the trifluoroacetyl (TFA) group used in US 5,151,507, the amino group is capped by acylation and is monofunctionalized thereafter. Later the TFA group is removed whereby the acetyl group remains bound and is difficult to remove. The advantage of the phthaloyl group for such purposes is obvious.

Hence, in one embodiment of the invention a building block for the synthesis of an oligonucleotide comprising the nucleosides adenosine or desoxyadenosine, guanosine or desoxyguanosine, isoguanosine or desoxyisoguanosine, cytidine or deoxycytidine, uridine or desoxyuridine, thymidine or desoxythymidine, or analogues of these nucleosides wherein a substituent is attached to the base moiety selected from the group of phthalimidoalkyl, phthalimidoalkenyl, or phthalimidoalkynyl groups with the following formulas



wherein Nuc is the position of attachment of the substituent to the base moiety and n is any integer from 1 to 18, with the proviso that the phthalimidoalkynyl or phthalimidoalkyl group is not attached to the C5-atom of deoxyuridine and the proviso that the phthalimidoalkynyl group is not attached to the C7-atom of 7-deaza-deoxyguanosine. In a

preferred embodiment the analogue of these nucleosides is a pyrazolo[3,4-d]pyrimidine analogue. In a further preferred embodiment, the building block according to the invention is a phosphoramidite derivative.

- Further the invention contemplates the use of a building block according to the invention
- 5 for the synthesis of an oligonucleotide. Further, the use of a phthaloyl group as a protecting group is contemplated in a method for the synthesis of an oligonucleotide from building blocks comprising the nucleosides adenosine or desoxyadenosine, guanosine or desoxyguanosine, isoguanosine or desoxyisoguanosine, cytidine or deoxycytidine, uridine or desoxyuridine, thymidine or desoxythymidine, or analogues of these nucleosides
- 10 wherein the amino groups of $-(CH_2)_n-NH_2$, $-CH=CH-(CH_2)_n-NH_2$, or $-C\equiv C-(CH_2)_n-NH_2$ attached to the base moiety are derivatized with the phthaloyl group and n is any integer from 1 to 18, with the proviso that the $-(CH_2)_n-NH_2$ or $-C\equiv C-(CH_2)_n-NH_2$ group is not attached to the C5-atom of deoxyuridine and the proviso that the $-C\equiv C-(CH_2)_n-NH_2$ group is not attached to the C7-atom of 7-deaza-deoxyguanosine. In a preferred
- 15 embodiment the analogue of these nucleosides is a pyrazolo[3,4-d]pyrimidine analogue. In a further preferred embodiment, the building block is a phosphoramidite derivative. In yet another embodiment of the invention, a method for the synthesis of an oligonucleotide from building blocks according to the invention is disclosed.
- 20 These examples are intended to illustrate possible applications and should not limit the scope of the invention.

The present invention is explained in more detail by the following examples:

Examples

5 Example 1: Synthesis and data on the 7-bromo-8aza-7-deazapurin-2,6-diamine nucleosides

1.1. Synthesis and Properties of Monomers

- The alkoxy nucleosides 4a,b [18] served as precursor for the synthesis of the 8-aza-7-deazapurin-2,6-diamine (pyrazolo[3,4-d]pyrimidin- 4,6-diamine) nucleosides 2a,b (purine numbering is used throughout the discussion section) The amination was performed in a steel bomb (4 days, 25% aq. NH₃, 70°). Both nucleosides (2a,b) were isolated crystalline. A few related 8-aza-7-deazapurin-2,6-diamine nucleosides have been prepared earlier [19 - 22].


15

Treatment of compound 2a with adenosine deaminase (ADA) resulted in the formation of 8-aza-7-deaza-2'-deoxyguanosine [18]. The reaction was followed UV-spectrophotometrically (Figure 1a).

20 The time-dependent spectra show two isosbestic points ($\lambda = 234$ nm and 267 nm) indicating the conversion of the starting material into only one reaction product. The deamination of 2a occurs much slower than that of the purine nucleoside 1. The 7-bromo derivative 2b [18] was not deaminated under these conditions even at high enzyme concentrations as it was observed earlier [20].

25 Scheme 1

- 91 -



Next, the half-lives of the N-glycosylic bonds of compounds 2a,b in acidic medium were measured and compared with that of compound 1. The reaction was performed in 0.5 N HCl at room temperature and was followed UV-spectrophotometrically as well as by HPLC-analysis (Table1).

Table 1. Half-life Values (τ) of 2'-Deoxyadenosine Derivatives in HCl at 25°.

Compound	τ [min] ^{a)}	λ (nm)	Compound	τ [min] ^{a)}	λ (nm)
1	6	252	2b	Stable ^{b)}	242
2a	91	255	2b	87 ^{c)}	242

^{a)} Measured in 0.5 N HCl. ^{b)} Within 4 h. ^{c)} Measured in 2 N HCl.

^{d)} Determined by HPLC.

- From the data of the table it is apparent that the 8-aza-7-deazapurin-2,6-diamine nucleoside (2a) shows an about 15-fold higher glycosylic bond stability than the parent 2-amino-2'-deoxyadenosine (1). The glycosylic bond of 2b is stable under these conditions. Hydrolysis occurs when the reaction is performed in 2 N HCl (Table 1). Furthermore, the 7-bromo substituent increases the lipophilic character of the molecule thereby decreasing the chromatographic mobility of compound 2b, in comparison to compounds 2a or 1 on reverse phase HPLC (Figure 1b).

In order to study the influence of the nucleobases on the N \leftrightarrow S pseudorotational equilibrium of the sugar moiety, the ^1H -NMR spectra of the nucleosides 1 – 3 were measured in D_2O . The analysis was performed on the basis of five vicinal ^1H , ^1H coupling

constants using the PSEUROT program [23]. According to Table 2 the 8-aza-7-deazapurin-2,6-diamine nucleoside 2a shows a more pronounced N-conformer population than the corresponding purine nucleoside 1, while the N-type population of the related 7-deazapurine nucleoside 3 is decreased. This is in line with an increase of the π -electron deficiency of the 8-aza-7-deazapurine system. An additional effect of the electron-withdrawing 7-bromo substituent is not observed (2b) but was found for the more electron-attracting cyano group introduced into the 7-position of 8-aza-7-deaza-2'-deoxyguanosine [24]. The conformation around the C(4')-C(5') bond indicates that the 8-aza-7-deazapurin-2,6-diamine nucleosides 2a and b as the 8-aza-7-deaza-2'-deoxyguanosines [24], prefer the γ^t -(-sc)-rotamer population, while for the regular purine nucleosides the $\gamma^{(+)\text{g}}$ -(+sc)- or the $\gamma^{(-)\text{g}}$ -(ap)-conformation is predominant [25].

Table 2. 3J (H, H) Coupling Constants of the Sugar Moieties and N/S-Conformer Populations of the 2'-Deoxyribonucleosides 1 – 3 at 303 K. ^{a)}

	$^3J_{\text{H,H}}/\text{Hz}$							Conformation				
	1',2'	1',2''	2',3'	2'',3'	3',4'	4',5'	4',5''	%N	%S	$\gamma^{(+)\text{g}}$	γ^t	$\gamma^{(-)\text{g}}$
1	7.30	6.10	7.00	3.10	3.40	3.20	4.30	31	69	62	25	13
2a	6.60	6.80	6.90	3.70	3.60	4.00	5.80	37	63	36	42	22
2b	6.60	6.70	6.80	3.70	3.90	4.30	5.90	37	63	32	43	25
3	7.90	6.40	6.20	3.10	3.00	3.87	4.82	25	75	48	32	20

^{a)} Solvent D₂O; r.m.s. < 0.4 Hz; $|\Delta J_{\text{max}}| < 0.4$ Hz.


15

In the past several laboratories have reported on a straightforward protection protocol for 2-amino-2'-deoxyadenosine (1). Drastic hydrolysis conditions are necessary for the complete removal of benzoyl protecting groups [4] while the more labile phenoxyacetyl (pac) residues were removed without difficulty [26], [27]. Nevertheless, the formation of the N-bis-acylated derivatives is encountered with difficulties due to the monoprotection of the molecule [28] and an increased tendency of the acylated derivatives to be subjected to depurination [4]. According to the observation that the nucleosides 2a,b are significantly more stable than the purine nucleoside 1 (Table 1) the N-acyl derivatives of 2a,b should show similar properties. Therefore, the phenoxyacetyl derivative 5a as well as the benzoyl compound 5b were prepared employing the protocol of transient protection [29]. 2,4,5-Trichlorophenylphenoxy acetate [27] or benzoyl chloride [30] were used as acylation reagents. The bis-phenoxyacylated derivative 5a was formed as it was described for the purine compound 1 [19]. However, the yield of 5a was rather low (30%). The bis-

25

benzoylated nucleoside 5b was isolated in much better yield (63%) (Scheme 2; a) 6a: Pyridine; Me₃SiCl, 2,4,5-trichlorophenylphenoxyacetate, 40°, 12 h. 6b: Pyridine, Me₃SiCl, PhCOCl, r.t., 12 h. b) Pyridine/(MeO)₂Tr-Cl, r.t., 4 h. c) THF, 2-cyanoethyl diisopropylphosphoramidochloridite, r.t. 30 min). Both compounds (5a,b) were converted into the DMT-derivatives 6a,b using the standard reaction conditions [31]. Phosphitylation in the presence of 2-cyanoethyl diisopropylphosphoroamidochloridite and (i-Pr)₂EtN afforded the phosphoramidites 7a,b [32]. However, as the benzoylated derivative 6b is poorly soluble in THF-solution, a large volume had to be used. According to this the phosphitylation was less effective and the yield of the phosphoramidite which is normally in the range of 80% was decreased to 67% (Scheme 2).

Scheme 2




Deprotection of compound 5a (25% aq. NH₃, 40°, HPLC-monitoring) showed that the first pac-group is removed within 5 min, while the removal of the second pac-group afforded 20 min. At r.t. a complete deprotection takes place in less than one hour. The complete removal of the two benzoyl groups of compound 5b (25% aq. NH₃, 40°, HPLC monitoring) requires 8 – 9 h while the deprotection of the bis-benzoylated purine nucleoside 1 afforded several days [4] [7]. From this point of view the benzoyl-protected phosphoramidite 7b represents a useful building block for the incorporation of compound 2a in oligonucleotides. Nevertheless, the low solubility of the intermediate 6b represents a problem.

In spite of this the synthesis of the N,N'-dialkylmethyliden derivatives was 8a,b undertaken. Original attempts to introduce N,N'-dimethylaminomethylidene residues into the purine nucleoside 1 failed due to the instability of the protecting groups [28]. The N,N'-dibutylaminomethyliden group [33] for the protection of the exocyclic amino function of an 8-aza-7-deazapurine analogue of 2'-deoxyisoguanosine was used previously [34]. The

- 94 -

same group was now introduced into the nucleosides 2a,b. The bis-amidines 8a,b were obtained as the major products (50% yield), while the mono adducts (9a,b) were formed as minor components (17%). The formation of the mono adduct might be circumvented when the more vigorous conditions are used. For the protected nucleoside 8a,b the time of complete deprotection (conc. ammonia, 40°, HPLC-monitoring at 260 nm) was determined to be 440 min for 8a and 450 min for 8b. A half-live value cannot be given as a mono-protected intermediate is formed.

Scheme 3



10

Subsequently, the 4,4'-dimethoxytrityl residues were introduced. After the work-up of the reaction followed by silica gel flash-chromatography one N,N'-dibutylaminomethylidene residue was hydrolyzed to give a formyl group (Scheme 3; a) Pyridine/(MeO)₂Tr-Cl, r.t., 4 h. b) THF, 2-cyanoethyl diisopropyl-phosphoramidochloridite, r.t., 30 min). This group is now protecting the 2-amino function, while the 6-amino group is still carrying the dibutylaminomethylidene residue. The structure of compounds 10a,b was established on the basis of NMR spectra. The amino group (10.77 ppm) and the proton of the formyl residue (9.56 ppm) are split in doublets with J = 9.88 Hz. This characteristic coupling

15

pattern has already been observed for the formyl derivatives of 5-aza-7-deaza-2'-deoxyguanosine [35] and 2-amino-8-aza-7-deazapurine 2'-deoxynucleoside [36].

Phosphitylation of the DMT-derivatives 10a,b was performed in THF in the presence of chloro(2-cyanoethyl)-(diisopropylamino)phosphine furnishing the phosphoramidites 11a,b (Scheme 3). These phosphoramidites as well as the corresponding building block 7b carrying benzoyl protecting groups can be efficiently used in solid-phase oligonucleotide synthesis resulting in high coupling yields. All compounds were characterized by ¹H-, ¹³C-, and ³¹P-NMR spectra and by elemental analysis (Table 3 and experimental part).

Table 3. ¹³C-NMR Chemical Shifts of Pyrazolo[3,4-d]pyrimidine 2'-Deoxyribonucleosides^{a)}

	C(2) ^{b,d)}	C(4) ^{d)}	C(5)	C(6) ^{d)}	C(7)	C=O	C=O	C(1')	C(2')	C(3')	C(4')	C(5')
)	C(7a)	C(3a)	C(4)	C(3)	/	/)))))
	C(6) ^{c)}))))	=CH	=CH)))))
2a	156.9	158.3	95.5	162.7	133.3	---	---	83.3	38.0	71.3	87.4	62.7
2b	157.4	157.6	94.5	162.7	119.2	---	---	83.0	37.5	70.9	87.3	62.4
5a	152.2	154.8	100.5	156.1	136.1	168.5	169.1	83.6	37.7	71.1	87.7	62.5
5b	153.3	155.2	102.2	156.1	134.0	165.6	166.3	83.5	37.6	70.9	87.5	62.3
6a	152.2	154.8	100.5	155.9	135.9	168.5	169.2	83.7	38.1	70.8	85.6	64.3
6b	153.3	155.3	102.3	156.1	132.2	165.7	166.4	83.7	38.0	70.9	85.6	64.4
8a ^{f)}	157.1	157.6	106.0	158.9	134.9	164.1	166.2	86.1	41.2	73.7	89.1	64.3
8b	156.1	156.1	103.6	157.9	121.4	162.1	164.4	83.3	37.5	70.8	87.5	62.2
8b ^{f)}	157.1	157.7	104.7	159.2	122.9	163.	166.7	86.2	41.0	73.8	89.0	64.2
9 ^{o)}	156.6	157.3	102.4	162.3	133.7	162.9	---	83.1	37.8	71.2	87.3	62.6
10a ^{f)}	155.5	155.5	107.1	156.8	134.9	161.8	163.7	84.2	38.3	73.6	86.7	64.9
10b	156.0	156.1	103.8	157.8	121.3	162.1	164.4	83.5	38.3	70.5	85.3	64.1

^{a)} Measured in (D₆)DMSO at 298 K. ^{b)} Purine numbering. ^{c)} Systematic numbering.

^{d)} Tentative. ^{e)} Superimposed by (D₆)DMSO. ^{f)} Measured in CDCl₃.

1.2. Oligonucleotides

1.2.1 Synthesis and Characterization. – Automated solid-phase synthesis of the oligonucleotides 102 - 116 (Tables 4 – 6) was performed using the phosphoramidites 7b and 11b as well as standard phosphoramidites. The oligonucleotides were detritylated and purified on oligonucleotide purification cartridges or by reversed-phase HPLC (conditions for purification see experimental part). The homogeneity of the compounds was proven by ion-exchange chromatography on a 4 X 250 mm NucleoPac PA-100 column (DIONEX,


P/N 043018, USA). The composition of the oligonucleotides was determined by tandem hydrolysis with snake venom phosphodiesterase and alkaline phosphatase as described (see Figure 2a,b) [2].

- 5 The 8-aza-7-deazapurin-2,6-diamine nucleosides 2a migrates slightly slower than dA. The bromo compound 2b is much more hydrophobic as it can be seen from the Figure 2b. From the modified oligonucleotides MALDI-TOF mass spectra were taken (Table 6, Experimental part). The correct mass was found in all cases which underlines that all protecting groups were split off during a 10 h ammonia treatment at 60°.


10

1.2.2 Duplex Stability


- Because of the presence of three hydrogen bonds the n^2A_d -dT base pair (II) is expected to show the same or a similar stability as a dG-dC pair (III). Experimental data obtained from DNA-melting experiments show that the thermodynamic stability of duplexes containing an n^2A_d -dT base pair is somewhat higher than that with a dA-dT pairs but still remains far below that of a dG-dC pair. Various studies have been performed to explain this dilemma but a convincing answer has not been given. A rather detailed examination of this matter has been undertaken by Sagy et al [11]. This author compared the thermal stabilities of a bidentate base pair represented by the motifs dA-dU, dI-dC, dA-dT and dI- m^5C_d with those of tridentate pairs such as n^2A_d -dT, dG-dC, dG- m^5C_d and others. The experiments were performed with alternating polynucleotides synthesized enzymatically. As long as a 2-amino group was absent (bidentate bases pairs), the various duplexes show all a very similar stability independently from the particular base structure. By insertion of a 2-amino group, which leads to the tridentate base pair this similarity disappears. The dG-dC base pair is now much more stable than the n^2A_d -dT pair. The authors noted also a significant effect of the 5-substituents attached to the pyrimidine moiety; while the presence of a 5-methyl group results in a rather small stabilization; larger alkyl groups destabilize the DNA.
- 25 Scheme 4




Watson Crick Base Pair I



Watson Crick Base Pair II



Watson Crick Base Pair III

Watson Crick Base Pair IV
R = H, Br

- In order to evaluate the influence of the 8-aza-7-deazapurin-2,6-diamine nucleosides 2a,b on the base pair stability in comparison to the purine nucleoside 1 all three compounds were incorporated into the non-self-complementary duplex 5'-d(TAGGTCAATACT) (102) and 5'-d(AGTATTGACCTA) (103). This duplex is used as a standard in our laboratory to study the influence of modified nucleosides on the thermal stability and the structural behavior of the helical formation. The T_m -value of (102•103) is 46° in 0.1 M NaCl in the presence of 10 mM $MgCl_2$ and 10 mM Na-cacodylate buffer. The incorporation of six n^2A_d -residues instead of six dA residues increases the T_m -value by only 4° (see duplex 104•105, Table 4). This corresponds to a 0.7° T_m -increase per residue. Similar findings have been reported from experiments performed in other laboratories [5 - 12]. When four 8-aza-7-deazapurin-2,6-diamine nucleosides 2a were replacing dA an increase of the T_m -value already from 46° to 52° was measured, which corresponds to a 1.5° increase per modified residue (compare duplexes 102•103 vs. 108•109). As the stability increase caused by an n^2A_d -residue was only 0.7°, the 8-aza-7-deazapurin-2,6-diamine forms a more stable tridentate base pairs with dT than that of the purin-2,6-diamine.

Table 4. T_m Value and Thermodynamic Data of Antiparallel Stranded Oligonucleotides Containing Purin-2,6-diamine 2'-Deoxyribonucleoside and Related Pyrazolo[3,4-d]-pyrimidine Analogs. ^{a)}

	T_m [°C]	ΔH^0 [kcal/mol]	ΔS^0 [cal/mol • K]	ΔG^0_{310} [kcal/mol]
5'-d(TAG GTC AAT ACT) (102) • (103) 3'-d(ATC CAG TTA TGA)	46	-86	-230	-10.3
5'-d(T1G GTC 11T 1CT) (104) • (105) 3'-d(ATC C1G TT1 TGA)	50	-58	-155	-11.2
5'-d(TAG GCC GGC ACT) (106) • (107)	65	-92	-247	-16.0

- 98 -

3'-d(ATC CGG CCG TGA)				
5'-d(TAG GTC 2a2aT ACT) (108) • (109)	52	-105	-299	-12.7
3'-d(ATC C2aG TT2a TGA)				
5'-d(TAG GTC 2b2bT ACT) (110) • (111)	68	-110	-297	-17.9
3'-d(ATC C2bG TT2b TGA)				
5'-d(TAG GTC 2b2bT ACT) (110) • (103)	56	-85	-232	-12.9
3'-d(ATC CAG TTA TGA)				
5'-d(TAG GTC AAT ACT) (102) • (111)	58	-88	-239	-13.4
3'-d(ATC C2bG TT2b TGA)				

^a) Measured at 260 nm in 0.1 M NaCl, 0.1 M MgCl₂, and 10 mM Na-cacodylate buffer, pH 7.0 with 5 + 5 µM oligomer concentration.


It has been shown that 7-substituents of 7-deazapurines as well as of 8-aza-7-deazapurine are well accommodated in the major groove of DNA [1 -3]. In particular, halogen substituents show very favorable properties with regard to duplex stability. This prompted us to replace dA-residues now by the 7-bromo derivative 2b. According to Table 4 the duplex (20•21) shows a remarkable stability. The T_m -value was 68° compared to 46° of the duplex with dA-dT pairs. The duplex stability is strengthened by 5.5 ° per modified base. The duplex (106•107) containing four dG-residues at the same positions lead to a T_m -value of 65° which corresponds to a 5° T_m -increase. Thus, the stability of a base pair motif IV (Scheme 4) which follows a dA-dT recognition motif is now as stable as a dG-dC pair (base pair III).

An explanation for this unexpected observation can be given on the basis of the basicity differences of the amino groups of the 8-aza-7-deazapurin-2,6-diamine nucleosides 2a,b compared to that of the purin-2,6-diamine nucleoside 1. As discussed above the third hydrogen bond of compound 1 contributes very little to the n^2A_d -dT base pair stability (Scheme 4, base pair II) while the 2-amino group of dG makes a large contribution to the stability of a dG-dC pair. With regard to the chemical structure of the molecule, both, the amino group of dG and isoG_d are part of an acylated guanidine moiety which is expected to be considerably less basic than the 2-amino group of n^2A_d which is also a part of guanidine system being now aromatic but is not connected to an electron-attracting moiety. These differences cause the low acidity of the 2-amino group of 1 compared to dG or isoG_d. This property is already noticeable in the extraordinary resistance of the acylated 2-amino group of 1 against alkaline deprotection (see above). Next, the question has to be answered why compounds 1 and 2a,b are different. Contrary to compound 1, one can draw

mesomeric structures of the base of compounds 2a,b with a positive charge either at the 2- or the 6-amino group and a negative one at the pyrazole nitrogen-2.

These mesomeric formulas explain the acidity (reduced basicity) of the 2-amino group of 2a over that of 1. When a bromo or other electron-attracting substituents are introduced at the 7-position this effect is even strengthened. Thus, the 2-amino-group of 2a will become a better donor for hydrogen bonds with dT than the 2-amino group of 1. The additional bromo substituent can cause several favorable effects: (i) It acidifies the 2-amino group further; (ii) it represents a hydrophobic group in the major groove of the DNA-duplex; (iii) it can form a hydrogen bond with the 2-amino group. Apart from these properties of the monomeric units the environmental conditions of the base within the nucleic acid duplexes have to modulate the stability of the n^2A -dT base pairs. Otherwise, it cannot be explained why the stability of such a base pair is different in DNA-duplexes, DNA-RNA hybrids or PNA-duplexes.

Scheme 5



Yet, it has not been proven whether the T_m -value increases linearly with the number of incorporated 8-aza-7-deazapurine residues. In the case of compound 1 such a study has been undertaken, which showed that in poly(dA-dT) a linear relationship exists when the replacement of dA by the n^2A_d -residues did not exceed 50%. However, there was no further T_m -increase at higher n^2A_d -incorporation when the T_m -values were measured at low salt concentration. A similar observation as made on synthetic oligonucleotides was obtained for DNA of the cyano phage S-2L which contains n^2A_d instead of dA [38].

- 100 -

In order to identify the helical structure of the duplexes CD-spectra of the duplexes 102•103 to 110•111 were measured [39]. The CD-spectra of all duplexes indicate a B-like DNA structure, with a positive Cotton effect around 270 nm to 290 nm and a negative lobe around 250 nm (see Figure 3a).

5

It has been reported that the purine nucleoside 1 forms more stable duplexes with complementary RNA than with DNA [40]. Therefore, oligonucleotides containing 1 or 2a,b were hybridized with a complementary oligoribonucleotide 114 (Table 5). According to Table 5 the DNA-RNA hybrid 104•114 shows a T_m -increase over that of the parent

- 10 102•114. The 8-aza-7-deazapurin-2,6-diamine residue 2a shows a similar effect as that of 1. An additional stabilization is observed when the bromo compound 2b was incorporated. All CD-spectra of the DNA-RNA hybrids (see Figure 3b) adopt the form of an A-DNA.

Table 5. T_m Value and Thermodynamic Data of RNA-DNA and DNA - DNA hybrids.^{a)}

	T_m [°C]	ΔH^0 [kcal/mol]	ΔS^0 [cal/mol • K]	ΔG_{310}^0 [kcal/mol]
5'-d(TAG GTC AAT ACT) (102) • (114) 3'-(AUC CAG UUA UGA)	45	-92	-264	-10.2
5'-d(T1G GTC 11T 1CT) (104) • (114) 3'-(AUC CAG UUA UGA)	48	-60	-162	-11.0
5'-d(TAG GTC 2a2aT ACT) (108) • (114) 3'-(AUC CAG UUA UGA)	48	-89	-263	-11.0
5'-d(TAG GTC 2b2bT ACT) (110) • (114) 3'-(AUC CAG UUA UGA)	53	-93	-258	-12.4

^{a)} See Table 4.

- 15 Finally, the base pairing properties of the nucleosides 1 and 2a,b were studied in parallel-stranded DNA (ps-DNA) [37], [42]. For this purpose it was necessary to exchange the dC-dG pair by a m^5C_d -dG and/or dC-iG_d pair. As standard duplexes the hybrids (102•115) or (116•103) were chosen. In both series of experiments the purin-2,6-diamine (1) destabilized the ps-hybrids (104•115 and 116•105). The nucleoside 2a already led to
- 20 slightly more stable duplexes (see Table 6). As in the case of antiparallel DNA the bromo derivative 2b resulted in a significant increase of the T_m -value (110•115 and 116•111). The

two sets of CD-spectra (Figure4a,b) of the ps-duplexes are rather different to those of aps-hybrids (see Figure4a).

From these results it is apparent that the 2-amino group causes destabilization in the series of purine compounds and reduces the stabilizing effect observed in aps-DNA significantly. This behavior is not surprising as it was already observed that mono-substituted purin-2-amine or 8-aza-7-deazapurin-2-amine have a rather unfavorable influence on the stability of the ps-DNA [36] [42].

Table 6. T_m value, Thermodynamic Data of Parallel Stranded Oligonucleotides Formed by Oligonucleotides Containing iG_d and m⁵iC_d. ^{a, b)}

	T_m [°C]	ΔH^0 [kcal/mol]	ΔS^0 [cal/mol • K]	ΔG^0_{310} [kcal/mol]
5'-d(TAG GTC AAT ACT) (102)•(115) [37] 5'-d(ATiC iCAiG TTA TiGA)	39	-74	-211	-8.8
5'-d(T1G GTC 11T 1CT) (104) • (115) 5'-d(ATiC iCAiG TTA TiGA)	36	-76	-200	-7.9
5'-d(TAG GTC 2a2aT ACT) (108) • (115) 5'-d(ATiC iCAiG TTA TiGA)	41	-62	-171	-9.0
5'-d(TAG GTC 2b2bT ACT) (110) • (115) 5'-d(ATiC iCAiG TTA TiGA)	45	-67	-184	-10.2
5'-d(TiCA TAA iCTiG iGAT) (116) • (103)[37] 5'-d(AGT ATT GAC CTA)	44	-85	-242	-10.3
5'-d(TiCA TAA iCTiG iGAT) (116) • (105) 5'-d(AGT 1TT G1C CTA)	39	-61	-170	-8.4
5'-d(TiCA TAA iCTiG iGAT) (116) • (109) 5'-d(AGT 2aTT G2aC CTA)	45	-80	-230	-10.3
5'-d(TiCA TAA iCTiG iGAT) (116) • (111) 5'-d(AGT 2bTT G2bC CTA)	48	-68	-186	-10.9

^{a)} Measured at 260 nm in 1 M NaCl, 0.1 M l₂, 60 mM Na-cacodylate buffer, pH 7.0 with 5 + 5 μ M oligomer concentration. ^{b)} d(iC) = m⁵iC_d = 2'-deoxy-5-methylisocytidine.

- 102 -

The two amino groups of the 8-aza-7-deazapurine nucleosides 2a,b can be protected with either benzoyl residues or dibutylaminomethylene groups without causing depurination problems during the acidic detritylation or the deprotection under standard conditions in ammonia. The problems of depurination and an extraordinary high stability of the 2-amino protecting groups of the purin-2,6-diamine nucleoside 1 is not observed.

1.3 Experimental Part

Monomers: General. See [2]. Flash chromatography (FC): 0.4 bar on silica gel 60 H (Merck, Darmstadt, Germany). Thin-layer chromatography (TLC): Aluminium sheets silica gel 60 F₂₅₄ (0.2 mm, Merck, Germany). TLC Scanning: CS-930 TLC scanner (Shimadzu, Japan). Solvent systems for flash chromatography (FC) and TLC: CH₂Cl₂ / MeOH 98:2 (A) CH₂Cl₂ / MeOH 95:5 (B), CH₂Cl₂ / MeOH 9:1 (C), CH₂Cl₂ / acetone 95:5 (D), CH₂Cl₂ / acetone 9:1 (E), CH₂Cl₂ / EtOAc 85:15 (F). M.p.: Büchi-SMP-20 apparatus (Büchi, Switzerland); uncorrected. NMR Spectra: Avance -DPX-250 and AMX-500 spectrometers (Bruker, Germany); δ values are in ppm downfield from internal SiMe₄ (¹H, ¹³C). Microanalyses were performed by Mikroanalytisches Labor Beller (Göttingen, Germany).

Oligonucleotides:

Synthesis and purification of Oligonucleotides 102 – 116. The synthesis was carried out on an automated DNA synthesizer (Applied Biosystems, model ABI 392-08 for phosphoramidite chemistry) in a 1 μ mol scale with 3'-phosphoramidites of [(MeO)₂Tr]ib²G_d, [(MeO)₂Tr]bz⁶A_d, [(MeO)₂Tr]bz⁴C_d and [(MeO)₂Tr]T_d, together with the phosphoramidites of the derivatives 7b and 11b. The synthesis of the oligomers followed the regular protocol of the DNA synthesizer for phosphoramidites. After cleavage from the solid-support, the oligonucleotides were deprotected in 25% NH₃/H₂O (12 – 15 h at 60°). The purification of the 5'-(dimethoxytrityl)-oligomers were performed by OPC cartridges as well as by reversed-phase HPLC (RP-18). The following solvent gradient was used (A, MeCN; B, 0.1 M (Et₃NH)OAc (pH 7.0)/ MeCN 95:5): 3 min 20% A in B, 12 min 15 – 40% A in B with a flow rate of 1.0 ml/min. To remove the 4,4'-dimethoxytrityl residues they were treated with 2.5% CHCl₂COOH/CH₂Cl₂ for 5 min at r.t. The detritylated oligomers were purified by reversed-phase HPLC with the gradient: 20 min 0 – 20% A in B with a flow rate of 1 ml/min. The oligomers were desalted on a short column (RP-18, silica gel) using H₂O for the elution of the salt, while the oligomers were eluted with MeOH/H₂O 3:2. The oligomers were lyophilized on a Speed-Vac evaporator to yield colorless solids which were frozen at –24°.

Table 7. Molecular Masses M⁺ of Oligonucleotides 12,13 and 18 - 21 determined by

MALDI-TOF Mass Spectroscopy

Oligomer	M+ (calc.)	M+ (found)
5'-d(TAG GTC AAT ACT) (102)	3644.4	3645
5'-d(AGT ATT GAC CTA) (103)	3644.4	3645
5'-d(TAG GTC 2a2aT ACT) (108)	3674.4	3677
5'-d(AGT 2aTT G2aC CTA) (109)	3674.4	3675
5'-d(TAG GTC 2b2bT ACT) (110)	3832.5	3830
5'-d(AGT 2bTT G2bC CTA) (111)	3832.5	3832

The MALDI-TOF-spectra were measured on a Biflex III by Dr. T. Wenzel (Bruker Saxonia, Germany). Nucleoside-Composition Analysis. The analyses were performed as described in [37]. Extinction coefficients of the nucleoside constituents: ϵ_{260} : dA 15400, dT 8800, dG 11700, dC 7600, iG_d 7400, m⁵iC_d 6300, 2a 8800, 2b 8700. Snake-venom phosphodiesterase (EC 3.1.4.1., *Crotallus durissus*) and alkaline phosphatase (EC 3.1.3.1., *E. coli*) were obtained from Roche Diagnostics GmbH, Germany.

Determination of T_m Values and Thermodynamic Data. Absorbance vs. temperature profiles were measured on a Cary-1/1E UV/VIS spectrometer (Varian, Australia) equipped with a Cary thermoelectrical controller. The T_m values were determined in the reference cell with a Pt-100 resistor, and the thermodynamic data were calculated using the Meltwin 3.0 program [43].

Circular Dichroism Spectra. The CD-spectra were recorded with a Jasco-600 (Jasco, Japan) spectropolarimeter with thermostatically (Lauda-RCS-6 bath) controlled 1cm cuvettes. UV-Spectra: 150-20 spectrometer (Hitachi, Japan).

3-Bromo-1-[2-deoxy- β -D-erythro-pentofuranosyl]-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine (2b). A soln. of compd 4b [18] (1.0 g, 2.6 mmol) in aq. 25% NH₃ (80 ml) was heated at 70° for 4 d in a steel vessel. The solvent was evaporated, and the residue was dissolved in hot H₂O. Crystallization occurred upon cooling. Colorless needles (646 mg, 72%). M.p. 155°. TLC (C): R_f 0.2. UV (MeOH): 260 (8700), 278 (9000). ¹H-NMR ((D₆)DMSO): 2.16, 2.68 (2m, H₂-C(2')); 3.38, 3.48 (2m, H₂-C(5')); 3.77 (m, H-C(4')); 4.36 (m, H-C(3')); 4.72 (t, J = 4.9, HO-C(5')); 5.19 (d, J = 4.1, HO-C(3')); 6.32 ('t', J = 6.5, H-C(1')); 6.39 (s, NH₂); 6.77 (br, NH₂). Anal. calc. for C₁₀H₁₃BrN₆O₃ (345.2): C 34.80, H 3.80, N 24.35; found: C 34.97, H 3.97, N 24.21.

- 104 -

N^4,N^6 -Bis(phenoxyacetyl)-1-[2-deoxy- β -D-erythro-pentofuranosyl]-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine (5a). Compd. 2a [15] (1.5 g, 5.6 mmol) was dried by coevaporation with pyridine. The residue was dissolved in pyridine (25 ml) and trimethylsilyl chloride (3.3 ml, 26 mmol) was added at r.t while stirring. Stirring was continued for 15 min and a
5 soln. of 2,4,5-trichlorophenyl phenoxyacetate (5.4 g, 16.4 mmol) [27] in pyridine (15 ml) was added in one portion. The reaction mixture was stirred for 16 h at 40°. It was cooled (ice bath) and H₂O (4.2 ml) was added. After 5 min, aq. 25% NH₃ (6 ml) was introduced, and the mixture was concentrated to dryness. The residue was applied FC (eluant: A \rightarrow C) yielding a colorless foam (900 mg, 30%). TLC (C): R_f 0.4. UV (MeOH): 266 (9100). ¹H-NMR ((D₆)DMSO): 2.26, 2.82 (2m, H₂-C(2')); 3.34, 3.48 (2m, H₂-C(5')); 3.82 (m, H-C(4')); 4.44 (m, H-C(3')); 4.73 (t, J = 5.5, HO-C(5')); 5.14 (s, OCH₂); 5.17 (d, J = 4.3, HO-C(3')); 6.60 ('t', J = 6.3, H-C(1')); 6.92 - 7.32 (m, arom. H); 8.41 (s, H-C(3)); 10.77 (s, NH); 11.30 (s, NH). Anal. calc. for C₂₆H₂₆N₆O₇ (534.5): C 58.42, H 4.90, N 15.72; found: C 58.68, H 4.78, N 15.20.

15

N^4,N^6 -Bis(phenoxyacetyl)-1-[2-deoxy-5-O-(4,4'-dimethoxytrityl)- β -D-erythro-pentofuranosyl]-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine (6a). Compd. 5a (0.8 g, 1.5 mmol) was coevaporated twice with pyridine. The residue was dissolved in pyridine (2 ml) and 4,4'-dimethoxytrityl chloride (0.6 g, 1.77 mmol) was added. Stirring at r.t. was
20 continued for 4 h than the soln. was diluted with MeOH (5 ml) and washed with 5% aq. sodium bicarbonate (3 x 20 ml). The organic layer was dried (Na₂SO₄) and concentrated to dryness. The residue was purified by FC (eluant: D \rightarrow E) yielding a colorless foam (310 mg, 25%). TLC (E): R_f 0.3. UV (MeOH): 266 (8900). ¹H NMR: ((D₆)DMSO): 2.32, 2.81 (2m, H₂-C(2')); 2.99 (2m, H₂-C(5')); 3.67 (2s, OCH₃); 3.95 (m, H-C(4')); 4.53 (m, H-C(3'));
25 5.15 (s, OCH₂); 5.33 (d, J = 4.7, HO-C(3')); 6.62 ('t', J = 6.3, H-C(1')); 6.69 - 7.31 (m, arom. H); 8.36 (s, H-C(3)); 10.83 (s, NH); 11.56 (s, NH). Anal. calc. for C₄₇H₄₄N₆O₉ (836.9): C 67.45, H 5.30, N 10.04; found: C 66.95, H 5.41, N 10.15.

N^4,N^6 -Bis(phenoxyacetyl)-1-[2-deoxy-5-O-(4,4'-dimethoxytrityl)- β -D-erythro-pentofuranosyl]-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine 3'-[(2-Cyanoethyl) N,N-Diisopropyl-phosphoramidite] (7a). To a soln. of compd. 7a (0.15 g, 0.18 mmol) in THF (0.5 ml), (¹Pr)₂EtN (0.12 ml, 0.71 mmol) and 2-cyanoethyl diisopropylphosphoramidochloridite (64 mg, 0.27 mmol) were added. Stirring under argon atmosphere at r.t. was continued for 30 min. Then, the mixture was diluted with
35 CH₂Cl₂ (20 ml) and 5% aq. NaHCO₃. The mixture was extracted with CH₂Cl₂ (3 x 15 ml).

The combined organic layers were dried (Na_2SO_4) and evaporated to give an oil. The residue was submitted to FC (eluant: F) yielding (80 mg, 43%) of a colorless foam. TLC (F): R_f 0.8. ^{31}P -NMR (CDCl_3): 149.2, 149.4.

- 5 N^4, N^6 -Bis(benzoyl)-1-[2-deoxy- β -D-erythro-pentofuranosyl]-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine (5b). Compd. 2a [17] (1 g, 3.8 mmol) was coevaporated twice with toluene and dissolved in pyridine (40 ml) and TMS-Cl (4.0 g, 36.8 mmol). The reaction mixture was stirred under argon atmosphere, cooled to 0° and PhCOCl (6.6 ml, 57 mmol) were added drop-wise to the soln. over a period of 30 min. After stirring overnight at r.t.
- 10 the mixture was diluted in EtOAc (200 ml) and washed with a sat. sodium bicarbonate soln. (200 ml) and ice cold H_2O (200 ml). The aq. layer was extracted with EtOAc (2 x 400 ml). The combined org. layers were evaporated to dryness and dissolved in THF/ MeOH/ H_2O 5:4:1 (250 ml). The dark orange soln. was cooled to 0° , and 2 N NaOH (25 ml) was added while stirring was continued for 40 min. The residue was purified by FC (eluant: A
- 15 \rightarrow B) yielding 1.1 g (61%) of an amorphous solid. TLC (B): R_f 0.3. UV (MeOH): 245 (16400), 274 (15200). ^1H -NMR ((D_6) DMSO): 2.13, 2.67 (2m, $\text{H}_2\text{-C}(2')$); 3.38, 3.52 (2m, $\text{H}_2\text{-C}(5')$); 3.84 (m, $\text{H-C}(4')$); 4.46 (m, $\text{H-C}(3')$); 4.72 (t, $J = 5.7$, $\text{HO-C}(5')$); 5.29 (d, $J = 4.4$, $\text{HO-C}(3')$); 6.66 (t', $J = 6.5$, $\text{H-C}(1')$); 7.51 - 8.11 (m, arom. H); 8.40 (s, $\text{H-C}(3)$); 10.95 (s, NH); 11.49 (s, NH). Anal. calc. for $\text{C}_{24}\text{H}_{22}\text{N}_6\text{O}_5$ (474.5): C 60.75, H 4.67, N 17.71; found:
- 20 C 61.03, H 4.70, N 17.58.

- N^4, N^6 -Bis(benzoyl)-1-[2-deoxy-5-O-(4,4'-dimethoxytrityl)- β -D-erythro-pentofuranosyl]-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine (6a). As described for 6a with 5a (500 mg, 1.05 mmol) and DMT-Cl (460 mg, 1.35 mmol) in pyridine (3 ml). The residue was purified by
- 25 FC (eluant: D \rightarrow E) yielding 400 mg (49%) of a colorless foam. TLC (E): R_f 0.4. UV (MeOH): 244 (16400), 275 (15200). ^1H -NMR ((D_6) DMSO): 2.35, 2.67 (2m, $\text{H}_2\text{-C}(2')$); 3.07, 3.10 (2m, $\text{H}_2\text{-C}(5')$); 3.68, 3.69 (2s, OCH_3); 3.97 (m, $\text{H-C}(4')$); 4.56 (m, $\text{H-C}(3')$); 5.35 (d, $J = 4.8$, $\text{HO-C}(3')$); 6.72 - 8.11 (m, arom. H); 8.36 (s, $\text{H-C}(3)$); 11.01 (s, NH); 11.54 (s, NH).

30

N^4, N^6 -Bis(benzoyl)-1-[2-deoxy-5-O-(4,4'-dimethoxytrityl)- β -D-erythro-pentofuranosyl]-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine 3'-[(2-Cyanoethyl) N,N-Diisopropylphosphoramidite] (7b). As described for 7a with 6b (110 mg, 0.14 mmol), $(^i\text{Pr})_2\text{EtN}$ (75 μl , 0.43 mmol) and 2-cyanoethyl diisopropylphosphoramidochloridite (38 μl ,

- 106 -

0.17 mmol) in THF (3 ml) at 30°. The oily residue was submitted to FC (eluant: F) yielding 92 mg (67%) of a colorless foam. TLC (F): R_f 0.8. ^{31}P -NMR (CDCl_3): 149.82, 149.63.

N^4, N^6 -Bis((di-n-butylamino)methylene)-1-[2-deoxy- β -D-erythro-pentofuranosyl]-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine (8a). To a stirred suspension of 2a (300 mg, 1.13 mmol) in MeOH (5 ml) N,N-di-n-butylformamide dimethyl acetal (790 μl , 3.39 mmol) was added. Stirring was continued for 2 h at 40°. The reaction mixture was evaporated to dryness and the residue adsorbed on silica gel. FC (eluant: A \rightarrow B) afforded two main products. From the faster migrating zone a colorless foam of 8a (330 mg, 53%) was isolated. TLC (B): R_f 0.4. UV (MeOH): 235 (22800), 274 (13800), 322 (25 700). ^1H -NMR (CDCl_3): 0.87 - 0.95 (m, CH_2CH_3); 1.26 - 1.38 (m, CH_2CH_3); 1.57 - 1.60 (m, CH_2CH_2); 2.42 - 2.92 (m, $\text{H}_2\text{-C}(2')$); 3.26 - 3.72 (2m, $\text{H}_2\text{-C}(5')$, NCH_2); 3.88 (m, $\text{H-C}(4')$); 4.39 (m, $\text{H-C}(3')$); 4.78 (t, $J = 5.8$, $\text{HO-C}(5')$); 5.21 (d, $J = 4.8$, $\text{HO-C}(3')$); 6.55 ('t', $J = 6.9$, $\text{H-C}(1')$); 7.93 (s, $\text{H-C}(3)$); 8.66 (s, N=CH); 8.88 (s, N=CH). Anal. calc. for $\text{C}_{28}\text{H}_{49}\text{N}_8\text{O}_3$ (544.7): C 61.74, H 8.88, N 20.57; found: C 61.71, H 8.91, N 20.57. From the slower migrating zone 1-[2-deoxy- β -D-erythro-pentofuranosyl]- N^6 -((di-n-butylamino)-methylene)-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine (9a) (80 mg, 17%) was obtained. TLC (B): R_f 0.3. UV (MeOH): 235 (23000), 275 (14900), 320 (22100). ^1H -NMR ((D_6) DMSO): 0.88 - 0.95 (m, CH_2CH_3); 1.27 - 1.36 (m, CH_2CH_3); 1.53 - 1.62 (m, CH_2CH_2); 2.21 (m, $\text{H-C}(2')$); 2.91 (m, $\text{H-C}(2')$); 3.38 - 3.59 (2m, $\text{H}_2\text{-C}(5')$, NCH_2); 3.79 (m, $\text{H-C}(4')$); 4.40 (m, $\text{H-C}(3')$); 4.78 (t, $J = 5.8$, $\text{HO-C}(5')$); 5.19 (d, $J = 4.4$, $\text{HO-C}(3')$); 6.36 (s, NH_2); 6.40 ('t', $J = 6.5$, $\text{H-C}(1')$); 7.77 (s, $\text{H-C}(3)$); 8.69 (s, N=CH). Anal. calc. for $\text{C}_{19}\text{H}_{31}\text{N}_7\text{O}_3$ (405.5): C 56.28, H 7.71, N 24.18; found: C 55.98, H 7.52, N 24.05.

N^4, N^6 -Bis((di-n-butylamino)methylene)-3-bromo-1-[2-deoxy- β -D-erythro-pentofuranosyl]-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine (8b). As described for 8a, with 2b (350 mg, 1 mmol) and N,N-di-n-butylformamide dimethyl acetal (720 μl , 3.1 mmol) in MeOH (7 ml) for 2 h at 40°. After FC (eluant: A \rightarrow B) two main products were isolated. The faster migrating zone gave 8b as a foam (310 mg, 50%). TLC(B): R_f 0.3. UV (MeOH): 235 (22500), 275 (14500), 320 (22100). ^1H -NMR ((D_6) DMSO): 0.94 - 0.99 (m, CH_2CH_3); 1.32 - 1.44 (m, CH_2CH_3); 1.59 - 1.73 (m, CH_2CH_2); 2.21 (m, $\text{H-C}(2')$); 2.91 (m, $\text{H-C}(2')$); 3.32 - 3.78 (2m, $\text{H}_2\text{-C}(5')$, NCH_2); 3.89 (m, $\text{H-C}(4')$); 4.40 (m, $\text{H-C}(3')$); 4.78 (t, $J = 5.8$, $\text{HO-C}(5')$); 5.19 (d, $J = 4.4$, $\text{HO-C}(3')$); 6.81 ('t', $J = 6.5$, $\text{H-C}(1')$); 8.69 (s, N=CH); 8.94 (s, N=CH). Anal. calc. for $\text{C}_{28}\text{H}_{47}\text{BrN}_8\text{O}_3$ (623.6): C 53.93, H 7.60, N 17.97; found: C 54.01, H 7.52, N 18.05. The slower zone furnished 3-bromo-1-[2-deoxy- β -D-erythro-

pentofuranosyl]-N⁶-((di-n-butylamino)methylene)-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine (9b) (75 mg, 16%). TLC(B): R_f 0.3. UV (MeOH): 236 (21600), 276 (14000), 320 (21900). ¹H-NMR ((D₆)DMSO): 0.90 - 0.96 (m, CH₂CH₃); 1.30 - 1.39 (m, CH₂CH₃); 1.59 - 1.73 (m, CH₂CH₂); 2.23 (m, H-C(2'')); 2.89 (m, H-C(2'')); 3.35 - 3.78 (2m, H₂-C(5'), NCH₂); 3.84 (m, H-C(4'')); 4.43 (m, H-C(3'')); 4.80 (t, J = 5.7, HO-C(5'')); 5.23 (d, J = 4.6, HO-C(3'')); 6.40 (s, NH₂); 6.81 ('t', J = 6.4, H-C(1'')); 8.65 (s, N=CH). Anal. calc. for C₁₉H₃₀BrN₇O₃ (484.4): C 47.11, H 6.24, N 20.24; found: C 47.23, H 6.52, N 20.35.

1-[2-Deoxy-5-O-(4,4'-dimethoxytrityl)-β-D-erythro-pentofuranosyl]-N⁴-((di-n-butylamino)methylene)-N⁶-formyl-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine (10a). As described for 6a with 8a (80 mg, 0.15 mmol), DMT-Cl (60 mg, 0.18 mmol) in pyridine (0.5 ml). The residue was purified by FC (eluant: D → E) yielding 85 mg (77%) of a colorless foam. TLC (E): R_f 0.3. UV (MeOH): 235 (22800), 276 (20000), 320 (24800). ¹H-NMR (CDCl₃): 0.98 - 1.05 (m, CH₂CH₃); 1.14 - 1.48 (m, CH₂CH₃); 1.65 - 1.78 (m, CH₂CH₂); 2.42 - 2.92 (m, H₂-C(2'')); 3.35 - 3.70 (2m, H₂-C(5'), NCH₂); 3.71 (s, OCH₃); 4.06 (m, H-C(4'')); 4.39 (m, H-C(3'')); 5.21 (d, J = 4.8, HO-C(3'')); 6.55 ('t', J = 6.6, H-C(1'')); 6.76 - 7.96 (m, arom. H); 7.94 (d, J = 10.5, NH); 7.96 (s, H-C(3)); 8.81 (s, N=CH); 9.57 (d, J = 10.5, CHO). Anal. calc. for C₄₁H₄₉N₇O₆ (735.9): C 66.92, H 6.71, N 13.32; found: C 66.85, H 6.56, N 13.40.

20 3-Bromo-1-[2-deoxy-5-O-(4,4'-dimethoxytrityl)-β-D-erythro-pentofuranosyl]-N⁴-((di-n-butylamino)methylene)-N⁶-formyl-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine (10b). As described for 6a with 8b (310 mg, 0.5 mmol), DMT-Cl (202 mg, 0.6 mmol) in pyridine (2 ml). The residue was purified twice by FC (eluant: D → E) yielding 250 mg (61%) of a colorless foam. TLC (E): R_f 0.3. UV (MeOH): 235 (22900), 276 (21100), 320 (25300). ¹H-NMR (D₆)DMSO: 0.90 - 0.96 (m, CH₂CH₃); 1.28 - 1.37 (m, CH₂CH₃); 1.61 - 1.68 (m, CH₂CH₂); 2.21 (m, H-C(2'')); 2.91 (m, H-C(2'')); 3.03 (2m, H₂-C(5'')); 3.35 - 3.52 (m, NCH₂); 3.69 (s, OCH₃); 3.90 (m, H-C(4'')); 4.48 (m, H-C(3'')); 5.32 (d, J = 4.7, HO-C(3'')); 6.46 ('t', J = 6.2, H-C(1'')); 6.73 - 7.31 (m, arom. H); 8.99 (s, N=CH); 9.56 (d, J = 9.89, NH); 10.77 (d, J = 9.88, CHO). Anal. calc. for C₄₁H₄₈BrN₇O₆ (814.8): C 60.44, H 5.94, N 12.03; found: C 60.36, H 5.73, N 11.85.

1-[2-Deoxy-5-O-(4,4'-dimethoxytrityl)-β-D-erythro-pentofuranosyl]-N⁴-((di-n-butylamino)methylene)-N⁶-formyl-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine 3'-[(2-Cyanoethyl) N,N-Diisopropylphosphoramidite] (11a). As described for 7a with 10a (110

mg, 0.15 mmol), (ⁱPr)₂EtN (78 μl, 45 mmol) and 2-cyanoethyl diisopropylphosphoramidochloridite (47 μl, 0.2 mmol) in THF (2 ml). The oily residue was submitted to FC (eluant: F) yielding 75 mg (53%) of a colorless foam. TLC (F): R_f 0.8. ³¹P-NMR (CDCl₃): 149.58, 149.52.

5

3-Bromo-1-[2-deoxy-5-O-(4,4'-dimethoxytrityl)-β-D-erythro-pentofuranosyl]-N⁴-((di-n-butylamino)methylene)-N⁶-formyl-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine 3'-[(2-Cyanoethyl) N,N-Diisopropylphosphoramidite] (11b). As described for 7a with 10b (90 mg, 0.11 mmol), (ⁱPr)₂EtN (63 μl, 0.36 mmol) and 2-cyanoethyl diisopropylphosphoramido-chloridite (31 μl, 0.14 mmol) in THF (1 ml). The oily residue was submitted to FC (eluant: F) yielding 80 mg (72%) of a colorless foam. TLC (F): R_f 0.8. ³¹P-NMR (CDCl₃): 149.58, 149.53.

10

15

Example 2: Synthesis of the 7-iodo-derivatives

Oligonucleotides were synthesized containing halogenated "purine" and pyrimidine bases. Bromo and iodo substituents were introduced into the 7-position of 8-aza-7-deazapurin-2,6-diamine (2b, c) or into the 5-position of uracil residues (3b, c). Phosphoramidites were synthesized using the isobutryl-residue for the protection of 2b and the benzoyl group for 2c. Duplexes containing the residues 2b or 2c gave always higher T_m values than those of the non-modified counterparts containing 2'-deoxyadenosine, the purin-2,6-diamine 2'-deoxyribonucleoside (1) or 2a at the same positions. Six 2b-residues replacing dA in the duplex 5'-d(TAGGTCAATACT) (102) • 5'-d(AGTATTGACCTA) (103) raised the T_m-value from 48° to 75° (4.5° per modification). Contrary to this, incorporation of the 5-halogenated 2'-deoxyuridines 3b or 3c into oligonucleotide duplexes showed very little influence on the thermal stability, regardless which "purine" nucleoside is located opposite to them. The positive effects on the thermal stability of duplexes observed in DNA were also found in DNA-RNA hybrids or DNA with parallel chain orientation.

20


25

30


The following data compare the effects of halogen substituents such as bromine and iodine introduced into the 7-position of 8-aza-7-deazapurin-2,6-diamine 2'-deoxyribonucleosides 2b, c and the 2'-deoxyuridine derivatives 3b, c (purine numbering is used ; see Scheme 6). These

nucleosides will be incorporated into various positions of duplex DNA, and the number of incorporations will be increased, stepwise. As the halogen substituents are directed into the major groove, in both the series, the "purine" and pyrimidine nucleosides, it was of interest to quantify the effects. Furthermore, the effect of multiple incorporations will be investigated as it is known from other duplex-stabilizing nucleosides, that the stabilizing effects level off with an increasing number of modified nucleoside incorporations (C. Bailly, M. J. Waring, *Nucleic Acids Res.* 1998, 26, 4309 and references therein; J. Sági, E. Szakonyi, M. Vorlickova, J. Kypr, *J. Biomolec. Struct. Dyn.* 1996, 13, 1035.). Oligonucleotide duplexes incorporating 7-halogenated 7-deazapurines (= pyrrolo[2,3-d]pyrimidines) show differences in the series of 7-iodo and 7-bromo compounds (F. Seela, M. Zulauf, *Chem. Eur. J.* 1998, 4, 1781), which result from the spatial requirements of the halogens (van der Waals radii: Br = 1.85 Å; I = 1.98 Å (A. Bondi, *J. Phys. Chem.* 1964, 68, 441) and/or from the more hydrophobic character of the iodo substituents compared to the bromo residues. Thus, the 7-iodo derivative of the 8-aza-7-deazapurin-2,6-diamine nucleoside (2c) was synthesized, converted into a phosphoramidite, and a number of oligonucleotides were synthesized containing the 7-iodinated 8-aza-7-deazapurine-2,6-diamine nucleoside 2c. A comparison of the thermal stability of these duplexes will illustrate the striking differences in duplex stabilization among halogens introduced into pyrimidine or 8-aza-7-deazapurine residues located in a similar environment of the major groove of a B-DNA.


20 Scheme 6



purine numbering



systematic numbering



2.1. Synthesis and Properties of Monomers

The 7-iodinated 8-aza-7-deazapurin-2,6-diamine 2'-deoxyribonucleoside 2c was prepared from the 6-isopropoxy compound 4 (F. Seela, G. Becher, *Synthesis* 1998, 2, 207) upon treatment with 25% aq. NH₃ at 70°C for 4 days in an autoclave. The halogenated nucleosides 2a, b F. (Seela, G. Becher, *Synthesis* 1998, 2, 207) as well as the 2'-deoxyuridine derivatives 3a-c have been described elsewhere (J. Asakua, M. J. Robins, *J. Org. Chem.* 1990, 55, 4928). The halogen substituents change the mobility of the nucleosides on an RP-HPLC column with the iodinated 2c and 3c as the slowest migrating compounds (Figure 5a, b). The retention times refer to the hydrophobic character of the nucleosides; the data were used later for the composition analysis of the base-modified oligonucleotides.

Data of the Iodo compound - Table 8. *Half-life Values (τ) of 2c at 25°.*

Compound	τ [min] ^{a)}	λ (nm)
I ⁷ NH ₂ ² c ⁷ z ⁸ A _d (2c)	stable ^{b)}	242
I ⁷ NH ₂ ² c ⁷ z ⁸ A _d (2c)	61 ^{c)}	242

^{a)} Measured in 0.5 N HCl. ^{b)} Within 3h. ^{c)}

Measured in 2 N HCl.

The nucleobases of the modified nucleosides 2a-c and 3a-c influence the N ↔ S pseudorotational equilibrium of the sugar moiety. This affects the conformation of the sugar phosphate backbone of DNA. For this reason the ¹H NMR spectra of the pyrazolo[3,4-d]pyrimidine nucleoside 2c as well as of the pyrimidine nucleosides 3a-c were measured in D₂O and ³J [¹H, ¹H] NMR coupling constants were determined. The conformational analysis was performed on the basis of ³J [¹H, ¹H] couplings using the program PSEUROT (Van J. Wijk, C. Altona, 'PSEUROT 6.2 – A Program for the Conformational Analysis of the Five-Membered Rings', University of Leiden, July 1993) According to Table 9 the 8-aza-7-deazapurin-2,6-diamine nucleoside 2a shows a higher N-conformer population than the corresponding purine nucleoside 1. The conformation around the C(4')-C(5') bond indicates that the 8-aza-7-deazapurin-2,6-diamine nucleoside 2c as the 8-aza-7-deaza-2'-

deoxy-guanosines (F. Seela, G. Becher, H. Rosemeyer, H. Reuter, G. Kastner, I. A. Mikhailopulo, *Helv. Chim. Acta* 1999, 82, 105) prefers the γ^t -(-sc)-rotamer population, while for the regular purine nucleosides the $\gamma^{(+)}g$ -(+sc)- or the $\gamma^{(-)}g$ -(ap)-conformation is predominant (G. Blackburn, M. J. Gait, 'Nucleic Acids in Chemistry and Biology', IRL Press, Oxford University Press 1990, p. 28) The

5 conformation of the sugar moieties of the pyrimidine nucleosides 3a-c is also influenced by the substitution at the 5-position (Table 9). The main change is observed between 3a without a 5-substituent and the derivatives 3b, c with Me or halogen substituents.

10 Table 9. 3J (H, H) Coupling Constants of the Sugar Moieties and N/S-Conformer Populations of the 2'-Deoxyribonucleosides 1 – 3 at 303 K ^{a)}

	$^3J_{H,H}/\text{Hz}$								Conformation			
	1',2'	1',2''	2',3'	2'',3'	3',4'	4',5'	4',5''	%N	%S	$\gamma^{(+)}g$	γ^t	$\gamma^{(-)}g$
1	7.30	6.10	7.00	3.10	3.40	3.20	4.30	31	69	62	25	13
2a	6.60	6.80	6.90	3.70	3.60	4.00	5.80	37	63	36	42	22
2c	6.60	6.70	6.85	3.90	3.85	4.30	5.90	38	62	31	43	26
dT [16]	---	---	---	---	---	---	---	36	64	---	---	---
3a	6.70	6.60	6.50	4.30	4.10	3.60	5.20	30	70	47	35	18
3b	6.45	6.50	6.50	4.70	4.50	3.40	4.70	34	66	55	30	15
3c	6.50	6.50	6.50	4.50	4.40	3.40	4.80	33	67	54	31	15

^{a)} Solvent D₂O; r.m.s. < 0.4 Hz; $|\Delta J_{\text{max}}| < 0.4 \text{ Hz}$.

[16] C. Thibaudeau, J. Plavec, J. Chattopadhyaya, *J. Org. Chem.* 1996, 61, 266.

15 As the reactivity of the amino groups of the nucleoside 2b is rather different to 1, various residues were studied for the base protection. Earlier, the nucleoside 2b was protected with N,N-di-(n-butyl)formamidine (dnb) residue. Now, the isobutyryl group was introduced employing the protocol of transient protection (2b → 13a) (G. S. Ti, B. L. Gaffney, R. A. Jones, *J. Am. Chem. Soc.* 1982, 104, 1316) As a side product the mono-protected nucleoside 14 was isolated (22%). The

20 formation of mono-acylated compounds has been observed in the case of other 2,6-diaminopurine nucleosides (I. Luyten, A. V. Aerschot, J. Rozenski, R. Busson, and P. Herdewijn, *Nucleosides Nucleotides* 1997, 16, 1649) When performing the alkaline hydrolysis of compounds 13a or 14 in 25% aq. NH₃ at 40°, a fast deprotection was observed (See Scheme 7). The half-life for 13a and 14 were determined and found to be 4.5 min and 20.5 min, respectively. It indicates that this

25 protecting group is appropriate for the solid-phase oligonucleotide synthesis. For the protection of

- 112 -




the iodo nucleoside 2c the transient protection protocol was used as in 2b, but in this case a benzoyl group was chosen (2c → 13b). The time for complete deprotection of 13b (25% aq. NH₃, 40°C, HPLC-monitoring at 260 nm) was 450 min. A half-life was not determined because of the step-wise reaction.

5

The base-protected nucleosides 13a, b as well as the pyrimidine nucleosides 3a-c were converted into the DMT-derivatives 15a, b, and 17a-c using standard reaction conditions (Y. S. Sanghvi, G. D. Hoke, S. M. Freier, M.C. Zounes, C. Gonzalez, L. Cummins, H. Sasmor, P. D. Cook, *Nucleic Acids Res.* 1993, 21, 3197) Phosphitylation of the DMT-derivatives 15a, b was performed in THF in the
10 presence of 2-cyanoethyl diisopropylphosphoramidochloridite furnishing the phosphoramidites 16a, b (Scheme 7); the pyrimidine building blocks 18a-c were prepared from 17a-c in dichloromethane (Y. S. Sanghvi, G. D. Hoke, S. M. Freier, M.C. Zounes, C. Gonzalez, L. Cummins, H. Sasmor, P. D. Cook, *Nucleic Acids Res.* 1993, 21, 3197) These phosphoramidites (16a, b and 18a-c) were employed in solid-phase oligonucleotide synthesis. All compounds were characterized by
15 ¹H-, ¹³C-, and ³¹P-NMR spectra and by elemental analysis (Table 10). Table 11 summarizes the ¹³C NMR data of 8-aza-7-deazapurine as well as those of pyrimidine nucleosides. The assignment was made according to gated-decoupled spectra. The NMR data of 3a-c were included as a search of the literature gave little information on that matter.

20 Scheme 7

- 113 -



(i) **13a:** Me_3SiCl , pyridine, isobutyric anhydride, r.t., 4h; **13b:** Me_3SiCl , pyridine, benzoyl chloride, r.t., 24h. (ii) $(\text{MeO})_2\text{TrCl}$, pyridine, r.t., 4h; (iii) 2-cyanoethyl diisopropylphosphoramidochloridite, CH_2Cl_2 , r.t., 30 min.

Table 10. ¹³C-NMR Chemical Shifts of Pyrazolo[3,4-d]pyrimidine-4,6-diamine 2'-Deoxyribo-nucleosides^{a)}

	C(2) ^{c)}	C(4) ^{c)}	C(5) ^{c)}	C(6) ^{e)}	C(7) ^{d)}	C=O/ =CH	C=O/ =CH	C(1')	C(2')	C(3')	C(4')	C(5')
	C(6) ^{d)}	C(7) ^{d)}	C(3 ^o) ^{d)}	C(4) ^{d)}	C(3) ^{d)}							
2b	157.6	157.4	94.5	162.7	119.2			83.0	37.5	70.9	87.3	62.4
2c	157.0	157.6	91.2	162.2	98.3			83.1	37.6	70.9	87.3	62.4
3a	163.2	150.5	101.8	140.6				84.2	38.5	70.5	87.4	61.3
3b	159.1	149.7	95.6	140.6				84.7	38.4	69.9	87.5	60.7
3c	160.4	150.0	69.9	145.0				84.6	^{g)}	69.2	87.4	60.7
13a	155.8	153.8	104.7	156.2	121.4	175.2	176.5	83.6	37.5	70.7	87.7	62.2
13b	149.6	152.1	105.7	153.4	96.3	169.4	177.5	84.4	38.7	71.6	88.6	63.1
14	156.0	156.5	97.4	157.5	119.1	175.3		83.3	37.5	70.8	87.5	62.2
15a	155.6	154.9	98.5	157.5	117.2	174.7	175.2	83.5	37.6	72.2	85.4	63.6
15b	149.5	152.1	105.8	153.2	96.2	---	177.5	84.8	^{f)}	71.6	86.7	65.3
17a	163.0	158.0	101.5	144.7				84.1	^{f)}	69.9	85.3	63.4
17b	158.0	159.1	96.0	144.7				84.9	38.4	70.3	85.7	63.6
17c	158.0	160.5	69.8	144.2				84.8	38.4	70.4	85.8	63.7

^{a)} Measured in (D₆)DMSO at 303 K. ^{c)} Purine numbering. ^{d)} Systematic numbering. ^{e)} Tentative.

^{f)} Superimposed by (D₆)DMSO.

2.2. Oligonucleotides

5 2. 2.1 Synthesis and Characterization

Automated solid-phase synthesis of the oligonucleotides (Tables 11-17) was performed using the phosphoramidites 16a, b and 18a-c as well as the standard building blocks. The syntheses followed the standard protocol (S. L. Beauge, M. Caruthers, *Tetrahedron Lett.* 1981, 22, 1859) and the coupling yields were always higher than 97%. Oligonucleotides containing halogenated dU-residues
 10 require the use of the 4-tert-butylphenoxyacetyl (tac) groups for the protection of the canonical phosphoramidites (E. Ferrer, C. Fàbrega, R. G. Garcia, F. Azorín, R. Eritja, *Nucleosides Nucleotides*

1996, 15, 907; J. C. Schulhof, D. Molko, and R. Teoule, *Nucleic Acids Res.* 1987, 15, 397; R. D. Sheardy and N. C. Seeman, *J. Org. Chem.* 1986, 51, 4301). In these cases the deprotection was performed with conc. ammonia at room temperature, while in all other cases the deprotection was carried out at 60°C. The oligonucleotides were detritylated and purified on purification cartridges (Applied Biosystems, 'User's Manual for Oligonucleotide Purification Cartridges') or by reversed-phase HPLC (conditions for purification see Exper. Part). The homogeneity of the compounds was proven by ion-exchange chromatography (see Exper. Part). The composition of the oligonucleotides was determined by tandem hydrolysis with snake venom phosphodiesterase and alkaline phosphatase followed by RP-18 HPLC as described (F. Seela, C. Wei, *Helv. Chim. Acta* 1999, 82, 726). Typical examples are shown in Figure 5c,d. The newly incorporated idonucleosides 2c or 3c migrate slower than the canonical DNA constituents. The oligonucleotides were also characterized by MALDI-TOF mass spectra. The detected masses were in good agreement with the calculated values (Table 18).

15 Table 18. Molecular Masses (MH^+) of Oligonucleotides Measured by MALDI-TOF Mass Spectrometry.

Oligomer	MH^+ (calc.)	MH^+ (found)
5'-d(TAG GTC AAT ACT) (102)	3644.4	3645
5'-d(AGT ATT GAC CTA) (103)	3644.4	3645
5'-d(TAG GTC 2a2aT ACT) (108)	3674.4	3677
5'-d(AGT 2aTT G2aC CTA) (109)	3674.4	3675
5'-d(TAG GTC 2b2bT ACT) (110)	3832.5	3830
5'-d(AGT 2bTT G2bC CTA) (111)	3832.5	3832
5'-d(T2bG GTC 2b2bT 2bCT) (201)	4020	4021
5'-d(AGT ATT G2cC CTA) (202)	3786 ?	3787
5'-d(AGT 2cTT GAC CTA) (203)	3786?	3792
5'-d(AGT 2cTT G2cC CTA) (100)	3926.5	3927
5'-d(TAG GTC 2c2cT ACT) (204)	3927	3931
5'-d(T2cG GTC 2c2cT 2cCT) (205)	4206.3	4210

- 116 -

5'-d(TAG G3aC AA3a ACT) (300)	3613.1	3616
5'-d(AGT A3a3a GAC CTA) (301)	3613.1	3615
5'-d(TAG G3bC AA3b ACT) (302)	3774.1	3775
5'-d(AGT A3b3b GAC CTA) (303)	3774.1	3772
5'-d(TAG G3cC AA3c ACT) (304)	3868.1	3871
5'-d(AGT A3c3c GAC CTA) (305)	3868.1	3871

2.2.2 Base Pair Stabilities of the Oligonucleotides Duplexes


The 7-bromo nucleoside 2b was found to stabilize DNA duplexes strongly while the non-halogenated compound 2a contributes very little to the duplex stability (see also F. Seela, G. Becher, *Helv. Chim. Acta* 2000, 83, 928; F. Seela, G. Becher, M. Zulauf, *Nucleosides Nucleotides* 1999, 18, 1399). The contribution of the purin-2,6-diamine nucleoside 1 on the duplex stability is even lower (C. Cheong, I. J. Tinoco, A. Chollet, *Nucleic Acids Res.* 1988, 16, 5115; J. D. Hoheisel, H. Lehrach, *FEBS Lett.* 1990, 274, 103). Thus, DNA duplexes containing compound 1-thymine base pairs are only slightly more stable than those with dA-dT pairs. As it was not known whether the stabilizing effect of the bromo nucleoside 2b will increase continuously by an increasing number of incorporations or will level off by multiple incorporations – as it is reported for other modified nucleosides (C. Bailly, M. J. Waring, *Nucleic Acids Res.* 1998, 26, 4309 and references therein)- a series of oligonucleotides were synthesized containing the halogenated compound 2b in a consecutive manner or in distant position. The modified residues were incorporated in one or both strands of a double-stranded DNA. The total number of incorporations was increased in duplexes from 1 to 6. The non-halogenated duplexes containing compound 2a were prepared for comparison. Apart from the incorporation of the bromo nucleoside 2b, the iodo compound 2c was also studied. For all experiments the non-self-complementary duplex 5'-d(TAGGTCAATACT) (102) • 5'-d(AGTATTGACCTA) (103) was chosen.

The non-halogenated nucleoside 2a increases the T_m -value of the standard duplex 102•103 by only 1° per modified residue (Table 11 and F. Seela, G. Becher, 2001, submitted). Contrary, the bromo compound 2b contributes a 4-5° stabilization per modified residue which represents an outstanding high stabilization induced by a non-canonical base. The strength of this effect is sequence-

dependent but the stability of the duplexes increases with an increasing number of 2b -
 incorporations as shown in Figure 6. When the iodo nucleoside 2c was replacing the bromo
 compound 2b, a similar effect regarding duplex stabilization is observed (Table 11). The effects of
 halogen substituents introduced into the pyrazolo[3,4-d]pyrimidine derivatives of dG (F. Seela, G.
 5 Becher, *Helv. Chim. Acta* 1999, 82, 1640) or dA (F. Seela, M. Zulauf, *J. Chem. Soc. Perkin Trans. 1*
 1999, 479) amounts only to 2° per modified residue.

A similar set of experiments as described for the duplexes containing the halogenated 8-aza-7-
 deazapurine nucleosides 2b or 2c was performed with the halogenated 2'-deoxyuridine derivatives
 3b or 3c (Table 12). Neither the bromo nucleoside 3b nor the iodo nucleoside 3c increases the
 10 stability of the duplexes significantly compared to that of the non-halogenated 3a or dT. Thus, only
 the base pairs of type I (Scheme 8) with halogen substituents located at the 7-position of the 8-aza-
 7-deazapurine (pyrazolo[3,4-d]pyrimidine) moiety are stabilized by the halogen substituents while
 those of the type II (Scheme 8) with the halogens attached to the 5-positions of the pyrimidine base
 exert little influence compared to those with dA-dT pairs.

15 Scheme 8



20 Table 11. T_m Values and Thermodynamic Data of Duplex Formation of Oligonucleotides Containing the Pyrazolo[3,4-d]pyrimidine Nucleosides 2a-c Opposite to dT^{a)}

Compound	T_m [°C]	ΔH° [kcal/mol]	ΔS° [cal/mol K]	ΔG°_{310} [kcal/mol]
5'-d(TAG GTC AAT ACT) (102)	47	-83.8	-235.9	-10.6
3'-d(ATC CAG TTA TGA) (103)				
5'-d(TAG GTC 2a2aT ACT) (108)	50	-93.3	-263.09	-11.7

- 118 -

3'-d(ATC CAG TTA TGA) (103)				
5'-d(TAG GTC AAT ACT) (102)	50	-98.79	-279.63	-12.06
3'-d(ATC C2aG TT2a TGA) (109)				
5'-d(TAG GTC 2a2aT ACT) (108)	51	-98.66	-278.58	-12.26
3'-d(ATC C2aG TT2a TGA) (109)				
	54	-99.96	-280.74	-12.89
5'-d(TAG GTC 2bAT ACT) (200)				
3'-d(ATC CAG TTA TGA) (103)				
5'-d(TAG GTC 2b2bT ACT) (110)	56	-91.42	-251.66	-13.37
3'-d(ATC CAG TTA TGA) (103)				
5'-d(TAG GTC AT ACT) (102)	59	-91.85	-251.25	-13.92
3'-d(ATC C2bG TT2b TGA) (111)				
5'-d(TAG GTC 2bAT ACT) (200)	63	-100.70	-274.04	-15.70
3'-d(ATC C2bG TT2b TGA) (111)				
5'-d(TAG GTC 2b2bT ACT) (110)	67	-105.40	-285.02	-17.00
3'-d(ATC C2bG TT2b TGA) (111)				
5'-d(T2bG GTC 2b2bT 2bCT)(201)	64	-99.08	-268.47	-15.81
3'-d(ATC CAG TTA TGA) (103)				
5'-d(T2bG GTC 2b2bT 2bCT)(201)	75	-107.43	-283.41	-19.53
3'-d(ATC C2bG TT2b TGA) (111)				
5'-d(TAG GTC AAT ACT) (102)	51	-90.65	-254.58	-11.69
3'-d(ATC C2cG TTA TGA) (202)				
5'-d(TAG GTC AAT ACT) (102)	54	-93.22	-260.43	-12.45
3'-d(ATC CAG TT2c TGA) (203)				
5'-d(TAG GTC AAT ACT) (102)	57	-95.09	-263.29	-13.43
3'-d(ATC C2cG TT2c TGA) (100)				
5'-d(TAG GTC 2c2cT ACT)(204)	55	-96.46	-268.72	-13.12
3'-d(ATC CAG TTA TGA) (102)				
5'-d(TAG GTC 2c2cT ACT)(204)	59	-102.95	-284.63	-14.67
3'-d(ATC C2cGT T A TGA) (202)				
5'-d(TAG GTC 2c2cT ACT)(204)	66	-104.93	-284.73	-16.62
3'-d(ATC C2cGT T2c TGA) (100)				
5'-d(T2cG GTC2c2cT 2cCT) (205)	72	—	—	—
3'-d(ATC C2cG TT2c TGA) (100)				

5 a) Thermodynamic parameters are derived from the fitting of melting curves measured at 260 nm in 0.1 M NaCl, 10mM MgCl₂, and 10 mM Na-cacodylate buffer, pH 7.0 with 5 μ M + 5 μ M single strand concentration. The ΔG° are taken directly from the program Meltwin 3.0 referring to 310°. Earlier publications of our laboratory using the fitting program refer to the same temperature and not to 298° as indicated. The thermodynamic data determined from the van't Hoff plots using the concentration dependence of the T_m -values are consistent with those obtained from the curve fitting within 15%. The van't Hoff data of the formation of the duplex 102•103 are the following: $\Delta H^\circ = 86.8$ kcal/mol; $\Delta S^\circ = 243.7$ cal/K mol; $\Delta G^\circ_{310} = 11.3$ kcal/mol.


Table 12. T_m Values and Thermodynamic Data of Duplex Formulation of Oligonucleotides Containing the Pyrimidine Nucleosides 3a-c Opposite to dA ^{a)}

Compound	T_m [°C]	ΔH° [kcal/mol]	ΔS° [cal/mol K]	ΔG°_{310} [kcal/mol]
5'-d(TAG G3aC AA3a ACT) (300) 3'-d(ATC CAG TTA TGA) (103)	47	-86.35	-244.74	-10.45
5'-d(TAG GTC AAT ACT) (102) 3'-d(ATC CAG 3a3aA TGA) (301)	48	-92.11	-261.95	-10.86
5'-d(TAG G3aC AA3a ACT) (300) 3'-d(ATC CAG 3a3aA TGA) (301)	46	-86.37	-245.42	-10.25
5'-d(TAG G3bC AA3b ACT) (302) 3'-d(ATC CAG TTA TGA) (103)	48	-95.76	-271.88	-11.44
5'-d(TAG GTC A A T ACT) (102) 3'-d(ATC CAG 3b3bA TGA) (303)	49	-97.43	-276.95	-11.53
5'-d(TAG G3bC AA3b ACT) (302) 3'-d(ATC CAG 3b3bA TGA) (303)	49	-94.25	-267.57	-11.26
5'-d(TAG G3cC AA3c ACT) (304) 3'-d(ATC CAG TTA TGA) (103)	49	-96.38	-273.73	-11.48
5'-d(TAG GTC AAT ACT) (102) 3'-d(ATC CAG 3c3cA TGA) (305)	50	-94.99	-269.20	-11.50
5'-d(TAG G3cC AA3c ACT) (304) 3'-d(ATC CAG 3c3cA TGA) (305)	50	-99.14	-281.47	-11.85

^{a)} See Table 11.

- 5 As the stability of the base pairs of the halogenated pyrimidine nucleosides 3b or 3c with dA was low compared to the halogenated pyrazolo[3,4-d]pyrimidine compounds 2b or 2c incorporated opposite to dT, tridentate base pairs are formed in which the halogenated pyrimidine nucleosides 3a-c are located opposite to the purine-2,6-diamine nucleoside 1 instead of dA (Scheme 9, bp type III). In this case a strengthening of the base pair can be expected if the formation of a third
- 10 hydrogen bond is possible. However, the T_m -values of those duplexes were also not influenced significantly by the replacement of dA-residues by the nucleoside 1 (Table 13). This indicates that the 2-amino group does not participate in possible base pairs as shown in IIIa-c (Scheme 9), a finding which is similar to that observed for a base pair between 1 and dT (F. Seela, G. Becher, 2001, submitted; C. Cheong, I. J. Tinoco, A. Chollet, *Nucleic Acids Res.* 1988, 16, 5115; J. D. Hoheisel, H.
- 15 Lehrach, *FEBS Lett.* 1990, 274, 103). A similar behaviour was found in the case of duplexes containing the base pair IVa (Scheme 9) formed between nucleoside 2a and the pyrimidine nucleosides 3a-c (Table 13). Nevertheless, the duplexes incorporating the base pairs IVb,c are slightly more stable than those containing the base pairs IIIb,c.

Scheme 9



5 Table 13. T_m Values and Thermodynamic Data of Oligonucleotide Duplexes Containing the Halogenated Pyrimidine Nucleosides 3a-c Opposite to the Purin-2,6-diamine Nucleoside 1 or the Pyrazolo[3,4-d]pyrimidin-4,6-diamine Nucleoside 2a^o)

	T_m [°C]	ΔH° [kcal/mol]	ΔS° [cal/mol K]	ΔG°_{310} [kcal/mol]
5'-d(TAG GTC AAT ACT) (102) 3'-d(ATC C1G TT1 TGA) (105)	49	-83.15	-232.81	-10.94
5'-d(TAG G3aC AA3a ACT) (300) 3'-d(ATC C1G TT1 TGA) (105)	48	-73.03	-202.44	-10.25
5'-d(TAG G3bC AA3b ACT) (302) 3'-d(ATC C1G TT1 TGA) (105)	49	-73.84	-203.94	-10.59
5'-d(TAG G3cC AA3c ACT) (304) 3'-d(ATC C1G TT1 TGA) (105)	49	-77.60	-215.48	-10.76
5'-d(T1G GTC 11T 1CT) (104) 3'-d(ATC CAG TTA TGA) (103)	48	-54.74	-144.65	-9.88
5'-d(T1G GTC 11T 1CT) (104) 3'-d(ATC CAG 3a3aA TGA) (301)	48	-60.03	-161.49	-9.94
5'-d(T1G GTC 11T 1CT) (104) 3'-d(ATC CAG 3b3b TGA) (303)	48	-60.09	-160.37	-10.35
5'-d(T1G GTC 11T 1CT) (104) 3'-d(ATC CAG 3c3cA TGA) (305)	48	-62.35	-167.33	-10.45
5'-d(TAG GTC 2a2aT ACT) (108) 3'-d(ATC CAG 3a3aA TGA) (301)	49	-87.32	-245.54	-11.16
5'-d(T1G GTC 2a2aT ACT) (108) 3'-d(ATC CAG 3b3b TGA) (303)	50	-89.11	-250.34	-11.47
5'-d(TAG GTC 2a2aT ACT) (108) 3'-d(ATC CAG 3c3cA TGA) (305)	51	-81.32	-225.60	-11.35
5'-d(TAG G3aC AA3a ACT) (300) 3'-d(ATC C2aG TT2a TGA) (109)	49	-90.30	-253.96	-11.53
5'-d(TAG G3bC AA3b ACT) (302) 3'-d(ATC C2aG TT2a TGA) (109)	51	-93.05	-261.91	-11.82
5'-d(TAG G3cC AA3c ACT) (304) 3'-d(ATC C2aG TT2a TGA) (109)	50	-90.58	-254.11	-11.77

^{a)} See Table 11.

A significant duplex stabilization was observed when the halogenated 8-aza-7-deazapurine nucleosides 2b or 2c were incorporated opposite to the pyrimidine nucleosides 3a-c (Table 14 and Scheme 10, bp of type V and VI). The T_m -increase amounts to about 4-5° per incorporated residue of the halogenated nucleosides 2b or 2c. All duplexes containing the halogenated nucleosides 2b or 2c (Scheme 10) give very similar T_m -values, no matter which pyrimidine monomer is located opposite to it. Thus, only halogen substituents attached to the modified purine residues lead to a duplex stabilization while halogens linked to the 5-position of the 2'-deoxyuridine moiety contribute very little to the duplex stability (S. M. Freier and K.-H. Altman, *Nucleic Acids Res.* 1997, 25, 4429; F. Seela, Y. He, *Helv. Chim. Acta* 2000, 83, 2527; S. Wang, E. T. Kool, *Biochemistry* 1995, 34, 4125)

Scheme 10




Table 14. T_m Values and Thermodynamic Data of Oligonucleotides Containing the Pyrazolo[3,4-d]pyrimidin-4,6-diamine Nucleosides 2b-c Opposite to dT and 3a-c ^{a)}

Duplex	T_m [°C]	ΔH° [kcal/mol]	ΔS° [cal/mol K]	ΔG°_{310} [kcal/mol]
5'-d(TAG GTC 2bAT ACT) (200) 3'-d(ATC CAG 3a3aA TGA) (301)	52	-92.88	-260.97	-11.94
5'-d(TAG GTC 2b2bT ACT) (110) 3'-d(ATC CAG 3a3aA TGA) (301)	55	-96.78	-269.67	-13.14
5'-d(T2bG GTC 2b2bT 2bCT) (201) 3'-d(ATC CAG 3a3aA TGA) (301)	64	-101.57	-276.51	-15.82
5'-d(TAG GTC 2bAT ACT) (200) 3'-d(ATC CAG 3b3bA TGA) (303)	54	-98.06	-275.03	-12.76
5'-d(TAG GTC 2b2bT ACT) (110) 3'-d(ATC CAG 3b3b TGA) (303)	55	-95.42	-266.68	-12.71

- 122 -

5'-d(T2bG GTC 2b2bT 2bCT) (201)	65	-106.49	-290.34	-16.44
3'-d(ATC CAG 3b3bA TGA) (303)				
5'-d(TAG GTC 2bAT ACT) (200)	54	-87.97	-244.29	-12.20
3'-d(ATC CAG 3c3cA TGA) (305)				
5'-d(TAG GTC 2b2bT ACT) (110)	55	-89.17	-246.44	-12.74
3'-d(ATC CAG 3c3cTGA) (305)				
5'-d(T2bG GTC 2b2bT 2bCT) (201)	65	-101.57	-275.34	-16.18
3'-d(ATC CAG 3c3cA TGA) (305)				
5'-d(TAG G3aC AA3a ACT) (300)	56	-95.76	-265.45	-13.43
3'-d(ATC C2bG TT2b TGA) (111)				
5'-d(TAG G3bC AA3b ACT) (302)	56	-96.53	-268.70	-13.20
3'-d(ATC C2bG TT2b TGA) (111)				
5'-d(TAG G3cC AA3c ACT) (304)	57	-97.07	-269.07	-13.62
3'-d(ATC C2bG TT2b TGA) (111)				
5'-d(TAG G3aC AA3a ACT) (300)	55	-96.93	-270.69	-12.98
3'-d(ATC C2cG TT2c TGA) (100)				
5'-d(TAG G3bC AA3b ACT) (302)	55	-100.82	-282.19	-13.29
3'-d(ATC C2cG TT2c TGA) (100)				
5'-d(TAG G3cC AA3c ACT) (304)	55	-98.00	-273.26	-13.25
3'-d(ATC C2cG TT2c TGA) (100)				

^a) See Table 11.

2.2.3 Base Discrimination

- In order to investigate the discrimination of the iodo nucleoside 2c towards the four canonical DNA – constituents, hybridization experiments were performed according to Table 15. As expected, the base pair 2c-dT is the strongest (102 • 100, $T_m = 59^\circ$) while those of the duplexes forming mismatches melt at a significantly lower temperature (Table 15). The discrimination of the iodo nucleoside 2c is similar to that of the canonical nucleosides except that the duplex 401 • 100 ($T_m = 52^\circ$) shows a 7° lower T_m – value than the duplex 102 • 100 ($T_m = 59^\circ$), while duplex 401 • 103 ($T_m = 46^\circ$) has almost the same stability as the parent duplex 102 • 103 ($T_m = 48^\circ$). The high T_m - value of 401 • 103 is the result of the formation of the dG-dA Hoogsteen pair which is obviously not formed between dG and compound 2c.

Table 15. T_m - Values of Oligonucleotides Containing the Pyrazolo[3,4-d]pyrimidin-4,6-diamine Nucleoside 2c Opposite to the Four Canonical Nucleosides ^a)

15

and nucleosides 2c appears to be the four canonical nucleosides.)

Duplex	T_m [°C]	T_m [°C]
5'-d(TAG GTC AAT ACT) (102) 3'-d(ATC C2cG TT2c TGA) (100)	57	5'-d(TAG GTC AAT ACT) (102) 3'-d(ATC CAG TTA TGA) (103)
5'-d(TAG GAC AAT ACT) (400) 3'-d(ATC C2cG TT2c TGA) (100)	45	5'-d(TAG GAC AAT ACT) (400) 3'-d(ATC CAG TTA TGA) (103)

- 123 -

5'-d(TAG GGC AAT ACT) (401)	50	5'-d(TAG GGC AAT ACT) (401)	46
3'-d(ATC C2cG TT2c TGA) (100)		3'-d(ATC CAG TTA TGA) (103)	
5'-d(TAG GCC AATACT) (402)	45	5'-d(TAG GCC AATACT) (402)	36
3'-d(ATC C2cG TT2c TGA) (100)		3'-d(ATC CAG TTA TGA) (103)	

^a) see Table 11.

The CD-spectra of the duplexes containing the 2-amino-8-aza-7-deazaadenine derivatives 2a, b were measured next. A B-like DNA structure can be deduced from the curves displayed in Figure 7a. A positive Cotton effect around 270 to 290 nm and a negative lobe at 250 nm are observed for the standard duplex (102•103). The CD spectrum of the duplex 104•105 containing 1 shows significant differences similar to that of 110•111 while the duplex 108•109 shows a stronger negative Cotton effect at 260 nm and a positive one at 235 nm. Similar differences were reported for the oligonucleotides containing 7-substituted 8-aza-7-deaza-7-iodoguanine (N. Ramzaeva, F. Seela, *Helv. Chim. Acta* 1996, 79, 1549).

10

2.4 DNA-RNA Hybrids

In order to study the influence of the 2,6-diamino nucleosides 1 and 2a,b on the stability of DNA-RNA hybrids, the oligodeoxyribonucleotides 110, 300, 302, 304, 104 and 102 were hybridized with the oligoribonucleotide 403. Table 16 shows that the various modifications exert a significant influence on the duplex stability of the DNA-RNA hybrids. While the incorporation of the non-halogenated nucleosides 1 or 2a stabilizes the DNA-RNA structure very little, the hybridization of the oligonucleotide containing 2b with 403 (110 • 403) show a significant increase of the duplex stability as in DNA. Compounds 3b and 3c enhance the thermal stability of DNA-RNA to a small degree, which is also observed for other modifications of 2'-deoxyuridine (B. C Froehler, S. Wadwani, T. J. Terhorst, S. R. Gerrard, *Tetrahedron* 1992, 33, 5307; J. Sági, A. Szemző, K. Ébinger, A. Szabolcs, G. Sági, É. Ruff, L. Ötvös, *Tetrahedron Lett.* 1993, 34, 2191). The hypochromicities of the chimeric hybrids are slightly decreased over those of the DNA-DNA duplexes (data not shown). From the CD-spectra of Figure 7b it can be seen that the DNA-RNA hybrids adopt the A-form (N. Ramzaeva, C. Mittelbach, F. Seela, *Helv. Chim. Acta* 1997, 80, 1809).

25

Table 16. T_m Values and Thermodynamic Data of DNA-RNA Hybrids ^{a)}

Duplex	T_m [°C]	ΔH^0 [kcal/mol]	ΔS^0 [cal/mol • K]	ΔG_{310}^0 [kcal/mol]
5'-d(TAG GTC AAT ACT) (102) 3'-(AUC CAG UUA UGA) (403)	45	-92.1	-264.0	-10.2
5'-d(T1G GTC 11T 1CT) (104) 3'-(AUC CAG UUA UGA) (403)	48	-60.46	-162.12	-10.18
5'-d(TAG GTC 2a2aT ACT) (108) 3'-(AUC CAG UUA UGA) (403)	48	—	—	—
5'-d(TAG GTC 2b2bT ACT) (110) 3'-(AUC CAG UUA UGA) (403)	53	-98.64	-276.97	-12.74
5'-d(TAG G3aC AA3a ACT) (300) 3'-(AUC CAG UUA UGA) (403)	44	-74.43	-208.26	-9.84
5'-d(TAG G3bC AA3b ACT) (302) 3'-(AUC CAG UUA UGA) (403)	48	-77.19	-215.13	-10.47
5'-d(TAG G3cC AA3c ACT) (304) 3'-(AUC CAG UUA UGA) (403)	47	-78.77	-220.52	-10.37

^{a)} See Table 11.

2.5 Duplexes with Parallel Chain Orientation

- 5 The base pairing of the nucleosides 1, 2a-c as well as 3b, c was also investigated in parallel stranded DNA [29]. For this purpose it was necessary to replace the dC-dG pair by a m⁵iC_d-iG_d pair. The duplexes 102 • 115 and 116 • 103 served as standards (F. Seela, C. Wei, G. Becher, M. Zulauf, P. Leonard, *Bioorg. Med. Chem. Lett.* 2000, 10, 289). Substitution of dA-residues by the diamino nucleoside 1 resulted in a slight stability decrease of the ps-duplexes 104 • 115 and 116 • 105 (Table
- 10 17). The incorporation of 2a-c into the ps-DNA resulted in a significant increase of the T_m -values. The incorporation of the substituted pyrimidine nucleosides 3b, c in place of dT leads only to minor changes of the T_m -values. From this it can be concluded that similar to DNA with antiparallel chain orientation, the halogen substituents of the "purine" nucleosides 2b, c stabilize these duplexes while the halogen substituents of the pyrimidine nucleosides contribute very little to
- 15 the duplex stability.

Table 17. T_m -Values and Thermodynamic Data of Parallel-Stranded Oligonucleotide Duplexes Containing the Nucleosides 1, 2a-c and 3a-c ^{a, b)}

Duplex	T_m [°C]	ΔH^0 [kcal/mol]	ΔS^0 [cal/mol • K]	ΔG_{310}^0 [kcal/mol]
5'-d(TAG GTC AAT ACT) (102)	39	-74.4	-212.3	-8.5

- 125 -

5'-d(ATiC iCAiG TTA TiGA) (115)				
5'-d(T1G GTC 11T 1CT) (104)	36	-48.12	-129.68	-7.90
5'-d(ATiC iCAiG TTA TiGA) (115)				
5'-d(TAG GTC 2a2aT ACT) (108)	41	-61.77	-170.95	-8.75
5'-d(ATiC iCAiG TTA TiGA) (115)				
5'-d(TAG GTC 2b2bT ACT) (110)	45	-66.59	-183.94	-9.54
5'-d(ATiC iCAiG TTA TiGA) (115)				
5'-d(TAG G3bC AA3b ACT) (302)	37	-50.64	-137.87	-7.88
5'-d(ATiC iCAiG TTA TiGA) (115)				
5'-d(TAG G3cC AA3c ACT) (304)	36	-55.09	-152.14	-7.90
5'-d(ATiC iCAiG TTA TiGA) (115)				
5'-d(TiCA TAA iCTiG iGAT) (116)	44	-85.0	-242.0	-10.0
5'-d(AGT ATT GAC CTA) (103)				
5'-d(TiCA TAA iCTiG iGAT) (116)	39	-61.00	-169.68	-8.38
5'-d(AGT 1TT G1C CTA) (105)				
5'-d(TiCA TAA iCTiG iGAT) (116)	45	-80	-230	-10.3
5'-d(AGT 2aTT G2aC CTA) (109)				
5'-d(TiCA TAA iCTiG iGAT) (116)	48	-68.34	-186.29	-10.56
5'-d(AGT 2bTT G2bC CTA) (111)				
5'-d(TiCA TAA iCTiG iGAT) (116)	43	-76.03	-215.51	-9.19
5'-d(AGT A3b3b GAC CTA) (303)				
5'-d(TiCA TAA iCTiG iGAT) (116)	42	-67.15	-187.02	-9.15
5'-d(AGT A3c3c GAC CTA) (305)				

^a) Measured at 260 nm in 1 M NaCl, 100 mM MgCl₂, 60 mM Na-cacodylate buffer, pH 7.0 with 5 + 5 μM oligomer concentration. ^b) d(iC) = m⁵iC_d = 2'-deoxy-5-methylisocytidine.

2.3. Conclusion

The halogen substituents introduced into the 7-position of the 8-aza-7-deazapurine 2'-deoxynucleoside (2b, c) or the 5-position of 2'-deoxyuridine residues (3a-c) have very different influences on the stability of nucleic acid duplexes. In the case of the 7-halogenated 8-aza-7-deazapurine 2'-deoxynucleosides 2b, c, each monomer contributes about 4°-5° to the duplex stabilization, while the pyrimidine nucleosides 3a-c show very little influence or even no effect. This is surprising as the halogens in both series of nucleobases are directed into the major groove of B-DNA both being in a not identical but a very similar environment. When both, the halogenated pyrimidines and the halogenated 8-aza-7-deazapurine nucleosides are present in a DNA-duplex only the latter make a contribution to the duplex stabilization. Consequently, the increase of the hydrophobic character of the major groove induced by the lipophilic halogen substituents and the expelling of water molecules is not the major effect induced by the halogenation of the major

groove. Apparently, stacking interaction between the nearest neighbours are strengthened in the case of 2b, c but not with 3b, c.

2.4. Experimental Part

- 5 *Monomers. General.* See preceeding manuscripts (F. Seela, G. Becher, *Helv. Chim. Acta* 2000, 83, 928; F. Seela, G. Becher, M. Zulauf, *Nucleosides Nucleotides* 1999, 18, 1399; F. Seela, M. Zulauf, *J. Chem. Soc. Perkin Trans. 1* 1999, 479.) Flash chromatography (FC): 0.4 bar on silica gel 60 H (Merck, Darmstadt, Germany). Thin-layer chromatography (TLC): Aluminum sheets, silica gel 60 F₂₅₄ (0.2 mm, Merck, Germany). Solvent systems for FC and TLC: CH₂Cl₂ / MeOH 9:1 (A), CH₂Cl₂ /
- 10 MeOH 95:5 (B), CH₂Cl₂ / acetone 9:1 (C), CH₂Cl₂ / EtOAc 85:15 (D), CH₂Cl₂ / acetone 95:5 (E),. M.p.: Büchi-SMP-20 apparatus (Büchi, Switzerland); uncorrected. NMR Spectra: Avance -DPX-250 and AMX-500 spectrometers (Bruker, Germany); δ values are in ppm downfield from internal SiMe₄ (¹H, ¹³C). Microanalyses were performed by Mikroanalytisches Labor Beller (Göttingen, Germany).
- 15
- Oligonucleotides.* - Oligonucleotide synthesis was performed on a DNA synthesizer, model 392 (Applied Biosystems, Weiterstadt, Germany). Melting curves were measured with a Cary-1/3 UV/VIS spectrophotometer (Varian, Australia) equipped with a Cary thermoelectrical controller. The temperature was measured continuously in the reference cell with a Pt-100 resistor, and the
- 20 thermodynamic data of duplex formation were calculated using the Meltwin 3.0 program (J. A. McDowell, D. H. Turner, *Biochemistry* 1996, 35, 14077). The CD-spectra were recorded with a Jasco-600 (Jasco, Japan) spectropolarimeter with thermostatically (Lauda-RCS-6 bath) controlled 1 cm cuvettes. UV-Spectra: 150-20 spectrometer (Hitachi, Japan). The enzymatic hydrolysis of the oligomers was performed as described (F. Seela, G. Becher, *Helv. Chim. Acta* 2000, 83, 928; F.
- 25 Seela, G. Becher, M. Zulauf, *Nucleosides Nucleotides* 1999, 18, 1399) using the following extinction coefficients: ϵ_{260} : ⁷G_d 2700, dT 8800, dC 7300, dA 15400, dG 11700, 2a 8800, 2b 8700, 2c 8700, 3a 10000, 3b 4800, 3c 3700. Snake-venom phosphodiesterase (EC 3.1.15.1, *Crotallus durissus*) and alkaline phosphatase (EC 3.1.3.1, *E. coli*) were generous gifts from Roche Diagnostics GmbH,

Germany. The MALDI-TOF-spectra were measured on a *Biflex III* spectrometer (*Bruker Saxonia, Leipzig, Germany*).

Synthesis and Purification of Oligonucleotides. - The synthesis was carried out in a 1 - μ mol scale using 3'-phosphoramidites of [(MeO)₂Tr]ib²G_d, [(MeO)₂Tr]bz⁶A_d, [(MeO)₂Tr]bz⁴C_d and [(MeO)₂Tr]T_d for the synthesis of oligonucleotides containing 2a-c. and [(MeO)₂Tr]tac²G_d, [(MeO)₂Tr]tac⁶A_d, [(MeO)₂Tr]tac⁴C_d and [(MeO)₂Tr]T_d for the synthesis of oligonucleotides containing 3a-c. After cleavage of the oligonucleotides from the solid support, the first were deprotected in 25% aq. NH₃ for 12 - 15 h at 60°. The latter were incubated in 25% aq. NH₃ for 1.5 - 2 h at r. t. for deprotection. The purification of the 5'-(dimethoxytrityl)-oligomers was performed by reversed-phase HPLC (RP-18). The following solvent gradient was used (A, 0.1 M (Et₃NH)OAc (pH 7.0)/ MeCN 95:5; B, MeCN): 3 min 20% B in A, 12 min 20 - 40% B in A with a flow rate of 1.0 ml/min. The concentrated oligonucleotide solutions were treated with 2.5% CHCl₂COOH/CH₂Cl₂ for 5 min at r.t. to remove the 4,4'-dimethoxytrityl residues. The detritylated oligomers were purified by reversed-phase HPLC with the gradient: 20 min 0 - 20% B in A with a flow rate of 1 ml/min. The oligomers were desalted on a short column (RP-18, silica gel) and then lyophilized on a *Speed-Vac* evaporator to yield colorless solids which were frozen at -24°. The purified oligomers were dissolved in 100 μ l of double-distilled H₂O, and the purity was controlled by ion-exchange chromatography on a *Dionex-Nucleopac-PA-100* HPLC column (4 x 250 mm, P/N 043010; *Dionex, Idstein, Germany*).

Nucleoside-Composition Analysis. - The oligonucleotides were dissolved in 0.1M Tris-HCl buffer (pH 8.3, 200 μ l), and treated with snake-venom phosphodiesterase (3 μ l) at 37° for 45 min, and then alkaline phosphatase (3 μ l) at 37° for another 30 min. The reaction mixtures were analyzed on reversed-phase HPLC (RP-18, at 260 nm, gradient A, 0.7 ml/min). The retention time of 2a-c and 3a-c were used as standards (Figure 5). The extinction coefficients of the nucleosides and the peak areas were used for quantification of the composition of the oligonucleotides (Figure 5a-d).

1-(2-Deoxy- β -D-erythro-pentofuranosyl)-3-iodo-1H-pyrazolo[3,4-d]-pyrimidin-4,6-diamine (2c). A soln. of compd. 6-amino-1-[2-deoxy- β -D-erythro-pentofuranosyl]-3-iodo-4-isopropoxy-1H-pyrazolo[3,4-d]pyrimidine 12 (F. Seela, G. Becher, *Synthesis* 1998, 2, 207.) (1 g, 2.3 mmol) in a aq. 25% NH₃ soln. (80 ml) was heated at 70°C for 4 d. The solvent was evaporated to dryness, the residue dissolved in hot water and crystallized. Colorless needles (640 mg, 71%). M.p. 154°. TLC (A): R_f 0.2. UV (MeOH): 223 (31800), 260 (8700), 278 (9100). ¹H-NMR ((D₆)DMSO): 2.12 (m, H_α-C(2'')); 2.67 (m, H_β-C(2'')); 3.38, 3.44 (m, H₂-C(5'')); 3.73 (m, H-C(4'')); 4.33 (m, H-C(3'')); 4.73 (t, J = 5.7, OH-C(5'')); 5.17 (d, J = 4.3, OH-C(3'')); 6.27 (t, J = 6.5, H-C(1'')); 6.34 (br, NH₂); 6.62 (br, NH₂). Anal. calc. for C₁₀H₁₃N₆O₃ (392.2): C 30.63, H 3.34, N 21.43; found: C 30.91, H 3.61, N 21.27.

3-Bromo-1-(2-deoxy- β -D-erythro-pentofuranosyl)-4,6-[(2-methylpropanoyl)amino]-1H-pyrazolo[3,4-d]pyrimidine (12a). Compd. 2b (F. Seela, G. Becher, *Synthesis* 1998, 2, 207) (0.74 g, 2.14 mmol) was co-evaporated with anhydr. pyridine for three times and dissolved in anhydr. pyridine (5 ml) while stirring at r.t. Me₃SiCl (1.37 ml, 10.8 mmol), and after 15 min isobutyric anhydride (3.56 ml, 21.5 mmol) were added. Stirring was continued for 3 h. The reaction mixture was cooled in an ice-bath and diluted with H₂O (2.5 ml), 5 min later aq. 25% NH₃ (4.3 ml) was added. After stirring for 30 min the reaction mixture was evaporated to dryness and coevaporated with toluene (three times). The residue was purified by FC (CH₂Cl₂/MeOH 9:1) furnishing two zones. From the fast migrating zone compound 13a was obtained as a colorless amorphous solid (500 mg, 48%). R_f (A) 0.4. UV (MeOH): 284 (9500), 239 (33900). ¹H-NMR ((D₆)DMSO): 1.15 (m, 2 CH(CH₃)₂); 2.25 (m, H_α-C(2'')); 2.75 (m, H_β-C(2'')); 2.86 (m, 2 CH(CH₃)₂); 3.47 (m, H₂-C(5'')); 3.81 (m, H-C(4'')); 4.44 (m, H-C(3'')); 4.73 (t, J = 5.5, OH-C(5'')); 5.32 (d, J = 4.3, OH-C(3'')); 6.54 (t, J = 6.6, H-C(1'')); 10.59, 10.72 (2 s, 2 NH). Anal. calc. for C₁₈H₂₅BrN₆O₅ (485.3): C 44.55, H 5.19, N 17.32, found C 44.90, H 5.28, N 16.81.

4-Amino-3-bromo-1-(2-deoxy- β -D-erythro-pentofuranosyl)-6-[(2-methylpropanoyl)amino]-1H-pyrazolo[3,4-d]pyrimidine (14). The slower migrating zone from the above reaction afforded compound 14 as a colorless amorphous solid (0.2 g, 22%). R_f (A) 0.36. UV (MeOH): 282 (10900),

237 (49500). ¹H-NMR ((D₆)DMSO): 1.04, 1.07 (*m*, CH(CH₃)₂); 2.21 (*m*, H_α-C(2')); 2.71 (*m*, H_β-C(2')); 2.90 (*m*, CH(CH₃)₂); 3.46 (*m*, H₂-C(5')); 3.78 (*m*, H-C(4')); 4.38 (*m*, H-C(3')); 4.72 (*t*, *J* = 5.6, OH-C(5')); 5.28 (*d*, *J* = 4.3, OH-C(3')); 6.42 (*t'*, *J* = 6.4, H-C(1')); 6.96, 7.76 (*br*, NH₂); 10.08 (*br*, NH). Anal. calc. for C₁₄H₁₉BrN₆O₄ (415.2): C 40.49, H 4.61, N 20.24, found C 40.58, H 4.72, N 19.93.

1-[2-Deoxy-β-D-erythro-pentofuranosyl]-4,6-dibenzamido-3-iodo-1H-pyrazolo[3,4-d]-pyrimidine (13b). Compound 2c (1.0 g, 2.55 mmol) was co-evaporated twice with toluene. It was dissolved in anhydr. pyridine (40 ml), and TMS-Cl (3.25 ml, 25.5 mmol) was added while stirring. The reaction mixture was stirred under argon atmosphere, cooled to 0°, and PhCOCl (3.0 ml, 25.8 mmol) was added dropwise within 30 min. After stirring overnight at r.t., the mixture was diluted with EtOAc (200 ml), washed with a sat. aq. NaHCO₃ soln. (200 ml) and ice-cold H₂O (200 ml). The aq. phase was extracted with EtOAc (2 x 400 ml). The combined org. phases were evaporated to dryness and the residue dissolved in THF/MeOH/H₂O (250 ml, 5:4:1). The dark orange soln. was cooled to 0°C, then 2 N NaOH (25 ml) was added, and stirring was continued for another 40 min. The residue was purified by FC (CH₂Cl₂/MeOH, 98:2 → CH₂Cl₂/MeOH, 95:5) to yield an amorphous solid (1.15 g, 75%). TLC (B): R_f 0.4. UV (MeOH): 244 (17400), 276 (14200). ¹H-NMR ((D₆)DMSO): 2.13 (*m*, H_α-C(2')); 2.67 (*m*, H_β-C(2')); 3.38, 3.52 (*m*, H₂-C(5')); 3.84 (*m*, H-C(4')); 4.46 (*m*, H-C(3')); 4.72 (*t*, *J* = 5.7, OH-C(5')); 5.29 (*d*, *J* = 4.4, OH-C(3')); 6.66 (*t'*, *J* = 6.5, H-C(1')); 7.51 - 8.11 (*m*, arom. H); 10.54, 10.78 (*s*, 2 NH). Anal. calc. for C₂₄H₂₁IN₆O₅ (586.4): C 48.01, H 3.53, N 14.00; found: C 47.93, H 3.53, N 14.05.

3-Bromo-1-[2-deoxy-5-O-(4,4'-dimethoxytrityl)-β-D-erythro-pentofuranosyl]-4,6-diisobutylamino-1H-pyrazolo[3,4-d]pyrimidine (15a). Compound 13a (0.5 g, 1.03 mmol) was coevaporated with anhydr. pyridine for three times and dissolved in pyridine (1.5 ml). DMT-Cl (0.45 g, 1.33 mmol) was added, and the mixture was stirred at r.t. for 3 h. The reaction was quenched by addition of MeOH and the mixture evaporated to dryness and coevaporated with toluene for three times. FC (CH₂Cl₂: MeOH, 10:1) gave 15a as a colorless foam (0.57 g, 70%). R_f (B) 0.3. UV (MeOH): 237 (52000), 283 (10500). ¹H-NMR ((D₆)DMSO): 1.04 - 1.17 (*m*, 2 (CH₃)₂CH); 2.29 (*m*, H_α-C(2'));

- 2.85 (*m*, H_β-C(2')); 2.89 (*m*, 2 (CH₃)₂CH); 3.07 (*m*, H₂-C(5')); 3.71 (*s*, 2 MeO); 3.94 (*m*, H-C(4')); 4.46 (*m*, H-C(3')); 5.35 (*m*, OH-C(3')); 6.57 (*m*, H-C(1')); 10.57, 10.74 (*s*, 2 NH). Anal. calc. for C₃₉H₄₃BrN₆O₇ (787.2): C 59.47, H 5.46, N 10.67; found: C 59.08, H 5.37, N 10.39.
- 5 3-Bromo-1-[2-deoxy-5-O-(4,4'-dimethoxytriphenylmethyl)-β-D-erythro-pentofuranosyl]-4,6-diisobutyrylamino-1H-pyrazolo[3,4-d]pyrimidine 3'-[(2-Cyanoethyl)-N,N-diisopropylphosphoramidite (16a). To a soln. of compound 15a (0.24 g, 0.3 mmol) in anhydr. CH₂Cl₂ (3 ml) (Ar) (iPr)₂EtN (0.16 ml, 0.9 mmol) and 2-cyanoethyl diisopropylphosphoramidochloridite (91 □, 0.41 mmol) was added, and the mixture was stirred at
- 10 r. t. for 30 min. The reaction was monitored by TLC. Then, the reaction mixture was diluted with CH₂Cl₂, and the soln. washed with a 5% aq. NaHCO₃ soln. twice and with brine. The organic phase was dried (Na₂SO₄), concentrated, and the product was separated by FC to yield a colorless foam (0.25 g, 84%). R_f (E), 0.63, 0.69. UV (MeOH): 282 (10000), 237 (49500). ³¹P-NMR (CDCl₃), 149.61, 149.65. ¹H-NMR ((D₆)DMSO): 1.11-1.35 (*m*, 4 CH(CH₃)₂); 2.51 (*m*, H_α-C(2')); 2.66 (*m*, H_β-C(2')); 2.95-3.91 (*m*, H₂-C(5')); CH(CH₃)₂; CH₂CH₂; 3.80 (*s*, MeO), 4.26 (*m*, H-C(4')); 4.82 (*m*, H-C(3')); 6.71 (*m*, H-C(1')); 6.75-7.44 (*m*, arom. H); 8.39, 8.59 (*br*, 2 NH). Anal. calc. for C₄₈H₆₀BrN₈O₈P (921.2): C 58.36, H 6.08, N 11.35; found: C 58.86, H 6.18, N 11.55.
- 15 1-[2-Deoxy-5-O-(4,4'-dimethoxytrityl)-β-D-erythro-pentofuranosyl]-4,6-dibenzamido-3-iodo-1H-pyrazolo[3,4-d]-pyrimidine (15b). Compound 13b (450 mg, 0.75 mmol) was coevaporated twice with anhydr. pyridine. The residue was dissolved in pyridine (2 ml) and dimethoxytrityl chloride (305 mg, 0.9 mmol) was added. After 4 h stirring, the soln. was diluted with 5 ml of MeOH and washed with a 5% aq. NaHCO₃ soln. (3 x 20 ml). The organic phase was dried (Na₂SO₄) and concentrated to dryness. The residue was purified by FC (CH₂Cl₂/acetone, 95:5 → CH₂Cl₂/acetone,
- 20 9:1) yielding 375 mg (55%) of a colorless foam. TLC (CH₂Cl₂/acetone 9:1): R_f (B) 0.4. UV (MeOH): 244 (16900), 276 (16200). ¹H-NMR ((D₆)DMSO): 2.35 (*m*, H_α-C(2')); 2.67 (*m*, H_β-C(2')); 3.07, 3.09 (2 *m*, H₂-C(5')); 3.70 (*s*, 2 OCH₃); 3.96 (*m*, H-C(4')); 4.56 (*m*, H-C(3')); (*d*, *J* = 4.8, OH-C(3')); 6.72 - 8.11 (*m*, arom. H); 10.54, 10.78 (2 *s*, 2 NH). Anal. calc. for C₄₅H₃₉IN₆O₇ (902.73): C 59.87, H 4.35, N 9.31; found: C 59.93, H 4.33, N 9.39.
- 25











1-[2-Deoxy-5-O-(dimethoxytrityl)- β -D-erythro-pentofuranosyl]-4,6-dibenzamido-3-iodo-1H-pyrazolo[3,4-d]-pyrimidine 3'-[(2-Cyanoethyl-N,N-diisopropylphosphoramidite) (16b). Compound 15b (330 mg, 0.37 mmol) was dissolved in THF (5 ml). (iPr)₂EtN (186 μ l, 1.07 mmol) and 2-cyanoethyl diisopropylphosphoramidochloridite (108 μ l, 0.48 mmol) were added under argon atmosphere. After 30 min the mixture was diluted with CH₂Cl₂ (20 ml) and a 5% aq. NaHCO₃ soln (2 x 20 ml). The mixture was extracted with CH₂Cl₂ (3 x 15 ml). The combined organic layer was dried (Na₂SO₄) and evaporated to an oil. The residue was submitted to FC (CH₂Cl₂/EtOAc, 85:15) yielding 280 mg (69%) of a colorless foam. TLC (D): R_f 0.8. ³¹P-NMR (CDCl₃): 149.37, 149.38.

10

Example 3 –Investigation of array precursor compounds

According to standard methodology, the oligonucleotides shown in Table 19 were synthesized from the phosphoramidites according to this invention 19 (= 8a and 16a) to 23 in scheme 11 and standard A,G,C and T phosphoramidites as already described above. The phosphoramidites are also available from Glen Research and were used according to the manufacturer's instructions (Orderung numbers: 10-1906-02 (20), 10-1964-02 (21), 10-1056-02 (22), 10-1067-02 (23)). The hybridisation behaviour was investigated as described using UV/VIS spectroscopy measuring temperature dependent absorption at a wavelength of 260 nm.

20



- 5 Table 19. T_m -Values and Thermodynamic Data of Duplex Formation of Antiparallel or Parallel Oligonucleotide Duplexes Carrying Fluorescent Reporter Groups and/or Aminoalkyl Linkers for the Immobilization on Arrays^{a)}.

10

Oligonucleotide	T_m	ΔH°	ΔS°	ΔG°_{310}
-----------------	-------	------------------	------------------	------------------------

- 133 -

	[°C]	[kcal/mol]	[cal/K mol]	[kcal/mol]
5'-d(TAG GTC AAT ACT) (102) • (103) 3'-d(ATC CAG TTA TGA)	47	-86.8	-243.7	-11.3
5'-d(TAG GTC 2b2bT ACT) (110) • (103) 3'-d(ATC CAG TTA TGA)	56	-91.4	-251.7	-13.4
5'-d(24 TAG GTC AAT ACT) (500) • (103) 3'-d(ATC CAG TTA TGA)	49	-90.0	-254.4	-11.1
5'-d(24 TAG GTC 2b2bT ACT) (501) • (103) 3'-d(ATC CAG T T A TGA)	56	-87.2	-239.3	-13.0
5'-d(24 TAG GTC AAT ACT) (500) • (502) 3'-d(ATC CAG TTA TGA 25)	49	-95.2	-270.8	-11.2
5'-d(24 TAG GTC 2b2bT ACT) (501) • (502) 3'-d(ATC CAG T T A TGA 25)	56	-92.7	-256.6	-13.1
5'-d(24 TAG GTC AAT ACT) (500) • (503) 3'-d(ATC CAG TTA TGA 26)	50	-86.5	-242.7	-11.2
5'-d(24 TAG GTC 2b2bT ACT) (501) • (503) 3'-d(ATC CAG T T A TGA 26)	58	-95.1	-262.1	-13.8
5'-d(T 27 A TAA 27 T 27 27 TA) (504)•(505) 5'-d(A G T ATT G A G G AT)	42	-77.1	-220.6	-8.7
5'-d(24 T 27 A TAA 27 T 27 27 TA)(506)•(507) 5'-d(A G T ATT G A G G AT 26)	43	-71.9	-201.8	-9.3
5'-d(24 T 27 A T2b2b 27 T 27 27 TA) (508)•(507) 5'-d(A G T AT T G A G G AT 26)	48	-93.7	-265.4	-11.3

^{a)} Measured UV-spectrophotometrically at 260 nm in 10 mM Na-cacodylate, 10 mM MgCl₂, 100 mM NaCl (pH 7) at 5 μ M + 5 μ M of single strand concentration. The thermodynamic data were calculated using the program *MeltWin* (3.0).

5


It could be shown that the melting behaviour is very similar in the parallel and antiparallel mode even when labelled with reporter groups. The situation is similar to the situation when the nucleic acid binding compound according to the invention is bound to surface or builds up an array or when it binds to an array of natural or non-natural nucleic acid binding compounds exemplifying the usefulness of the present invention. This is exemplified by the oligonucleotide 501 containing a linker by which it can be attached to a solid phase and hybridizes to a labelled target nucleic acid like the oligonucleotide 503.

10


15 Example 4 – Visualization of antiparallel and parallel hybridization

The hydrogen bonding interaction pattern in parallel or antiparallel duplexes is visualized in scheme 12 and 13.


Scheme 12




Watson Crick AT-Base Pair I
antiparallel




Reverse Watson Crick AT-Base Pair II
parallel




Watson Crick n^2 AT-Base Pair III
antiparallel



Reverse Watson Crick n^2 AT-Base Pair IV
parallel



Watson Crick PT-Base Pair V
R = H, Br, I
parallel




Reverse Watson Crick PT-Base Pair VI
R = H, Br, I
parallel

5


Scheme 13

10


- 135 -




Watson Crick GC-Base Pair VII
antiparallel



Reverse Watson Crick iGC-Base Pair VIII
parallel



Reverse Watson Crick GiC-Base Pair IX
parallel



Reverse Watson Crick G^{me}iC-Base Pair X
parallel

References


- [1] N. Ramzaeva, F. Seela, *Helv. Chim. Acta* 1996, 79, 1549.
- [2] F. Seela, M. Zulauf, *J. Chem. Soc., Perkin Trans. 1* 1999, 479.
- [3] F. Seela, G. Becher, *Helv. Chim. Acta* 1999, 82, 94.
- 5 [4] B. L. Gaffney, L. A. Marky, R. A. Jones, *Tetrahedron* 1984, 40, 3.
- [5] C. Bailly, M. J. Waring, *Nucleic Acids Res.* 1998, 26, 4309.
- [6] A. Chollet, E. H. Kawashima, *Nucleic Acids Res.* 1988, 16, 305.
- [7] A. Chollet, A. Chollet - Damerius, E. H. Kawashima, *Chem. Scri.* 1986, 26, 37.
- [8] F. B. Howard, H. T. Miles, *Biochemistry* 1983, 22, 597.
- 10 [9] G. M. Lamm, B. J. Blencowe, B. S. Sproat, A. M. Iribarren, U. Ryder, A. I. Lamond, *Nucleic Acids Res.* 1991, 19, 3193.
- [10] J. D. Hoheisel, H. Lehrach, *FEBS Lett.* 1990, 274, 103.
- [11] J. Sági, E. Szakonyi, M. Vorlicková, J. Kypr, *J. Biomolec. Struct. Dyn.* 1996, 13, 1035.
- [12] S. Gryaznov, R. G. Schultz, *Tetrahedron Lett.* 1994, 35, 2489.
- 15 [13] G. Haaime, H. F. Hansen, L. Christensen, O. Dahl, P. E. Nielsen, *Nucleic Acids Res.* 1997, 25, 4639.
- [14] V. Boudou, L. Kerremans, B. De Bouvere, E. Lescrinier, G. Schepers, R. Busson, A. Van Aerschot, P. Herdewijn, *Nucleic Acids Res.* 1999, 27, 1450.
- [15] F. Seela, H. Driller, *Helv. Chim. Acta* 1988, 71, 757.
- 20 [16] G. Balow, V. Mohan, E. A. Lesnik, J. F. Johnston, B. P. Monia, O. L. Acevedo, *Nucleic Acids Res.* 1998, 26, 3350.
- [17] F. Seela, H. Steker, H. Driller, U. Bindig, *Liebigs Ann. Chem.* 1987, 15.
- [18] F. Seela, G. Becher, *Synthesis* 1998, 207.
- [19] L. D. Garaeva, I. V. Yartseva, S. Y. Melnik, *Nucleosides Nucleotides* 1991, 10, 1295.
- 25 [20] F. Oertel, H. Winter, Z. Kazimierczuk, J. A. Vilpo, P. Richter, F. Seela, *Liebigs Ann. Chem.* 1992, 1165.
- [21] J. Davoll, K. A. Kerridge, *J. Chem. Soc.* 1961, 2589.
- [22] L. D. Garaeva, I. A. Korbukh, Y. V. Dobrynin, T. G. Nikolaeva, M. N. Preobrazhenskaya, *Pharm. Chem. J. (Engl. Transl.)* 1988, 22, 523.
- 30 [23] J. van Wijk, C. Altona, 'PSEUROT 6.2 – A Program for the Conformational Analysis of the Five-Membered Rings', University of Leiden, July 1993; C. A. G. Haasnoot, F. A. A. M. de Leeuw, C. Altona, *Tetrahedron* 1980, 86, 2783.
- [24] F. Seela, G. Becher, H. Rosemeyer, H. Reuter, G. Kastner, I. A. Mikhailopulo, *Helv. Chim. Acta* 1999, 82, 105.

- [25] G. Blackburn, M. J. Gait, in 'Nucleic Acids in Chemistry and Biology', IRL Press, Oxford University Press 1990, p. 28.
- [26] B. S. Sproat, A. M. Iribarren, R. G. Garcia, B. Beijer, *Nucleic Acids Res.* 1991, 19, 733.
- [27] A. Cano, M. F. Goodman, R. Eritja, *Nucleosides Nucleotides* 1994, 13, 501.
- 5 [28] I. Luyten, A. Van Aerschot, J. Rozenski, R. Busson, P. Herdewijn, *Nucleosides Nucleotides* 1997, 16, 1649.
- [29] G. S. Ti, B. L. Gaffney, R. A. Jones, *J. Am. Chem. Soc.* 1982, 104, 1316.
- [30] K. Groebke, J. Hunzicker, W. Fraser, L. Peng, U. Diederichsen, K. Zimmerman, A. Holzner, Ch. Leumann, A. Eschenmoser, *Helv. Chim. Acta* 1998, 81, 175.
- 10 [31] B. C. Froehler, P. G. Ng, M. D. Matteucci, *Nucleic Acids Res.* 1986, 14, 5399.
- [32] N. D. Sinha, J. Biernat, J. McManus, H. Köster, *Nucleic Acids Res.* 1984, 12, 4539.
- [33] L. J. McBride, R. Kierzek, S. L. Beaucage, M. H. Caruthers, *J. Am. Chem. Soc.* 1986, 108, 2040.
- [34] F. Seela, C. Wei, A. Melenewski, Y. He, R. Kröschel, E. Feiling, *Nucleosides Nucleotides* 1999, 18, 1543.
- 15 [35] F. Seela, A. Melenewski, *Eur. J. Org. Chem* 1999, 485.
- [36] F. Seela, G. Becher, *Helv. Chim. Acta* 2000, 83, 928.
- [37] F. Seela, C. Wei, *Helv. Chim. Acta* 1999, 82, 726.
- [38] M. D. Kirnos, I. Y. Khudyakov, N. I. Alexandrushkina, B. F. Vanyushin, *Nature* 20 1977, 270, 369.
- [39] F. B. Howard, C. Chen, J. S. Cohen, H. T. Miles, *Biochem. Biophys. Res. Commun.* 1984, 118, 848.
- [40] E. A. Lesnik, S. M. Freier, *Biochemistry*, 1995, 34, 10807.
- [41] Ribozyme Pharmaceuticals Inc., 2950 Wilderness Place, Boulder, CO 80301, USA.
- 25 [42] F. Seela, C. Wei, G. Becher, M. Zulauf, P. Leonard, *Bio. Org. Med. Chem. Lett.* 2000, 10, 289.
- [43] J. A. McDowell, D. H. Turner, *Biochemistry*, 1996, 35, 14077.

Claims:

1. A nucleic acid binding compound comprising a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic group is a group of the general formula I

Formula I



wherein

R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of

- (1) -F, -Cl, -Br or -I,
 - (2) Nitro
 - (3) Cyano
 - (4) $-\text{COO}^-$
 - (5) $-(\text{C}_1\text{-C}_{10})$ -alkyl substituted according to (10)
 - (6) $-(\text{C}_2\text{-C}_{10})$ -alkenyl substituted according to (10)
 - (7) $-(\text{C}_2\text{-C}_{10})$ -alkynyl substituted according to (10)
 - (8) $-(\text{C}_6\text{-C}_{22})$ -aryl substituted according to (10)
 - (9) $-\text{W}-(\text{C}_1\text{-C}_{10})$ -alkyl, $-\text{W}-(\text{C}_2\text{-C}_{10})$ -alkenyl, $-\text{W}-(\text{C}_2\text{-C}_{10})$ -alkynyl, $-\text{W}-(\text{C}_6\text{-C}_{22})$ -aryl or W-H, wherein W = -S-, -O-, -NH-, -S-S-, -CO-, -COO-, -CO-NH-, -NH-CO-, -NH-CO-NH-, -NH-CS-NH-, $-(\text{CH}_2)_n\text{-[O-(CH}_2)_r\text{]}_s-$, where r and s are, independently of each other, an integer between 1 to 18 and n is 0 or 1 independently from r and s,
 - (10) substituents (5) to (9) wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -SH, $-\text{NO}_2$, -CN, -S- $(\text{C}_1\text{-C}_6)$ -alkyl, $-\text{NR}^5\text{R}^6$, $-\text{N}^+\text{R}^5\text{R}^6\text{R}^{12}$, $-\text{OR}^{12}$, $-\text{COR}^{11}$, $-\text{NH-CO-NR}^5\text{R}^6$, $-\text{NH-CS-NR}^5\text{R}^6$ and $-(\text{CH}_2)_n\text{-[O-(CH}_2)_r\text{]}_s-\text{NR}^5\text{R}^6$, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s, wherein R^{11} is selected from the group consisting of $-\text{NHR}^{12}$, OR^{12} , and $-\text{SR}^{12}$
- wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of -H, $-(\text{C}_1\text{-C}_{10})$ -alkyl, $-(\text{C}_2\text{-C}_{10})$ -alkenyl, $-(\text{C}_2\text{-C}_{10})$ -alkynyl, -

(C₆-C₂₂)-aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,

5 said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,

10 with the proviso that R⁵, R⁶ or R¹² is not a reporter group if the radicals (5) to (7) are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;

R², R³ is independent from X, R¹, R² and R³ and is selected from the group of,

- (1) -H
- 15 (2) (C₁-C₁₀)-alkyl,
- (3) (C₂-C₁₀)-alkenyl,
- (4) (C₂-C₁₀)-alkynyl,
- (5) (C₆-C₂₂)-aryl,
- (6) -Z-(C₁-C₁₀)-alkyl, -Z-(C₂-C₁₀)-alkenyl, -Z-(C₂-C₁₀)-alkynyl, -Z-(C₆-C₂₂)-aryl or Z-H, wherein Z = -CO-, -CO-NH-, -CS-NH-, -(CH₂)_n-[O-(CH₂)_r]_s-, where r and s are, independently of each other, an integer between 1 to 18 and n is 1 or 2 independently from r and s,
- 20 (7) substituents (2) to (6)

25 wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -NO₂, -OR¹², -CN, -SH, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,

30 wherein R¹¹ is selected from the group consisting of -NHR¹² and OR¹²,

wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group,

35 said alkyl, alkenyl, alkynyl or aryl in substituents (2) to (7) being unsubstituted or substituted by one or more moieties selected from the group consisting of -

- 140 -

halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s;

5 X is independent from R¹, R² or R³ and is selected from the group consisting of N and CH₃; and

D is the position of attachment of the group to the rest of the nucleic acid binding compound

or any salts thereof.

10 2. The nucleic acid binding compound of claim 1

wherein R¹ is alkynyl-amino -C≡C-E-NR⁵R⁶, alkenyl-amino -CH=CH-E-NR⁵R⁶ and -E-NR⁵R⁶ wherein E is -[(CH₂)_rF]_s-(CH₂)_r- wherein F = O or S and r and s are independently from one another an integer from 1 to 18,

15 wherein R⁵ and R⁶ are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group,


wherein R¹¹ is selected from the group consisting of -NHR¹² and OR¹²,

20 wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group,

said alkyl, alkenyl, alkynyl or aryl being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s.

25 3. The nucleic acid binding compound accordig to any of the claims 1 to 2, wherein said nucleic acid compound comprises one or more moieties of the general formula II

Formula II



- 141 -

wherein


- A is selected from the group consisting of O, S, CH₂, N-CO-(C₁-C₁₀)-alkyl,
 L is selected from the group consisting of oxy, sulfanediyl, -CH₂- and -NR²²-,
 T is selected from the group consisting of oxo, thioxo and selenoxo,
 5 U is selected from the group consisting of -OH, O⁻, -O-reporter group, -SH, -S
 reporter group -SeH, -(C₁-C₁₀)-alkoxy, (C₁-C₁₀)-alkyl, -(C₆-C₂₂)-aryl, -(C₆-
 C₁₄)-aryl-(C₁-C₁₀)-alkyl, -NR²³R²⁴, and -(O-(C₁-C₁₀)-alkyl)_n-R²⁵, wherein n
 can be any integer between 1 and 6, or wherein -NR²³R²⁴ can together with N
 be a 5-6-membered heterocyclic ring,
 10 V is selected from the group consisting of oxy, sulfanediyl, -CH₂-, or -NR²²-,
 R¹⁴ is selected from the group consisting of -H, -OH, -(C₁-C₆)-alkyl, -(C₁-C₁₀)-
 alkoxy, -(C₂-C₁₀)-alkenyloxy, -halogen, -azido, -O-allyl, -O-alkynyl, and -NH₂,
 R²² is independently selected from the group of -H and -(C₁-C₁₀)-alkyl,
 R²³ and R²⁴ are independently selected from the group consisting of -(C₁-C₁₀)-alkyl, -
 15 (C₁-C₂₀)-aryl, -(C₆-C₁₄)-aryl-(C₁-C₁₀)-alkyl, -(C₁-C₆)-alkyl-[NH(CH₂)_c]_d-
 NR²⁶R²⁷ and a reporter group,
 R²⁵ is selected from the group consisting of -H, -OH, -halogen, -amino,
 -(C₁-C₁₈)-alkylamino, -COOH, -CONH₂ and -COO(C₁-C₄)-alkyl and a
 reporter group,
 20 R²⁶ and R²⁷ are independently selected from the group consisting from -H,
 -(C₁-C₆)-alkyl, and -(C₁-C₄)-alkoxy-(C₁-C₆)-alkyl and a reporter group,
 c is an integer from 2 to 6,
 d is an integer from 0 to 6, and
 B is a moiety of formula I with R¹, R², R³ and X as specified in claim 1,
 25
 wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties
 selected from the group consisting of -halogen, -SH, -NO₂, -CN, -S-(C₁-C₆)-alkyl, -
 NR⁵R⁶, -N⁺R⁵R⁶R¹², -OR¹², -COR¹¹, -NH-CO-NR⁵R⁶, -NH-CS-NR⁵R⁶ and -(CH₂)_n-
 [O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to
 30 18 and n is 0 or 1 independently from r and s,
 wherein R¹¹ is selected from the group consisting of -NHR¹², OR¹², and -SR¹²
 wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of
 -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a
 reporter group, a group which facilitates intracellular uptake or a group that, when
 35 the nucleic acid binding compound hybridizes to its complementary nucleic acid,

attacks the latter while binding, cross-linking or cleaving,
or any salts thereof.

4. The nucleic acid binding compound according to any of the claims 1 to 3, wherein the backbone comprises a moiety of the general formula III, wherein t is 0 or 1,

5


Formula III



wherein in the case that t=1,

- A is selected from the group consisting of O, S, CH₂ and N-(C₁-C₆)-alkyl,
M is selected from the group consisting of oxy, sulfanediyl, -NR²²-, -(C₁-C₁₀)-alkyl-, or -O-(C₁-C₁₀)-alkyl-O-, and -S-(C₁-C₁₀)-alkyl-O- and -NR²²-(C₁-C₆)-alkyl-O-,
R²² is selected from the group of -H, -(C₁-C₁₀)-alkyl, a protecting group and a reporter group,
R¹⁴ is selected from the group consisting of -H, -OH, -(C₁-C₃)-alkyl, -(C₁-C₆)-allyl, -(C₁-C₁₀)-alkoxy, -(C₂-C₁₀)-alkenyloxy, -(C₂-C₁₀)-alkynyloxy, -(O-CH₂)_n- wherein n may be an integer of from 1 to 18, -halogen, -azido, SH, -(C₁-C₁₀)-alkylmercapto, O-reporter group, O-solid phase and -NH₂,
R¹⁵ is selected from the group consisting of -H, -(C₁-C₆)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₂-C₁₀)-alkyl-carbonyl, -(C₃-C₁₉)-alkenyl-carbonyl, -(C₃-C₁₉)-alkynyl-carbonyl, -(C₆-C₁₄)-aryl-(C₁-C₁₀)-alkyl, a solid phase and a group of formula IV

Formula IV



wherein


- 143 -

- T is selected from the group consisting of oxo, thioxo and selenoxo, and
- U is selected from the group consisting of -OH, O⁻, -O-reporter group, -SH, -SeH, -(C₁-C₁₀)-alkoxy, -(C₁-C₁₀)-alkyl, -(C₆-C₂₂)-aryl, -(C₆-C₁₄)-aryl-(C₁-C₁₀)-alkyl, -NR²³R²⁴, and -(O-(C₁-C₁₀)-alkyl)-R²⁵, wherein N can be any integer between 1 and 6, or wherein NR²³R²⁴ can together with N be a 5-6-membered heterocyclic ring,
- R²³ and R²⁴ are independently selected from the group consisting of -(C₁-C₁₀)-alkyl, -(C₁-C₂₀)-aryl, -(C₆-C₁₄)-aryl-(C₁-C₁₀)-alkyl, -(C₁-C₆)-alkyl-[NH(CH₂)_c]_d-NR²⁶R²⁷,
- R²⁵ is selected from the group consisting of -H, -OH, -halogen, -amino, -(C₁-C₁₈)-alkylamino, -COOH, -CONH₂ and -COO(C₁-C₄)-alkyl,
- R²⁶ and R²⁷ are independently selected from the group consisting from -H, -(C₁-C₆)-alkyl, and -(C₁-C₄)-alkoxy-(C₁-C₆)-alkyl
- R²⁹ is selected from the group consisting of -OR³⁰ and -SR³⁰,
- R³⁰ is selected from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₆-C₂₂)-aryl, a protecting group, a solid phase and a reporter group
- c is an integer from 2 to 6,
- d is an integer from 0 to 6, and
- B is a moiety of formula I with R¹, R², R³ and X as specified in claim 1, wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -SH, -NO₂, -CN, -S-(C₁-C₆)-alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -OR¹², -COR¹¹, -NH-CO-NR⁵R⁶, -NH-CS-NR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s, wherein R¹¹ is selected from the group consisting of -NHR¹², OR¹², and -SR¹² wherein R⁵, R⁶ and R¹² are selected independently from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₆-C₂₂)-aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving, and wherein in the case that t=0, R¹⁵ is -H, or any salts thereof.
5. The nucleic acid binding compound of claim 4, wherein t=0, R¹⁵ is -H and R¹⁴ is -H.

- 144 -

6. The nucleic acid binding compound according to any of the claims 1 to 5, wherein said backbone comprises a moiety of the formula V

Formula V




5 wherein

- A is selected from the group consisting of O, S, CH₂ and N-(C₁-C₆)-alkyl,
M' is selected from the group consisting of oxy, sulfanediyl, -NR²²-, -(C₁-C₁₀)-alkyl, or -O-(C₁-C₁₀)-alkyl-O-, and -S-(C₁-C₁₀)-alkyl-O- and -NR²²-(C₁-C₆)-alkyl-O-,
10 R²² is selected from the group of -H, a protecting group, a reporter group and -(C₁-C₁₀)-alkyl,
R¹⁴ is selected from the group consisting of -H, -OH, -(C₁-C₁₀)-alkoxy, -(C₂-C₁₀)-alkenyloxy, -(C₂-C₁₀)-alkynyloxy, -halogen, azido, -SH, -S-(C₁-C₆)-alkylmercapto, O-reporter group, O-solid phase and NH₂,
15 R¹⁶ is selected from the group consisting of -H, -(C₁-C₈)-alkyl, -(C₂-C₁₈)-alkenyl, -(C₂-C₁₈)-alkynyl, -(C₂-C₁₈)-alkyl-carbonyl, -(C₃-C₁₉)-alkenyl-carbonyl, -(C₃-C₁₉)-alkynyl-carbonyl, -(C₆-C₁₄)-aryl-(C₁-C₈)-alkyl, a protective group or a compound of formula IV

20

Formula IV



wherein

- T is selected from the group consisting of oxo, thioxo and selenoxo,
U is selected from the group consisting of -OH, -SH, -SeH, -(C₁-C₁₀)-alkoxy, -(C₁-C₁₀)-alkyl, -(C₆-C₂₂)-aryl, -(C₆-C₁₄)-aryl-(C₁-C₁₀)-alkyl, -NR²³R²⁴, and -O-(C₁-C₁₀)-alkyl-R²⁵, wherein n can be any integer between 1 and 6, wherein
25 NR²³R²⁴ can together with N be a 5-6-membered heterocyclic ring,

- 145 -

- R^{23} and R^{24} are independently selected from the group consisting of $-(C_1-C_{10})$ -alkyl, $-(C_1-C_{20})$ -aryl, $-(C_6-C_{14})$ -aryl- $-(C_1-C_{10})$ -alkyl, $-(C_1-C_6)$ -alkyl- $[\text{NH}(\text{CH}_2)_c]_d$ - $\text{NR}^{26}\text{R}^{27}$,
- R^{25} is selected from the group consisting of -H, -OH, -halogen, -amino,
 5 $-(C_1-C_{18})$ -alkylamino, -COOH, -CONH₂ and -COO $-(C_1-C_4)$ -alkyl,
- R^{26} and R^{27} are independently selected from the group consisting from -H,
 $-(C_1-C_6)$ -alkyl, and $-(C_1-C_4)$ -alkoxy- $-(C_1-C_6)$ -alkyl
- R^{29} is selected from the group consisting of -OR³⁰ and -SR³⁰,
- R^{30} is selected from the group consisting of -H, $-(C_1-C_{10})$ -alkyl, $-(C_2-C_{10})$ -alkenyl,
 10 $-(C_6-C_{22})$ -aryl, a protecting group, a solid phase and a reporter group, and
- c is an integer from 2 to 6,
- d is an integer from 0 to 6, and
- B is a moiety of formula I with R^1 , R^2 , R^3 and X as specified in claim 1,
- 15 wherein any alkyl, alkenyl, alkynyl or aryl can be substituted by one or more moieties selected from the group consisting of -halogen, -SH, -NO₂, -CN, -S $-(C_1-C_6)$ -alkyl, -NR⁵R⁶, -N⁺R⁵R⁶R¹², -OR¹², -COR¹¹, -NH-CO-NR⁵R⁶, -NH-CS-NR⁵R⁶ and $-(\text{CH}_2)_n$ -[O $-(\text{CH}_2)_r$]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s ,
- 20 wherein R^{11} is selected from the group consisting of -NHR¹², OR¹², and -SR¹²
- wherein R^5 , R^6 and R^{12} are selected independently from the group consisting of -H, $-(C_1-C_{10})$ -alkyl, $-(C_2-C_{10})$ -alkenyl, $-(C_2-C_{10})$ -alkynyl, $-(C_6-C_{22})$ -aryl and a reporter group, a group which facilitates intracellular uptake or a group that, when the nucleic acid binding compound hybridizes to its complementary nucleic acid, attacks the latter while binding, cross-linking or cleaving,
- 25 and any salts thereof.
7. The compound of claim 6, wherein M^1 is O, R^{16} is H and R^{14} is selected from the group consisting of -H and -OH.
8. The nucleic acid binding compound according to any of the claims 1 to 7, wherein
- 30 the backbone comprises sugar and phosphate moieties.
9. The nucleic acid binding compound according to claim 8, wherein the sugar is in a locked conformation.
10. The nucleic acid binding compound according to any of the claims 1 to 9 wherein the nucleic acid binding compound comprises a heterocyclic group which is a pyrrolo-[2,3-d]-pyrimidine, a pyrazolo[3,4-d]-pyrimidine or an analogue thereof.
- 35


- 146 -

11. The nucleic acid binding compound according to any of the claims 1 to 10 wherein a protecting group substitutes one or two hydrogen atoms of a -OH, -SH, -NH₂, -NH-alkyl, -NH-alkenylene, -NH-alkynylene, or a NH-aryl group.
12. The nucleic acid binding compound of claim 1 to 11 wherein said nucleic acid
5 binding compound contains a reporter group.
13. The nucleic acid binding compound according to claim 12 wherein the heterocyclic group of formula I as defined in claim 1 is 1 to 5 nucleotides separated from the nucleotide to which the reporter group is attached.
14. The nucleic acid binding compound of any of claims 1 to 11 wherein a heterocyclic
10 group is 7-bromo-7-deaza-8-aza-2,6-diamino-purine or 7-iodo-7-deaza-8-aza-2,6-diamino-purine.
15. A composition for analyzing interactions between nucleic acid binding compounds comprising an array of a plurality of nucleic acid binding compounds having different sequences, wherein said plurality of nucleic acid binding compounds are
15 coupled to a solid substrate at known locations and are selected to bind to complementary nucleic acid binding compounds characterized in that
only the nucleic acid binding compounds or the nucleic acid binding compounds and the complementary nucleic acid binding compounds together are nucleic acid
20 binding compounds comprising a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases, wherein a heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof.
16. A composition according to claim 15 wherein the substituted pyrazolo[3,4-
d]pyrimidine analogue is a pyrazolo[3,4-d]pyrimidine analogue of adenine or
25 guanine, a substituted pyrazolo[3,4-d]pyrimidine analogue of adenine or guanine or a 7-substituted pyrazolo[3,4-d]pyrimidine analogue of adenine or guanine.
17. A composition according to claim 15 wherein only the nucleic acid binding compounds or the nucleic acid binding compounds and the complementary nucleic acid binding compounds are nucleic acid binding compounds according to any of the
30 claims 1 to 14.
18. A composition according to any of the claims 15 to 17, wherein said solid substrate is selected from the group consisting of silica, polymeric materials, glass, porous glass, beads, chips, and slides.
19. A composition according to any of the claims 15 to 18 wherein said composition
35 comprises an array of nucleic acid binding compounds 5 to 20 nucleotides in length.

- 147 -

20. The binding product of a first nucleic acid binding compound according to any of the claims 1 to 15 or a composition according to any of the claims 16 to 19 with a second nucleic acid binding compound or a second a nucleic acid binding compound according to any of the claims 1 to 14, wherein the first nucleic acid binding compound or the composition and the second nucleic acid binding compound are bound to each other by base pairing in parallel or antiparallel orientation.
21. A compound of the general formula VIII

Formula VIII



10

wherein


- PPP is a triphosphate group, a thiotriphosphate group or analogues thereof,
 R^{14} is selected from the group consisting of -H, -OH, $-(C_1-C_{10})$ -alkoxy,
 15 $-(C_2-C_{10})$ -alkenyloxy, $-(C_2-C_{10})$ -alkynyloxy halogen, -azido and NH_2 ,
 R^{36} is selected from the group of -H and -OH, and
 B is a group of formula I as defined in claim 1

and any salts thereof

- 20 and wherein optionally at least one protecting group substitutes one or two hydrogen atoms of a -OH, -SH, NH_2 , NH-alkyl, -NH-alkenylene, -NH-alkynylene, or a -NH-aryl group.
22. A compound of claim 21, wherein R^{14} is H and B is 7-Bromo-7-deaza-8-aza-2,6-diaminopurine or 7-iodo-7-deaza-8-aza-2,6-diaminopurine.
23. A compound of formula VI

- 148 -

Formula VI



wherein


A is selected from the group consisting of O, S, CH₂ and N-(C₁-C₆)-alkyl,

5 R¹⁴ is selected from the group consisting of -H, -OH, -(C₁-C₁₀)-alkoxy, O-protecting group, S-protecting group, NH-protecting group, -(C₂-C₁₀)-alkenyloxy, -halogen, -azido, -SH, -(C₁-C₆)-alkylmercapto, O-reporter group and -NH₂,

10 R¹⁵ and R¹⁶ are independently selected from the group consisting of -H, -(C₁-C₈)-alkyl, -(C₂-C₁₈)-alkenyl, -(C₂-C₁₈)-alkynyl, -(C₂-C₁₈)-alkyl-carbonyl, -(C₃-C₁₉)-alkenyl-carbonyl, -(C₃-C₁₉)-alkynyl-carbonyl, -(C₆-C₁₄)-aryl-(C₁-C₈)-alkyl, a protecting group or a compound of formula IV,

with the proviso that R¹⁵ or R¹⁶ is a group of formula IV,

Formula IV



15

wherein

T is selected from the group consisting of oxo, thioxo and selenoxo,


20 U is selected from the group consisting of -OH, -SH, -SeH, -(C₁-C₁₀)-alkoxy, -(C₁-C₁₀)-alkyl, -(C₆-C₂₂)-aryl, -(C₆-C₁₄)-aryl-(C₁-C₁₀)-alkyl, -NR²³R²⁴, and -O-(C₁-C₁₀)-alkyl-O-(C₁-C₁₀)-alkyl-R²⁵, or wherein NR²³R²⁴ can together with N be a 5-6-membered heterocyclic ring,

R²³ and R²⁴ are independently selected from the group consisting of -(C₁-C₁₀)-alkyl, -(C₁-C₂₀)-aryl, -(C₆-C₁₄)-aryl-(C₁-C₁₀)-alkyl, -(C₁-C₆)-alkyl-[NH(CH₂)_c]-NR²⁶R²⁷,

- 149 -

- R^{25} is selected from the group consisting of -H, -OH, -halogen, amino,
 -(C₁-C₁₈)-alkylamino, -COOH, -CONH₂ and COO(C₁-C₄)-alkyl,
 R^{26} and R^{27} are independently selected from the group consisting from -H,
 -(C₁-C₆)-alkyl, and -(C₁-C₄)-alkoxy-(C₁-C₆)-alkyl,
 5 R^{29} is selected from the group consisting of -OR³⁰ and -SR³⁰,
 R^{30} is selected from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -
 (C₆-C₂₂)-aryl, a protecting group, a phosphate, a diphosphate and a reporter
 group, and
 M and M' are independently selected from the group consisting of oxy, sulfanediyl, -
 10 NR^{22} , -(C₁-C₁₀)-alkyl, or -O-(C₁-C₁₀)-alkyl-O-, and -S-
 (C₁-C₁₀)-alkyl-O- and -NR²²-(C₁-C₆)-alkyl-O-,
 R^{22} is selected from the group of -H and -(C₁-C₁₀)-alkyl, and
 c is an integer from 2 to 6,
 d is an integer from 0 to 6, and
 15 B is a moiety of formula I,


Formula I



wherein

- R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of
 20 substituents (1) to (10) for R^1 in formula I in claim 1 and
 (11)

- 150 -



wherein Nuc is the link to formula I and n is any integer from 1 to 18
said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being
unsubstituted or substituted by one or more moieties selected from the group
consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -
COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r
and s are independently of each other an integer of from 1 to 18 and n is 0 or 1
independently from r and s,

with the proviso that R⁵, R⁶ or R¹² is not a reporter group if the radicals (5)
to (7) are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;

R², R³ is independent from X, R¹, R² and R³ and is selected from the group of
substituents (1) to (7) for R² and R³ in formula I in claim 1,


X is independent from R¹, R² or R³ and is selected from the group consisting of N
and CH

D is the position of attachment of the group to the rest of the nucleic acid binding
compound.

whereby optionally at least one protecting group substitutes one or two hydrogen
atoms of a -OH, -SH, NH₂, NH-alkyl, -NH-alkenylene, -NH-alkynylene, or a -NH-
aryl group,
and any salts thereof.

24. A compound of formula VII

- 151 -




wherein

- 5 A is selected from the group consisting of O, S, CH₂ and N-(C₁-C₆)-alkyl,
M and M' are independently selected from the group consisting of oxy, sulfanediyl, -
NR²², -(C₁-C₁₀)-alkyl, or -O-(C₁-C₁₀)-alkyl-O-, and -S-
(C₁-C₁₀)-alkyl-O- and -NR²²-(C₁-C₆)-alkyl-O-,
R²² is selected from the group of -H and -(C₁-C₁₀)-alkyl,
10 R¹⁴ is selected from the group consisting of -H, -OR³¹, -(C₁-C₁₀)-alkoxy,
-(C₂-C₁₀)-alkenyloxy, -(C₂-C₁₀)-alkynyloxy, -halogen, -azido, NHR³¹, SR³¹,
R³¹ is a protecting group or a reporter group,
R³² and R¹⁷ are independently selected from the group consisting of -H, -(C₁-C₁₀)-
alkyl, -(C₂-C₁₀)-alkenyl, -(C₆-C₂₂)-aryl, or wherein NR³²R¹⁷ can form together
15 with N a 5-6-membered heterocyclic ring,
R¹⁸ is selected from the group consisting of -(C₂-C₆)-alkenyloxy, substituted or
unsubstituted -(C₁-C₆)-alkyl, unsubstituted -(C₁-C₆)-alkoxy or -(C₁-C₆)-alkoxy
substituted one or more times by a group selected from the group consisting of
-halogen, p-nitroaryloxy and -cyano, and
20 B is a group of formula I

- 152 -


Formula I



wherein

R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of substituents (1) to (10) for R^1 in formula I in claim 1 and

(11)



wherein Nuc is the link to formula I and n is any integer from 1 to 18

said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being

unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C_1 - C_6)-alkyl, -(C_1 - C_6)-alkoxy, -OH, - NR^5R^6 , -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,

with the proviso that R^5 , R^6 or R^{12} is not a reporter group if the radicals (5) to (7) are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;

R^2 , R^3 is independent from X, R^1 , R^2 and R^3 and is selected from the group of substituents (1) to (7) for R^2 and R^3 in formula I in claim 1


;

- X is independent from R^1 , R^2 or R^3 and is selected from the group consisting of N and CH
 D is the position of attachment of the group to the rest of the nucleic acid binding compound.

5

with the proviso that one or two hydrogen atoms of any -OH, -SH, -NH₂, -NH-alkyl, -NH-alkenylene, -NH-alkynylene, or -NH-aryl group are substituted by a protecting group,
 and any salts thereof.

- 10 25. A compound of general formula IX



wherein


- 15 A is selected from the group consisting of O, S, CH₂ and N-(C₁-C₆)-alkyl,
 M and M' are independently selected from the group consisting of oxy, sulfanediyl, -NR²², -(C₁-C₁₀)-alkyl, or -O-(C₁-C₁₀)-alkyl-O-, and -S-(C₁-C₁₀)-alkyl-O- and -NR²²-(C₁-C₆)-alkyl-O-,
 R²² is selected from the group of -H and -(C₁-C₁₀)-alkyl,
 20 R¹⁴ is selected from the group consisting of -H, -OR³¹, -(C₁-C₁₀)-alkoxy, -(C₂-C₁₀)-alkenyloxy, -(C₂-C₁₀)-alkynyloxy, -halogen, -azido, NHR³¹, SR³¹, or O-reporter group,
 R³¹ is a protecting group or a reporter group,
 R³² and R¹⁷ are independently selected from the group consisting of -H, -(C₁-C₁₀)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₆-C₂₂)-aryl, or wherein NR³²R¹⁷ can form together
 25 with N a 5-6-membered heterocyclic ring,
 R¹⁸ is selected from the group consisting of -(C₂-C₆)-alkenyloxy, substituted or unsubstituted -(C₁-C₆)-alkyl, unsubstituted -(C₁-C₆)-alkoxy or -(C₁-C₆)-alkoxy

- 154 -

substituted one or more times by a group selected from the group consisting of -halogen, p-nitroaryloxy and -cyano, and

B is a group of formula I


Formula I



5 wherein

R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of substituents (1) to (10) for R^1 in formula I in claim 1 and

10 (11)



wherein Nuc is the link to formula I and n is any integer from 1 to 18 said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,

with the proviso that R⁵, R⁶ or R¹² is not a reporter group if the radicals (5) to (7) are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;

20

- 155 -

R^2, R^3 is independent from X, R^1 , R^2 and R^3 and is selected from the group of substituents (1) to (7) for R^2 and R^3 in formula I in claim 1

X is independent from R^1 , R^2 or R^3 and is selected from the group consisting of N and CH


5 D is the position of attachment of the group to the rest of the nucleic acid binding compound.

with the proviso that one or two hydrogen atoms of any -OH, -SH, -NH₂, -NH-alkyl, -NH-alkenylene, -NH-alkynylene, or -NH-aryl group are substituted by a protecting group, and any salts thereof.

10

26. A compound of general formula X

Formula X



15

wherein

M and M' are independently selected from the group consisting of oxy, sulfanediyl, -NR²², -(C₁-C₁₀)-alkyl, or -O-(C₁-C₁₀)-alkyl-O-, and -S-(C₁-C₁₀)-alkyl-O- and -NR²²-(C₁-C₆)-alkyl-O-,

20

R²² is selected from the group of -H and -(C₁-C₁₀)-alkyl,

R¹⁴ is selected from the group consisting of -H, -OR³¹, -(C₁-C₁₀)-alkoxy, -(C₂-C₁₀)-alkenyloxy, -(C₂-C₁₀)-alkynyloxy, -halogen, -azido, NHR³¹, SR³¹, or O-reporter,


R³¹ is a protecting group or a reporter group,

25

B is a group of formula I

- 156 -

Formula I




wherein

R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of substituents (1) to (10) for R^1 in formula I in claim 1 and

5

(11)



wherein Nuc is the link to formula I and n is any integer from 1 to 18
 said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being
 unsubstituted or substituted by one or more moieties selected from the group
 consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -
 COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r
 and s are independently of each other an integer of from 1 to 18 and n is 0 or 1
 independently from r and s,

15

with the proviso that R^5 , R^6 or R^{12} is not a reporter group if the radicals (5)
 to (7) are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;

R^2 , R^3 is independent from X, R^1 , R^2 and R^3 and is selected from the group of
 substituents (1) to (7) for R^2 and R^3 in formula I in claim 1


- 157 -

- X is independent from R^1 , R^2 or R^3 and is selected from the group consisting of N and CH
- D is the position of attachment of the group to the rest of the nucleic acid binding compound.

5

with the proviso that one or two hydrogen atoms of any -OH, -SH, -NH₂, -NH-alkyl, -NH-alkenylene, -NH-alkynylene, or -NH-aryl group are substituted by a protecting group, and any salts thereof.

- 10 27. A precursor for the synthesis of a nucleic acid binding compound comprising a backbone, wherein the backbone comprises a moiety of the general formula VI




wherein

- 15 A is selected from the group consisting of O, S, CH₂ and N-(C₁-C₆)-alkyl, M and M' are independently selected from the group consisting of oxy, sulfanediyl, -NR²²-, -(C₁-C₁₀)-alkyl-, or -O-(C₁-C₁₀)-alkyl-O-, and -S-(C₁-C₁₀)-alkyl-O- and -NR²²-(C₁-C₆)-alkyl-O-,
- 20 R²² is selected from the group of -H, -(C₁-C₁₀)-alkyl, a protecting group and a reporter group,
- R¹⁴ is selected from the group consisting of -H, -OR³¹, -(C₁-C₁₀)-alkoxy, O-protecting group, S-protecting group, NH₂-protecting group, -(C₂-C₁₀)-alkenyloxy, -(C₂-C₁₀)-alkynyloxy, -halogen, -azido, SH, -(C₁-C₁₀)-alkylmercapto, and -O-solid phase,
- 25 R¹⁵ and R¹⁶ are independently selected from the group consisting of -H, -(C₁-C₆)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₂-C₁₀)-alkyl-carbonyl, -(C₃-C₁₉)-alkenyl-carbonyl, -(C₃-C₁₉)-alkynyl-carbonyl, -(C₆-C₁₄)-aryl-(C₁-C₁₀)-alkyl, protecting group and a solid phase

- 158 -

B is the link to a moiety of formula I,


Formula I



wherein

5 R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of substituents (1) to (10) for R^1 in formula I in claim 1 and

(11)



10

15

wherein Nuc is the link to formula I and n is any integer from 1 to 18
 said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being
 unsubstituted or substituted by one or more moieties selected from the group
 consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -
 COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r
 and s are independently of each other an integer of from 1 to 18 and n is 0 or 1
 independently from r and s,
 with the proviso that R⁵, R⁶ or R¹² is not a reporter group if the radicals (5)
 to (7) are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;

- 159 -

R^2, R^3 is independent from X, R^1, R^2 and R^3 and is selected from the group of substituents (1) to (7) for R^2 and R^3 in formula I in claim 1


X is independent from R^1, R^2 or R^3 and is selected from the group consisting of N and CH

5 D is the position of attachment of the group to the rest of the nucleic acid binding compound.

with the proviso that one or two hydrogen atoms of any -OH, -SH, -NH₂, -NH-alkyl, -NH-alkenylene, -NH-alkynylene, or -NH-aryl group are substituted by a protecting group,

10 and any salts thereof.

28. A precursor or intermediate of a nucleic acid binding compound, wherein the backbone comprises a moiety of the general formula III



15

wherein

A is selected from the group consisting of O, S, CH₂ and N-(C₁-C₆)-alkyl,

20 M is selected from the group consisting of oxy, sulfanediyl, -NR²²-, -(C₁-C₁₀)-alkyl-, or -O-(C₁-C₁₀)-alkyl-O-, and -S-(C₁-C₁₀)-alkyl-O-, O-CO-, -NR²²-(C₁-C₆)-alkyl-O-,

R²² is selected from the group of -H, -(C₁-C₁₀)-alkyl, a protecting group and a reporter group,

25 R¹⁴ is selected from the group consisting of -H, -OH, -(C₁-C₁₀)-alkoxy, -(C₂-C₁₀)-alkenyloxy, -(C₂-C₁₀)-alkynyloxy, -halogen, -azido, SH, -(C₁-C₁₀)-alkylmercapto, O-reporter group, O-solid phase and -NH₂ linked to a protecting group,


- 160 -

R^{15} is selected from the group consisting of -H, -(C₁-C₆)-alkyl, -(C₂-C₁₀)-alkenyl, -(C₂-C₁₀)-alkynyl, -(C₂-C₁₀)-alkyl-carbonyl, -(C₃-C₁₉)-alkenyl-carbonyl, -(C₃-C₁₉)-alkynyl-carbonyl, -(C₆-C₁₄)-aryl-(C₁-C₁₀)-alkyl and a solid phase,

B is the link to a moiety of formula I,

5

Formula I




wherein

R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of substituents (1) to (10) for R^1 in formula I in claim 1 and

10

(11)



15

wherein Nuc is the link to formula I and n is any integer from 1 to 18
said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being
unsubstituted or substituted by one or more moieties selected from the group
consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -
COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r
and s are independently of each other an integer of from 1 to 18 and n is 0 or 1
independently from r and s,

20

- 161 -

with the proviso that R^5 , R^6 or R^{12} is not a reporter group if the radicals (5) to (7) are substituted by $-NR^5R^6$, NHR^{12} , OR^{12} , or SR^{12} ;

R^2 , R^3 is independent from X, R^1 , R^2 and R^3 and is selected from the group of substituents (1) to (7) for R^2 and R^3 in formula I in claim 1,


5 X is independent from R^1 , R^2 or R^3 and is selected from the group consisting of N and CH

D is the position of attachment of the group to the rest of the nucleic acid binding compound.

10 wherein R^{14} is O-solid phase or R^{15} is solid phase,
with the proviso that one or two hydrogen atoms of any $-OH$, $-SH$, $-NH_2$, $-NH$ -alkyl, $-NH$ -alkenylene, $-NH$ -alkynylene, or $-NH$ -aryl group are substituted by a protecting group,
and any salts thereof.

15 29. A precursor or intermediate compound comprising a backbone, said backbone having attached heterocyclic groups characterized in that a heterocyclic group is a group of the general formula I

Formula I




20 wherein

R^1 is independent from X, R^2 or R^3 and is selected from the group consisting of substituents (1) to (10) for R^1 in formula I in claim 1 and

(11)

- 162 -



wherein Nuc is the position of attachment of the group to the rest of the group of formula I and n is any integer from 1 to 18

said alkyl, alkenyl, alkynyl or aryl in substituents (5) to (10) being

5 unsubstituted or substituted by one or more moieties selected from the group consisting of -halogen, -SH, -S-(C₁-C₆)-alkyl, -(C₁-C₆)-alkoxy, -OH, -NR⁵R⁶, -COR¹¹, -NH-CONR⁵R⁶, -NH-CSNR⁵R⁶ and -(CH₂)_n-[O-(CH₂)_r]_s-NR⁵R⁶, r and s are independently of each other an integer of from 1 to 18 and n is 0 or 1 independently from r and s,

10 with the proviso that R⁵, R⁶ or R¹² is not a reporter group if the radicals (5) to (7) are substituted by -NR⁵R⁶, NHR¹², OR¹², or SR¹²;

R², R³ is independent from X, R¹, R² and R³ and is selected from the group of substituents (1) to (7) for R² and R³ in formula I in claim 1,

15 X is independent from R¹, R² or R³ and is selected from the group consisting of N and CH

D is the position of attachment of the group to the rest of the nucleic acid binding compound.

wherein said backbone is solid phase bound,
with the proviso that one or two hydrogen atoms of any -OH, -SH, -NH₂, -NH-alkyl,
-NH-alkenylene, -NH-alkynylene, or -NH-aryl group are substituted by a protecting
group,

5 and any salts thereof.

30. A method for the enzymatic synthesis of a nucleic acid binding compound according to any of the claims 1 to 14, comprising reacting a triphosphate subunit with a primer using a nucleic acid as a template for the elongation of the primer, wherein the triphosphate subunit is a compound according to claim 21 or 22.
- 10 31. A method for the chemical synthesis of a compound according to any of the claims 1 to 14 using activated subunits, wherein said subunits contain a compound according to any of the claims 23 to 29.
32. A method according to claim 31 comprising the additional step of reacting the synthesized nucleic acid binding compound with a further compound.
- 15 33. A method according to claim 32 wherein the further compound is a label.
34. Use of a first nucleic acid binding compound in a hybridization reaction to form a parallel or antiparallel duplex with a second nucleic acid binding compound wherein the first nucleic acid binding compound and/ or the second nucleic acid binding compound comprise a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof.
- 20 35. Use according to claim 34 wherein the hybridization reaction is a multiplex hybridization reaction.
36. Use of a nucleic acid binding compound having a backbone as a capture probe, whereby the backbone has attached thereto heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof.
- 25 37. Use according to any of the claims 34 to 36 wherein the nucleic acid binding compound or the first and/ or the second nucleic acid binding compound is a nucleic acid binding compound according to any of the claims 1 to 14.
- 30 38. Use of a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof in place of a heterocyclic group in a first nucleic acid binding compound to increase the melting temperature of a parallel or antiparallel duplex with a second nucleic acid binding compound whereby the increase in melting temperature is compared to the melting temperature of a duplex of the first nucleic acid binding compound with a second nucleic acid binding compound wherein the heterocyclic group in the first nucleic
- 35

acid binding compound is complementary to a heterocyclic group in the second nucleic acid binding compound.

39. Use of a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof in place of a heterocyclic group in a first nucleic acid binding compound used as probe in an amplification reaction, to increase the melting temperature of a duplex with a second nucleic acid binding compound in comparison to the melting temperature of a primer used in the amplification reaction, whereby the increase in melting temperature is compared to the melting temperature of a duplex of the first nucleic acid binding compound with the second nucleic acid binding compound wherein the heterocyclic group in the first nucleic acid binding compound is complementary to a heterocyclic group in the second nucleic acid binding compound.
40. Use of a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof in place of a heterocyclic group in a first nucleic acid binding compound to harmonize the contribution of each base pair to the melting temperature of a parallel or antiparallel duplex with a second nucleic acid binding compound.
41. Use of a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof in place of a heterocyclic group in a first nucleic acid binding compound for enhanced detection of sequences in a second nucleic acid binding compound having mismatches in a duplex with the first nucleic acid binding compound.
42. Use of a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof in place of a heterocyclic group in a nucleic acid binding compound to increase the melting temperature of an intramolecular duplex of the nucleic acid binding compound whereby the increase in melting temperature is compared to the melting temperature of the intramolecular duplex of the nucleic acid binding compound wherein the heterocyclic group in the nucleic acid binding compound is complementary to a heterocyclic group in the hybridizing part of the nucleic acid binding compound.
43. Use of a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof in place of a heterocyclic group in a nucleic acid binding compound for enhanced detection of subtypes in a target nucleic acid.
44. Use according to any of the claims 39 to 43 wherein one or two reporter groups are attached to the nucleic acid binding compound or the first and/or the second nucleic acid binding compound.
45. Use according to claim 44 wherein the substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof in place of a heterocyclic group is 1 to 5 nucleotides separated from the point of attachment of a reporter group.

- 165 -

46. Use according to any of the claims 34 to 45 wherein the substituted pyrazolo[3,4-d]pyrimidine analogue is a substituted pyrazolo[3,4-d]pyrimidine analogue of adenine or guanine or a 7-substituted pyrazolo[3,4-d]pyrimidine analogue of adenine or guanine.
- 5 47. Use according to any of the claims 34 to 45 wherein the substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof is a substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof or a 7-substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof.
48. Use according to any of the claims 47 wherein the substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof or the 7-substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof has the formula I with the substituents as defined in claim 1.
- 10 49. A method for the determination of the presence, absence or amount of a nucleic acid comprising the steps
- 15 - providing a sample suspected to contain the nucleic acid,
- providing a nucleic acid binding compound comprising a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof, which is essentially complementary to a part or all of the
- 20 nucleic acid,
- contacting said sample with the nucleic acid binding compound under conditions for binding the nucleic acid binding compound to the nucleic acid to form a duplex,
- determining the binding product or the degree of hybridization between the nucleic acid and the nucleic acid binding compound as a measure of the presence, absence or
- 25 amount of the nucleic acid.
50. A method for distinguishing related nucleotide sequences in a nucleic acid, the method comprising the following steps:
- a) providing a nucleic acid binding compound having a defined sequence, wherein the nucleic acid binding compound comprises a backbone, said backbone having
- 30 attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof;
- b) providing a nucleic acid with two related nucleotide sequences, each of which comprises a target sequence, wherein one of the nucleotide sequence is a target
- 35 sequence that is perfectly complementary to the nucleic acid binding compound and one other of the segments is a related target sequence;

- 166 -

- c) incubating the nucleic acid with the nucleic acid binding compound under hybridization conditions to form a duplex; and
- d) determining the degree of hybridization between the nucleic acid binding compound and each of the segments.
- 5 51. The method according to any of the claims 49 to 50, wherein the nucleic acid binding compound comprises a reporter group.
52. The method according to any of the claims 51 wherein the reporter group is a fluorescent label.
53. The method according to claim 52 wherein the nucleic acid binding compound
10 further comprises a quenching agent which quenches the fluorescence emission of the fluorescent label.
54. The method according to claim 53 wherein the fluorescent label is a fluorescein and wherein the quenching agent is a fluorescent rhodamine or cyanine.
- 15 55. The method according to any of the claims 53 to 54, further comprising the step of altering the spatial relationship between the fluorescent label and the quenching agent subsequent to hybridization.
56. The method according to claim 55, wherein alteration of the spatial relationship between the fluorescent label and the quenching agent is accomplished by
20 exonuclease hydrolysis of the nucleic acid binding compound, wherein release of label occurs as a result of exonuclease hydrolysis.
57. The method according to claim 56, wherein the degree of hybridization between the nucleic acid binding compound and the nucleic acid is determined by the quantity of label that is released from the nucleic acid binding compound subsequent to
25 hybridization.
58. The method according to claim 49 to 50, wherein the degree of hybridization between the nucleic acid binding compound and the nucleic acid is determined by the priming ability of the nucleic acid binding compound.
59. The method according to claim 58, wherein priming occurs as part of a polymerase
30 chain reaction.
60. The method according to claim 52, wherein more than one nucleic acid binding compound is used.
61. The method according to claim 60, wherein two nucleic acid binding compounds are used.
- 35 62. The method according to claim 61, wherein the first of the two nucleic acid binding compounds comprises a fluorescence donor and the second of the two nucleic acid


- binding compounds comprises a fluorescence acceptor, and wherein the emission wavelengths of the fluorescence donor overlap the absorption wavelengths of the fluorescence acceptor.
63. The method according to claim 62 wherein the degree of hybridization is measured
5 by the quantity of light transferred between the fluorescence donor and the fluorescence acceptor and emitted by the fluorescence acceptor.
64. The method according to claim 62 wherein the degree of hybridization is determined by the measurement of the melting temperature between the nucleic acid binding compound and the nucleic acid.
- 10 65. A method for detecting the presence of a target sequence in a nucleic acid, the method comprising the following steps:
- a) providing a nucleic acid which is to be tested for the presence of the target sequence;
 - b) providing a nucleic acid binding compound comprising a backbone, said
15 backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof, wherein the nucleic acid binding compound has a sequence that is substantially complementary to the target sequence;
 - c) incubating the nucleic acid and the nucleic acid binding compound under
20 hybridization conditions; and
 - d) identifying hybridized nucleic acids;
66. The method according to claim 65, wherein multiple nucleic acids are tested for the presence of the target sequence, and wherein the nucleic acids have related target sequences.
- 25 67. The method according to claim 65, wherein the nucleic acid binding compound is a primer comprising an extendible 3'-hydroxyl group.
68. The method according to claim 67, wherein the hybridized nucleic acids are identified by extending the primer with a polymerizing enzyme.
69. The method according to claim 67, wherein the nucleic acid binding compound is a
30 primer in a polymerase chain reaction.
70. A method for primer extension, the method comprising the following steps:
- a) providing a nucleic acid containing a target sequence,
 - b) providing one or more nucleic acid binding compound complementary to the
target sequence wherein the nucleic acid binding compound comprises a backbone,
35 said backbone having attached heterocyclic groups capable of base pairing to

- nucleobases characterized in that a heterocyclic groups is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof,
- c) providing a polymerizing enzyme and nucleotide substrates, and
- d) incubating the nucleic acid, the nucleic acid binding compounds, the enzyme and the substrates under conditions favorable for polymerization.
71. The method according to claim 70, wherein the method is part of a polymerase chain reaction.
72. The method according to claim 70, wherein the method is used in the synthesis of a cDNA molecule.
73. A method for determining the nucleotide sequence of a nucleic acid, the method comprising the following steps:
- a) providing an array of nucleic acid binding compounds having different known sequences, wherein the nucleic acid binding compound comprises a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof, with the proviso that the nucleic acid binding compounds do not contain a reporter group;
- b) incubating the nucleic acid with the array under hybridization conditions, and
- c) determining to which of the nucleic acid binding compounds in the array the nucleic acid hybridizes.
74. A method for determining the nucleotide sequence of a target sequence in a nucleic acid, the method comprising the following steps:
- a) providing a nucleic acid comprising the target sequence;
- b) providing at least two nucleic acid binding compounds comprising a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic group is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof with a known sequence and wherein one of the at least two nucleic acid binding compounds has a sequence that is perfectly complementary to the target sequence and at least one other of the nucleic acid binding compounds has a related target sequence;
- c) incubating the at least two nucleic acid binding compounds with the nucleic acid under hybridization conditions; and
- d) determining the degree of hybridization between each of the nucleic acid binding compounds and the nucleic acid.
75. The method according to claim 74, wherein the at least one other nucleic acid binding compounds has a single-nucleotide mismatch with the target sequence.

76. A method for examining gene expression in a cell, the method comprising the following steps:
- a) providing a population of nucleic acids representative of the genes expressed in the cell,
 - 5 b) providing an array of nucleic acid binding compounds having different sequences wherein the nucleic acid binding compounds comprise a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic groups is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof, and with the proviso that the nucleic acid binding compounds do not contain a reporter group;
 - 10 c) incubating the population of nucleic acids with the array under hybridization conditions, and
 - d) determining which of the nucleic acid binding compounds in the array become hybridized to nucleic acids.
- 15 77. A method for identifying a mutation in a target sequence of a gene of interest, the method comprising the following steps:
- a) providing a nucleic acid that comprises the target sequence.
 - b) providing an array of nucleic acid binding compounds comprising a backbone, said backbone having attached heterocyclic groups capable of base pairing to nucleobases characterized in that a heterocyclic groups is a substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof, wherein the nucleic acid binding compounds have different sequences, wherein the different sequences include the wild-type target sequence and different mutant target sequences, with the proviso that the nucleic acid binding compounds do not contain a reporter group;
 - 20 c) incubating the nucleic acid with the array under hybridization conditions, and
 - 25 d) determining which of the nucleic acid binding compounds in the array become hybridized to the nucleic acid.
78. Method according to any of the claims 49 to 77 wherein the substituted pyrazolo[3,4-d]pyrimidine analogue is a substituted pyrazolo[3,4-d]pyrimidine analogue of adenine or guanine or a 7-substituted pyrazolo[3,4-d]pyrimidine analogue of adenine or guanine.
- 30 79. Method according to any of the claims 49 to 77 wherein the substituted pyrazolo[3,4-d]pyrimidine or an analogue thereof is a substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof or a 7-substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof.
- 35

- 170 -

80. Method according to claim 79 wherein the substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof or the 7-substituted 7-deaza-8-aza-2,6-diamino-purine or a derivative thereof has the general formula I as defined in claim 1.
81. Method according to any of the claims 49 to 80 wherein the nucleic acid binding compound is a nucleic acid binding compound according to any of the claims 1 to 14.
82. Method according to any of the claims 49 to 80 wherein a parallel or an antiparallel duplex is formed.
83. A pharmaceutical composition comprising a nucleic acid binding compound according to any of the claims 1 to 14.
84. A nucleic acid binding compound according to any of the claims 1 to 14 for use in medicine.
85. A building block for the synthesis of an oligonucleotide comprising the nucleosides adenosine or desoxyadenosine, guanosine or desoxyguanosine, isoguanosine or desoxyisoguanosine, cytidine or deoxycytidine, uridine or desoxyuridine, thymidine or desoxythymidine, or analogues of these nucleosides wherein a substituent is attached to the base moiety selected from the group of phthalimidoalkyl, phthalimidoalkenyl, or phthalimidoalkynyl groups with the following formulas



wherein Nuc is the position of attachment of the substituent to the base moiety and n is any integer from 1 to 18, with the proviso that the phthalimidoalkynyl or phthalimidoalkyl group is not attached to the C5-atom of deoxyuridine and the proviso that the


- 171 -

phthalimidoalkynyl group is not attached to the C7-atom of 7-deaza-deoxyguanosine.

86. A building block according to claim 85 wherein the analogue of these nucleosides is a pyrazolo[3,4-d]pyrimidine analogue.
- 5 87. A building block according to any of the claims 85 to 86 wherein the building block is a phosphoramidite derivative.
88. Use of a building block according to any of the claims 85 to 87 for the synthesis of an oligonucleotide.
89. Use of a phthaloyl group as a protecting group in a method for the synthesis of an
10 oligonucleotide from building blocks comprising the nucleosides adenosine or desoxyadenosine, guanosine or desoxyguanosine, isoguanosine or desoxyisoguanosine, cytidine or deoxycytidine, uridine or desoxyuridine, thymidine or desoxythymidine, or analogues of these nucleosides wherein the amino groups of $-(CH_2)_n-NH_2$, $-CH=CH-(CH_2)_n-NH_2$, or $-C\equiv C-$
15 $(CH_2)_n-NH_2$ attached to the base moiety are derivatized with the phthaloyl group and n is any integer from 1 to 18,
with the proviso that the $-(CH_2)_n-NH_2$ or $-C\equiv C-(CH_2)_n-NH_2$ group is not attached to the C5-atom of deoxyuridine and the proviso that the $-C\equiv C-(CH_2)_n-NH_2$ group is not attached to the C7-atom of 7-deaza-deoxyguanosine.
- 20 90. Use according to claim 89 wherein the analogue of these nucleosides is a pyrazolo[3,4-d]pyrimidine analogue.
91. Use according to any of the claims 88 to 90 wherein the building block is a phosphoramidite derivative.
92. A method for the synthesis of an oligonucleotide from building blocks according to
25 any of the claims 85 to 87.

Figure 1

a)



b)





Figure 2

a)



b)





Figure 3

a)



b)





Figure 4

a)



5

b)





Figure 5a,b



5 Figure 5c,d




Figure 6




Figure 7a





Figure 7b



- 1 -

SEQUENCE LISTING

5 <110> Roche Diagnostics GmbH
F. HOFFMANN-LA ROCHE AG

<120> Oligonucleotides containing Pyrazolo (3,4-d) pyrimidine
analogues of Purin-2,6-diamine and their uses

10 <130> 5419/0A/

<140>
<141>

15 <150> EP00116816.0
<151> 2000-08-03

<160> 39

20 <170> PatentIn Ver. 2.1

<210> 1
<211> 12
<212> DNA

25 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
(100)

30 <220>
<223> n in positions 4 and 8 denotes compound 2c

<400> 1

35 agtnttgncc ta 12

<210> 2
<211> 12
<212> DNA

40 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
(102)

45 <400> 2
taggtcaata ct 12

50 <210> 3
<211> 12
<212> DNA

<213> Artificial Sequence

55 <220>
<223> Description of Artificial Sequence:oligonucleotide
(103)

60 <400> 3
agtattgacc ta 12

- 2 -

<210> 4
<211> 12
5 <212> DNA
<213> Artificial Sequence

<220>
10 <223> Description of Artificial Sequence:oligonucleotide
(104)

<220>
<223> n in positions 2, 7, 8 and 10 denotes compound 1

15 <400> 4
tnggtcnntn ct 12

<210> 5
20 <211> 12
<212> DNA
<213> Artificial Sequence

<220>
25 <223> Description of Artificial Sequence:oligonucleotide
(105)

<220>
30 <223> n in positions 4 and 8 denotes compound 1

<400> 5
agtnttgncc ta 12

35 <210> 6
<211> 12
<212> DNA
<213> Artificial Sequence

40 <220>
<223> Description of Artificial Sequence:oligonucleotide
(106)

<400> 6
45 taggccggca ct 12

<210> 7
<211> 12
50 <212> DNA
<213> Artificial Sequence

<220>
55 <223> Description of Artificial Sequence:oligonucleotide
(107)

<400> 7
agtgccggcc ta 12

60 <210> 8
<211> 12

- 3 -

<212> DNA
<213> Artificial Sequence

<220>
5 <223> Description of Artificial Sequence:oligonucleotide
(108)

<220>
10 <223> n in positions 7 and 8 denotes compound 2a

<400> 8
taggtcnnta ct 12

<210> 9
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
20 <223> Description of Artificial Sequence:oligonucleotide
(109)

<220>
25 <223> n in positions 4 and 8 denotes compound 2a

<400> 9
agtnttgncc ta 12

<210> 10
<211> 12
<212> DNA
<213> Artificial Sequence

<220>
35 <223> Description of Artificial Sequence:oligonucleotide
(110)

<220>
40 <223> n in positions 7 and 8 denotes compound 2b

<400> 10
taggtcnnta ct 12

<210> 11
<211> 12
<212> DNA
50 <213> Artificial Sequence

<220>
55 <223> Description of Artificial Sequence:oligonucleotide
(111)

<220>
60 <223> n in positions 4 and 8 denotes compound 2b

<400> 11
agtnttgncc ta 12

- 4 -

<210> 12
<211> 12
<212> RNA
<213> Artificial Sequence
5
<220>
<223> Description of Artificial Sequence:oligonucleotide
(114)
10 <400> 12
aguauugacc ua 12

<210> 13
15 <211> 12
<212> DNA
<213> Artificial Sequence
<220>
20 <223> Description of Artificial Sequence:oligonucleotide
(115)
<220>
<223> n in following positions denotes: 3 + 4 = isoC, 6
25 + 11 = isoG
<400> 13
atnnanttat na 12

30 <210> 14
<211> 12
<212> DNA
<213> Artificial Sequence
35 <220>
<223> Description of Artificial Sequence:oligonucleotide
(116)
40 <220>
<223> n in following positions denotes: 2 + 7 = isoC, 9
+ 10 = isoG
<400> 14
45 tnataantnn at 12

<210> 15
<211> 12
50 <212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence:oligonucleotide
55 (200)
<220>
<223> n in position 7 denotes compound 2b
60 <400> 15
taggtcnata ct 12

- 5 -

<210> 16
<211> 12
<212> DNA
5 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
(201)
10

<220>
<223> n in positions 2, 7, 8 and 10 denotes compound 2b

<400> 16
15 tnggtcnntn ct 12

<210> 17
<211> 12
20 <212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
25 (202)

<220>
<223> n in position 8 denotes compound 2c

<400> 17
30 agtattgncc ta 12

<210> 18
35 <211> 12
<212> DNA
<213> Artificial Sequence

<220>
40 <223> Description of Artificial Sequence:oligonucleotide
(203)

<220>
<223> n in position 4 denotes compound 2c
45

<400> 18
agtnttgacc ta 12

<210> 19
50 <211> 12
<212> DNA
<213> Artificial Sequence

<220>
55 <223> Description of Artificial Sequence:oligonucleotide
(204)

<220>
60 <223> n in positions 7 and 8 denotes compound 2c

<400> 19

- 6 -

taggtcnnta ct 12

5 <210> 20
<211> 12
<212> DNA
<213> Artificial Sequence

10 <220>
<223> Description of Artificial Sequence:oligonucleotide
(205)

15 <220>
<223> n in positions 2, 7, 8 and 10 denotes compound 2c
<400> 20
tnggtcnntn ct 12

20 <210> 21
<211> 12
<212> DNA
<213> Artificial Sequence

25 <220>
<223> Description of Artificial Sequence:oligonucleotide
(300)

30 <220>
<223> n in positions 5 and 9 denotes compound 3a
<400> 21
taggncaana ct 12

35 <210> 22
<211> 12
<212> DNA
<213> Artificial Sequence

40 <220>
<223> Description of Artificial Sequence:oligonucleotide
(301)

45 <220>
<223> n in positions 5 and 6 denotes compound 3a
<400> 22
agtanngacc ta 12

50 <210> 23
<211> 12
<212> DNA
<213> Artificial Sequence

55 <220>
<223> Description of Artificial Sequence:oligonucleotide
(302)

60 <220>
<223> n in positions 5 and 9 denotes compound 3b

- 7 -

5 <400> 23
taggncaana ct 12

5 <210> 24
 <211> 12
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <223> Description of Artificial Sequence:oligonucleotide
 (303)

15 <220>
 <223> n in positions 5 and 6 denotes compound 3b

20 <400> 24
agtanngacc ta 12

25 <210> 25
 <211> 12
 <212> DNA
 <213> Artificial Sequence

30 <220>
 <223> Description of Artificial Sequence:oligonucleotide
 (304)

35 <400> 25
taggncaana ct 12

40 <210> 26
 <211> 12
 <212> DNA
 <213> Artificial Sequence

45 <220>
 <223> Description of Artificial Sequence:oligonucleotide
 (305)

50 <220>
 <223> n in positions 5 and 6 denotes compound 3c

50 <400> 26
agtanngacc ta 12

55 <210> 27
 <211> 12
 <212> DNA
 <213> Artificial Sequence

60 <220>
 <223> Description of Artificial Sequence:oligonucleotide
 (400)

- 8 -

<400> 27
taggacaata ct 12

5 <210> 28
<211> 12
<212> DNA
<213> Artificial Sequence

10 <220>
<223> Description of Artificial Sequence:oligonucleotide
(401)

<400> 28
15 tagggcaata ct 12

<210> 29
<211> 12
20 <212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
25 (402)

<400> 29
taggcgaata ct 12

30 <210> 30
<211> 12
<212> RNA
<213> Artificial Sequence

35 <220>
<223> Description of Artificial Sequence:oligonucleotide
(403)

40 <400> 30
aguauugacc ua 12

<210> 31
45 <211> 13
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
50 (500)

<220>
<223> n in position 1 denotes compound 24
55

<400> 31
ntaggtcaat act 13

60 <210> 32
<211> 13
<212> DNA

- 9 -

<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
5 (501)

<220>
<223> n in position 1 denotes compound 24 and n in
10 positions 8 and 9 denotes compound 2b

<400> 32
ntaggtcnnt act 13

15 <210> 33
<211> 13
<212> DNA
<213> Artificial Sequence

20 <220>
<223> Description of Artificial Sequence:oligonucleotide
(502)

<220>
25 <223> n in position 1 denotes compound 25

<400> 33
nagtattgac cta 13

30 <210> 34
<211> 13
<212> DNA
<213> Artificial Sequence

35 <220>
<223> Description of Artificial Sequence:oligonucleotide
(503)

40 <220>
<223> n in position 1 denotes compound 26

<400> 34
nagtattgac cta 13

45 <210> 35
<211> 12
<212> DNA
50 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
(504)

55 <220>
<223> n in positions 2, 7, 9 and 10 denotes compound 27

<400> 35
60 tnataantnn ta 12

- 10 -

<210> 36
<211> 12
<212> DNA
<213> Artificial Sequence
5
<220>
<223> Description of Artificial Sequence:oligonucleotide
(505)
10 <400> 36
agttattgagg at 12

<210> 37
15 <211> 13
<212> DNA
<213> Artificial Sequence

<220>
20 <223> Description of Artificial Sequence:oligonucleotide
(506)

<220>
25 <223> n in positions 3, 8, 10 and 11 denotes compound 27
and n in position 1 denotes compound 24

<400> 37
ntnataantn nta 13
30
<210> 38
<211> 13
<212> DNA
<213> Artificial Sequence
35
<220>
<223> Description of Artificial Sequence:oligonucleotide
(507)
40
<220>
<223> n in position 13 denotes compound 26

<400> 38
45 agttattgagg atn 13

<210> 39
<211> 13
<212> DNA
50 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
(508)
55
<220>
<223> n in positions 3, 8, 10 and 11 denotes compound
27, n in positions 6 and 7 denotes compound 2b and
n in position 1 denotes compound 24
60
<400> 39
ntnatnnntn nta 13

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/08850

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07H19/16 C07H21/00 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07H C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	F. SEELA ET AL.: "8-Aza-7-deazapurine DNA: Synthesis and duplex stability of oligonucleotides containing 7-substituted bases" NUCLEOSIDES & NUCLEOTIDES, vol. 18, 1999, pages 1399-1400, XP002157790 cited in the application the whole document	1,3-11, 14,20, 23,24, 26-29, 31,34-48
A	WO 90 03370 A (MICROPROBE CORP) 5 April 1990 (1990-04-05) cited in the application the whole document	1,3-92
A	WO 99 51775 A (EPOCH PHARMACEUTICALS INC) 14 October 1999 (1999-10-14) the whole document	1,3-92
	--- -/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

6 November 2001

Date of mailing of the international search report

14/11/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

de Nooy, A

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 01/08850

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>OERTEL, FRANK ET AL: "Synthesis and properties of methylthiopyrazolo'3,4-d!pyrimidine 2'-deoxy-.beta.-D-ribonucleosides" LIEBIGS ANN. CHEM. (1992), (11), 1165-70, XP002157791 cited in the application page 1165, structure 3b</p>	1
A	<p>GARAEVA, L. D. ET AL: "Studies on glycosides of 3,4,6-trisubstituted pyrazolo'3,4-d!pyrimidines. Synthesis of 2'-deoxyribonucleosides" NUCLEOSIDES NUCLEOTIDES (1991), 10(6), 1295-303, XP000980295 cited in the application page 1297, structure 8</p>	1
A	<p>PETRIE C R ET AL: "A NOVEL BIOTINYLATED ADENYLATE ANALOGUE DERIVED FROM PYRAZOLOU3,4-DPYRIMIDINE FOR LABELING DNA PROBES" BIOCONJUGATE CHEMISTRY, US, AMERICAN CHEMICAL SOCIETY, WASHINGTON, vol. 2, no. 6, 1 November 1991 (1991-11-01), pages 441-446, XP000572789 ISSN: 1043-1802 abstract page 442, structure 9</p>	1
A	<p>F. SEELA ET AL.: "Duplex stability of oligonucleotides containing 7-substituted 7-deaza- and 8-aza-7-deazapurine nucleosides" NUCLEOSIDES & NUCLEOTIDES, vol. 16, 1997, pages 963-966, XP002157792 cited in the application abstract page 963-964, structure 1-11</p>	1
A	<p>G. BALOW ET AL.: "Biophysical and antisense properties of oligodeoxynucleotides containing 7-propenyl-, 7-iodo- and 7-cyano-7-deaza-2-amino-2'-deoxyadenosines " NUCLEIC ACIDS RESEARCH, vol. 26, no. 14, 1998, pages 3350-3357, XP002157793 cited in the application page 3351-3353, structures 1-21</p> <p style="text-align: center;">-/--</p>	1

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/08850

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	N. RAMZAEVA ET AL.: "7-Deaza-2'-deoxyguanosines functionalized with 7-(omega-aminoalk-1-ynyl) residues" NUCLEOSIDES & NUCLEOTIDES, vol. 18, 1999, pages 1439-1440, XP002157794 cited in the application the whole document -----	1,23
A	DE 198 23 454 A (EPIGENOMICS GMBH) 25 November 1999 (1999-11-25) abstract -----	15

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1 (in part), 2 (not searched), 3-13 (in part), 15-21 (in part), 23-92 (in part)

Present claims 1-13, 15-21, 23-92 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds, precursors, intermediates, nucleic acid binding compounds, building blocks containing at least one heterocyclic group which is a 7-halogen-7-deaza-8-aza-2,6-diamino-purine (that is, the compounds of Formula I of claim 1 where R1 is halogen (-F, -Cl, -Br, -I)).

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/08850

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9003370	A	05-04-1990	CA 1338379 A1	11-06-1996
			WO 9003370 A1	05-04-1990
			US 5824796 A	20-10-1998
			US 5849482 A	15-12-1998
WO 9951775	A	14-10-1999	US 6127121 A	03-10-2000
			AU 3472499 A	25-10-1999
			EP 1068358 A1	17-01-2001
			WO 9951775 A1	14-10-1999
DE 19823454	A	25-11-1999	DE 19823454 A1	25-11-1999
			AU 4896899 A	06-12-1999
			WO 9960156 A2	25-11-1999